

Noxious Weed Qualitative Risk Assessment

Oregon Department of Agriculture

Common name: North Africa grass, Ventenata grass

Scientific name: *Ventenata dubia*

Family: *Poaceae*

For use with plant species that occur or may occur in Oregon to determine their potential to become serious noxious weeds. For each of the following categories, select the number that best applies. Numerical values are weighted to increase priority categories over less important ones. Choose the best number that applies, intermediate scores can be used.

Total Score: 53 Risk Category: B

GEOGRAPHICAL INFORMATION

1) 6 Invasive in Other Areas

- 0 Low- not know to be invasive elsewhere.
- 2 Known to be invasive in climates dissimilar to Oregon's current climates.
- 6 Known to be invasive in geographically similar areas.

Comments: Reported from other temperate zone states in the northwest and northeast.

2) 6 Habitat Availability: Are there susceptible habitats for this species and how common or widespread are they in Oregon?

- 1 *Low* – Habitat is very limited, usually restricted to a small watershed or part of a watershed (e.g., tree fern in southern Curry County).
- 3 *Medium* – Habitat encompasses 1/4 or less of Oregon (e.g., oak woodlands, coastal dunes, eastern Oregon wetlands, Columbia Gorge).
- 6 *High* – Habitat covers large regions or multiple counties, or is limited to a few locations of high economic or ecological value (e.g., threatened and endangered species habitat).

Comments: Habitat covers much of eastern and western Oregon.

3) 0 Proximity to Oregon: What is the current distribution of the species?

- 0 *Present* – Occurs within Oregon.
- 1 *Distant* – Occurs only in distant US regions or foreign countries.
- 3 *Regional* – Occurs in Western regions of US but not adjacent to Oregon border.
- 6 *Adjacent* – Weedy populations occur adjacent (<50 miles) to Oregon border.

Comments: Occurs throughout Oregon, primarily east of the Cascades.

4) 3 Current Distribution: What is the current distribution of escaped populations in Oregon?

- 0 *Not present* – Not known to occur in Oregon.
- 1 *Widespread* – Throughout much of Oregon (e.g., cheatgrass).
- 5 *Regional* – Abundant (i.e., occurs in eastern, western, central, coastal, areas of Oregon) (e.g., gorse, tansy ragwort).
- 10 *Limited* – Limited to one or a few infestations in state (e.g., kudzu).

Comments: Abundant in eastern Oregon and has a lesser distribution through western Oregon.

BIOLOGICAL INFORMATION

- 5) 3 **Environmental Factors:** Do abiotic (non-living) factors in the environment effect establishment and spread of the species? (e.g., precipitation, drought, temperature, nutrient availability, soil type, slope, aspect, soil moisture, standing or moving water).
- 1 *Low* – Severely confined by abiotic factors.
 - 2 *Medium* – Moderately confined by environmental factors
 - 4 *High* – Highly adapted to a variety of environmental conditions (e.g., tansy ragwort, Scotch broom).

Comments: Although establishes in moist areas of clayey soil types, recent history has shown this plant to expand to drier sites

- 6) 3 **Reproductive Traits:** How does this species reproduce? Traits that may allow rapid population increase both on and off site.
- 0 *Negligible* – Not self-fertile, or is dioecious and opposite sex not present.
 - 1 *Low* – Reproduction is only by seed, produces few seeds, or seed viability and longevity are low.
 - 3 *Medium* – Reproduction is vegetative (e.g., by root fragments, rhizomes, bulbs, stolons).
 - 3 *Medium* – Produces many seeds, and/or seeds of short longevity (< 5 years).
 - 5 *High* – Produces many seeds and/or seeds of moderate longevity (5-10 years) (e.g., tansy ragwort).
 - 6 *Very high* – Has two or more reproductive traits (e.g., seeds are long-lived >10 years and spreads by rhizomes).

Comments: Plant reproduces via prolific seeding.

- 7) 4 **Biological Factors:** Do biotic (living) factors restrict or aid establishment and spread of the species? (What is the interaction of plant competition, natural enemies, native herbivores, pollinators, and pathogens with species?)
- 0 *Negligible* – Host plant not present for parasitic species.
 - 1 *Low* – Biotic factors highly suppress reproduction or heavily damage plant for an extended period (e.g., biocontrol agent on tansy ragwort).
 - 2 *Medium* – Biotic factors partially restrict or moderately impact growth and reproduction, impacts sporadic or short-lived.
 - 4 *High* – Few biotic interactions restrict growth and reproduction. Species expresses full growth and reproductive potential.

Comments: Herbivory minimal, highly competitive with perennial grasses, no biocontrols present.

- 8) 3 **Reproductive Potential and Spread After Establishment - Non-human Factors:** How well can the species spread by natural means?
- 0 *Negligible* – No potential for natural spread in Oregon (e.g., ornamental plants outside of climate zone).
 - 1 *Low* – Low potential for local spread within a year, has moderate reproductive potential or some mobility of propagules (e.g., propagules transported locally by animals, water movement in lakes or ponds, not wind blown).
 - 3 *Medium* - Moderate potential for natural spread with either high reproductive potential or highly mobile propagules (e.g., propagules spread by moving water, or dispersed over longer distances by animals) (e.g., perennial pepperweed).
 - 5 *High* – Potential for rapid natural spread throughout the susceptible range, high reproductive capacity and highly mobile propagules. Seeds are wind dispersed over large areas (e.g., rush skeletonweed).

Comments:

- 9) 4 **Potential of Species to be Spread by Humans.** What human activities contribute to spread of species? Examples include: interstate or international commerce; contaminated commodities; packing materials or products; vehicles, boats, or equipment movement; logging or farming; road maintenance; intentional introductions of ornamental and horticultural species, or biofuel production.
- 1 *Low* – Potential for introduction or movement minimal (e.g., species not traded or sold, or species not found in agricultural commodities, gravel or other commercial products).
 - 3 *Medium* – Potential for introduction or off-site movement moderate (e.g., not widely propagated, not highly popular, with limited market potential; may be a localized contaminant of gravel, landscape products, or other commercial products) (e.g., lesser celandine, Canada thistle).
 - 5 *High* – Potential to be introduced or moved within state high (e.g., species widely propagated and sold; propagules common contaminant of agricultural commodities or commercial products; high potential for movement by contaminated vehicles and equipment, or by recreational activities) (e.g., butterfly bush, spotted knapweed, Eurasian watermilfoil).

Comments: Hay and crop contaminate, fine structure and seed adhere easily as the plant occurs commonly in clayey soils.

IMPACT INFORMATION

- 10) 5 **Economic Impact:** What impact does/can the species have on Oregon's agriculture and economy?
- 0 *Negligible* – Causes few, if any, economic impacts.
 - 1 *Low* - Potential to, or causes low economic impact to agriculture; may impact urban areas (e.g., puncture vine, pokeweed).
 - 5 *Medium* – Potential to, or causes moderate impacts to urban areas, right-of-way maintenance, property values, recreational activities, reduces rangeland productivity (e.g., English ivy, Himalayan blackberry, cheatgrass).
 - 10 *High* – Potential to, or causes high impacts in agricultural, livestock, fisheries, or timber production by reducing yield, commodity value, or increasing production costs (e.g., gorse, rush skeleton weed, leafy spurge).

Comments: Similar impacts to that of cheatgrass.

- 11) 5 **Environmental Impact:** What risks or harm to the environment does this species pose? Plant may cause negative impacts on ecosystem function, structure, and biodiversity of plant or fish and wildlife habitat; may put desired species at risk.
- 0 *Negligible* – None of the above impacts probable.
 - 1 *Low* – Can or does cause few or minor environmental impacts, or impacts occur in degraded or highly disturbed habitats.
 - 4 *Medium* – Species can or does cause moderate impacts in less critical habitats (e.g., urban areas, sagebrush/ juniper stands).
 - 6 *High* – Species can or does cause significant impacts in several of the above categories. Plant causes severe impacts to limited or priority habitats (e.g., aquatic, riparian zones, salt marsh; or T&E species sites).

Comments: Causes reductions in species richness where infestations are large and dense, invasive to T&E habitats throughout the upper elevations of Blue Mountains and alters fire regimes.

- 12) 2 **Impact on Health:** What is the impact of this species on human, animal, and livestock health? (e.g., poisonous if ingested, contact dermatitis, acute and chronic toxicity to livestock, toxic sap, injurious spines or prickles, causes allergy symptoms.
- 0 *Negligible* – Has no impact on human or animal health.
 - 2 *Low* – May cause minor health problems of short duration, minor allergy symptoms (e.g., leafy spurge).
 - 4 *Medium* – May cause severe allergy problems, death or severe health problems through chronic toxicity, spines or toxic sap may cause significant injury. (e.g., giant hogweed, tansy ragwort).
 - 6 *High* – Causes death from ingestion of small amounts, acute toxicity (e.g. poison hemlock).

Comments: Recognized as a significant factor in changing fire regimes and increased danger in fire suppression activities.

CONTROL INFORMATION

- 13) 5 **Probability of Detection at Point of Introduction:** How likely is detection of species after introduction and naturalization in Oregon?
- 1 *Low* – Grows where probability of early detection is high, showy and easily recognized by public; access to habitat not restricted (e.g., giant hogweed).
 - 5 *Medium* – Easily identified by weed professionals, ranchers, botanists; some survey and detection infrastructure in place. General public may not recognize or report species (e.g., leafy spurge).
 - 10 *High* – Probability of initial detection by weed professionals low. Plant shape and form obscure, not showy for much of growing season, introduction probable at remote locations with limited access (e.g., weedy grasses, hawkweeds, skeletonweed).

Comments: Diffuse, fine structured plant makes it less conspicuous, may be difficult to differentiate from a similar native, the public may not notice the spread of the species early on.

- 14) 4 **Control Efficacy:** What level of control of this species can be expected with proper timing, herbicides, equipment, and biological control agents?
- 1 *Negligible* – Easily controlled by common non-chemical control measures (e.g., mowing, tillage, pulling, and cutting; biocontrol is very effective at reducing seed production and plant density) (e.g., tansy ragwort).
 - 2 *Low* – Somewhat difficult to control, generally requires herbicide treatment (e.g., mechanical control measures effective at preventing flowering and but not reducing plant density; herbicide applications provide a high rate of control in a single application; biocontrol provides partial control).
 - 4 *Medium* – Treatment options marginally effective or costly. Tillage and mowing increase plant density (e.g., causes tillering, rapid regrowth, spread from root fragments). Chemical control is marginally effective. Crop damage occurs or significant non-target impacts result from maximum control rates. Biocontrol agents ineffective.
 - 6 *High* – No effective treatments known or control costs very expensive. Species may occur in large water bodies or river systems where containment and complete control are not achievable. Political or legal issues may prevent effective control.

Comments: Herbicide effectiveness variable, no biocontrol.

Category Scores:

15 Geographic score (Add scores 1-4)

17 Biological Score (Add lines 5-9)

12 Impact Score (Add lines 10-12)

09 Control Score (Add Lines 13-14)

53 Total Score (Add scores 1-14 and list on front of form)

Risk Category: 55-89+ = **A** 24-54 = **B** < 24 = unlisted.

This Risk Assessment was modified by ODA from the USDA-APHIS Risk Assessment for the introduction of new plant species.

Vers. 3.8 1/31/2019