

**Vaccines for Children
Oregon State-Supplied Vaccine
Effective March 25, 2009**

Vaccines	Ages of Covered Children	All or High Risk?
DTaP	6 weeks through 6 years	All
Hepatitis A	1 year through 18 years	All
Hepatitis B	Birth through 18 years	All
Hib ₁	6 weeks through 59 months, certain 5-18 year olds	See footnote
HPV	9 years through 18 years, females	All
Influenza (2008-09 Season)	6 months through 18 years	All
IPV	6 weeks through 17 years	All
Meningococcal Vaccines (Both Polysaccharide and Conjugate), MPSV4 & MCV4 ₂	Ages 2 years through 18 years, as indicated	See footnotes
MMR	1 year through 18 years	All
MMR/V	1 year through 12 years (including 2 nd doses when indicated)	All
Pneumococcal Conjugate (PCV7)	6 weeks through 59 months	All
Pneumococcal Polysaccharide 23-valent (PPV23) ₃	2 through 18 years, high risk	See footnotes
Rotavirus	6 through 32 weeks	All
Td	7 through 18 years	All
Tdap	10 through 18 years	All
Varicella	Two doses for all 1 year through 18 years	All

1) One dose of Hib vaccine is available for unimmunized high-risk children ages 5 through 18 years, with either functional or anatomical asplenia (SSD, post-splenectomy);

immunodeficiency; HIV infection; or immunosuppression due to HIV infection or chemotherapy.

- 2) **MCV4 & MPSV4: MCV4 is the preferred vaccine, whenever possible.** Routine vaccination of all persons aged 11--18 years with 1 dose of **MCV4** at the earliest opportunity. Persons aged 11--12 years should be routinely vaccinated at the 11--12 years health-care visit as recommended by ACIP. **MCV4** is also available for children ages 2 through 10 years at high risk, including: persons with terminal complement component deficiencies; persons with anatomic or functional asplenia; travelers to or residents of Sub-Saharan Africa's "Meningitis Belt," during December to June; visitors to Mecca in Saudi Arabia during annual Hajj; countries in which *N. Meningitidis* is hyperendemic or epidemic; also recommended for use in control of meningococcal outbreaks. MPSV4 is available as above when MCV4 is not available or contraindicated.
- 3) Pneumococcal Polysaccharide (23-valent) vaccine is available for high-risk children 2 years through 18 years as follows:

IMMUNOCOMPETENT PERSONS	HIGH RISK PERSONS
Persons aged 2–18 years with chronic illness, including <ul style="list-style-type: none"> o Cardiovascular disease o Pulmonary disease (e.g. COPD, emphysema, NOT asthma) o Cochlear implants o Diabetes mellitus o Alcoholism, chronic liver disease o CSF leaks 	Persons 2–18 years of age immunocompromised due to: <ul style="list-style-type: none"> o HIV infection, o Hodgkins disease, multiple myeloma, generalized malignancy, leukemia o chronic renal failure or nephrotic syndrome o organ or bone marrow transplants o immunosuppression o functional or anatomic asplenia (sickle cell disease or splenectomy)
Persons aged 2–18 years living in nursing homes and other long-term care facilities	

If you have any questions about this chart or the footnotes, or would like the document in an alternate format (e.g. Braille), please call the Public Health Division Immunization Program at (971) 673-0300.