

Section E: Appendix

USDA and Adult Care Food Program (E28 – E30)

How do I sign up for the program?

- Child Care Centers (Replace phone number) Call (503) 947-5902
- Registered and certified family child care providers (Replace after first sentence) To locate a sponsor contact the Oregon Department of Education Child Nutrition Program. Call (503) 947-5902. Visit the ODE website <http://www.ode.state.or.us/search/results/?id=209> for more information about the USDA Child and Adult Care Food Program.

How much money will I receive?

- (Replace website) <http://www.ode.state.or.us/search/results/?id=209>

Specific Illnesses: E 31-61

Clarify Information about Varicella (Chicken Pox) (E34-35)

Infectious period (Replace)

The infectious period begins one to two days before the rash and continues until the last blister is scabbed over. Report outbreaks of several cases to the local health department immediately.

Ways to limit spread (Replace)

- Children and adults can return after the last blister has scabbed over.
- See C59 regarding vaccination

Steps child care can take (Add)

- Do not give aspirin, **or products containing aspirin (Pepto-Bismol, Kaopectate, Alka Seltzer)**, to the child, as there appears to be an association between aspirin use and Reyes Syndrome (a serious illness which can cause death).
- Child care providers who have not had chickenpox should be tested for immunity (through a blood test) and should receive the chickenpox/varicella vaccine if the test shows that they are not immune.

Clarify Information about Influenza/Flu (E 40)

Steps child care can take (Add)

- Do not give aspirin, **or products containing aspirin (Pepto-Bismol, Kaopectate, Alka Seltzer)**, to the child as there appears to be an association between aspirin use and Reye Syndrome (serious illness that can cause death).

Clarify Information about Hepatitis A (E 43)

Infectious period (Replace)

For one to two weeks before first stomach symptoms occur until one week after jaundice begins.

Ways to limit spread (Replace 3d sentence)

Routine vaccination of children is recommended.

Clarify Information about Hepatitis B (E 44)

Incubation period (Add)

45 to 180 days (**usually 2 to 3 months**)

Clarify Information about Measles (E 49)

Ways to limit spread (Add)

Measles is **highly contagious**. Children and staff with measles should be excluded from child care while they are infectious.

Steps child care can take (Add)

- Exclude while infectious.
- Children or staff without 2 doses of measles vaccine may be excluded from facility if there is not time to give a second dose following an exposure. Staff should have received 2 doses of MMR vaccine.

Clarify Information about Mumps (E 52)

Ways to limit spread (Replace)

Children should receive mumps vaccine according to the current childhood immunization schedule (mumps is included in the MMR vaccine). Adults who have never had mumps or who have not yet received a vaccine against mumps should also be immunized.

Steps child care can take (Add)

- Exclude people with mumps while infectious.

Clarify Information about Whooping Cough/Pertussis (E 60-61)

First paragraph

- Delete last sentence and replace with, “A booster form of the vaccine is now available.”

Steps child care can take (Add)

- Make sure that child care providers are immunized against pertussis
- Exposed people may need antibiotics to prevent disease, even if they have no symptoms, regardless of immunization history.