

# EXHIBIT 3

Referenced in OAR 333-200-0080(5)(b)

## HOSPITAL TRAUMA TEAM RESPONSE CRITERIA

Systolic blood pressure <90 mmHg; or  
Respiratory distress with rate <10 or >29; or  
Airway management required; or  
Glasgow Coma Scale  $\leq$  10; or  
Penetrating injury of the neck, torso, or groin; or  
Flail chest; or  
Two or more obvious long-bone (humerus/femur) fractures; or  
Spinal cord injury with limb paralysis.

**YES**



Activate  
Full Trauma Team

### Full Trauma Team

- General Surgeon
- Emergency Physician
- Anesthesiologist
- Emergency Nurse(s)
- Laboratory
- Radiology
- Respiratory Therapist

Glasgow Coma Scale of 11 or 12; or  
Isolated penetrating injury above base of skull; or  
Isolated amputation above the wrist or ankle; or  
Death of a same car occupant; or  
Ejection of patient from an enclosed vehicle; or  
Heavy extrication time >20 minutes; or  
EMT suspects life threatening injuries related to high energy transfer situations or to the presence of co-morbid factors.

**YES**



Activate  
Modified Trauma Team

### Modified Trauma Team

- Emergency Physician
- Emergency Nurse(s)
- Laboratory
- Radiology

## HIGH ENERGY TRANSFER SITUATION/CO-MORBID FACTORS

- Falls >20 feet; or
- Pedestrian hit at 20 mph or thrown 15 feet; or
- Rollover; or
- Motorcycle, ATV or bicycle crash; or
- Significant impact or intrusion into occupant space of vehicle; or
- Extremes of age <5 or >55 years; or
- Patient with bleeding disorder or patient on anticoagulants; or
- Medical illness: cardiac or respiratory disease, insulin-dependent diabetes, cirrhosis, or morbid obesity; or
- Pregnancy; or
- Immunosuppressed patients; or
- Presence of intoxicants.

Trauma system hospitals are responsible for eliciting either a full or modified trauma team response based on information obtained from the on-line prehospital report, as set forth in Exhibit 3 and 4 of these rules.

**Full Trauma Team** activation means the entire trauma team is activated, including the general surgeon as prescribed by OAR 333-200-0090, Exhibit 4.

**Modified Trauma Team** activation allows for the general surgeon to be activated at the discretion of the emergency physician at Level II, III and IV (if applicable) facilities. At Level I facilities, a modified response allows for anesthesia to be activated at the discretion of the emergency physician or trauma surgeon.

**Effective Date: May 4, 2000**