


Colorectal cancer screening among ages 50–75 years by county, Oregon, 2014–2017



Colorectal cancer screening among ages 50–75 years		
by county, Oregon, 2014–2017		
County	Percent of adults	Number of respondents (N)
Baker	67.9%	59
Benton	74.0%	210
Clackamas	75.0%	725
Clatsop	66.7%	106
Columbia	70.1%	155
Coos	54.6%	163
Crook	69.8%	67
Curry	66.1%	66
Deschutes	66.9%	311
Douglas	70.9%	292
Gilliam / Sherman / Wasco	66.5%	77
Grant	*	27
Harney	*	20
Hood River	59.2%	50
Jackson	67.4%	403
Jefferson	60.5%	50
Josephine	57.5%	190
Klamath	57.8%	149
Lake	28.3%†	31
Lane	65.6%	752
Lincoln	74.0%	156
Linn	71.7%	261
Malheur	55.5%	54
Marion	71.0%	536
Morrow	*	19
Multnomah	69.9%	1,240
Polk	73.5%	172
Tillamook	67.9%	71
Umatilla	70.1%	112
Union	59.5%	66
Wallowa	68.9%	37
Washington	74.1%	831
Wheeler	*	4
Yamhill	61.9%	165
Frontier ZIP codes	60.1%	234
Rural ZIP codes	66.1%	2,997
Urban ZIP codes	71.9%	4,211
Oregon State	68.7%	7,627

[†] Rates with a relative standard error (RSE) ≥ 30 are considered unreliable.

^{*} Rates based on a count < 3 or a denominator < 30 or with a relative standard error (RSE) \geq 50 are not reported.

About the Data

Data Source: Oregon Behavioral Risk Factor Surveillance System (BRFSS)

Data Notes:

Number of respondents (N) is unweighted; percentages are weighted

Colorectal cancer screening includes fecal occult blood test (FOBT) in the past year or a colonoscopy within 10 years or a sigmoidoscopy within the past 5 years with FOBT within the past 3 years.

Date: September 6, 2019