Year	Low Birthweight	First Trimester Care	No Care	Inadequate Care <sup>1</sup>	Third Trimester Care	Less than Five Visits
1000	50.4	700.0	F F	59.0	25.0	A 4 - A
1980 1981	50.4 48.5	780.8 775.6	5.5 8.9	58.0 63.1	35.2 38.6	41.4 43.0
1981	40.5	769.3	0.9 11.2	70.3	30.0 41.0	43.0 48.0
1982	50.0	775.3	11.2	66.5	38.5	48.0
1983	51.5	771.5	11.0	68.2	41.1	46.2
1904	51.5	111.5	11.0	00.2	41.1	40.2
1985	51.3	752.0	12.1	72.9	43.7	47.5
1986	51.3	738.7	11.7	83.3	52.1	54.6
1987	54.0	736.8	16.5	86.2	50.3	58.5
1988	52.6	738.8	13.8	83.6	49.9	54.7
1989	52.2	750.7	12.0	73.2	42.9	48.7
1990	50.1	757.1	10.7	70.0	43.4	45.1
1991	49.2	768.2	8.7	61.0	37.4	38.6
1992	51.8	787.0	8.2	52.6	31.4	34.0
1993	52.5	794.6	7.6	51.7	30.4	33.8
1994	53.0	790.9	8.5	57.8	34.3	36.4
1005					o ( =	
1995	54.9	787.7	8.6	58.4	34.7	38.2
1996	53.5	799.3	7.1	53.7	31.7	34.8
1997	55.0	811.2	6.7	50.0	29.6	32.3
1998	53.7	807.2	7.2	53.5	30.7	35.3
1999	53.9	809.9	7.3	53.7	29.6	35.7
2000	56.6	812.8	8.5	55.9	29.8	36.6
2001	55.6	815.2	8.0	50.5	28.7	33.1
2002	57.9	816.4	9.4	52.2	28.6	35.7
2003	61.6	810.7	11.7	55.5	28.6	38.4
2004	60.6	804.3	10.9	57.9	30.3	41.0
2005	61.2	810.0	8.9	58.3	30.1	40.8

## TABLE 1-6. Oregon Rates of Low Birthweight, and Measures of Prenatal Care, 1980-2005

<sup>1</sup> Inadequate prenatal care is defined as care that began in the third trimester or consisted of less than five prenatal visits.

All rates are per 1,000 live births. Rates and percentages are calculated excluding missing and unknown values.