

**Induced abortions by length of gestation, procedure type, complications, and age of patient, Oregon residents, 2012-2014**

**Wheeler County**

Medical and health characteristics	Total	Patient's age				
		10-14	15-17	18-19	20+	N.S.
<b>Total</b> .....	3	—	—	1	2	—
<b>Gestation (weeks)</b>						
< 9 .....	2	—	—	1	1	—
9-12 .....	1	—	—	—	1	—
13-16 .....	—	—	—	—	—	—
17-20 .....	—	—	—	—	—	—
21-22 .....	—	—	—	—	—	—
23+ .....	—	—	—	—	—	—
Unknown .....	—	—	—	—	—	—
<b>Contraceptive use</b>						
None .....	2	—	—	—	2	—
Birth control pills .....	—	—	—	—	—	—
Condoms .....	—	—	—	—	—	—
Other contraceptive .....	1	—	—	1	—	—
Unknown .....	—	—	—	—	—	—
<b>Procedure type</b>						
Suction curette .....	2	—	—	—	2	—
Medical (non-surgical) .....	1	—	—	1	—	—
Dilation & evacuation .....	—	—	—	—	—	—
Intra-uterine instillation .....	—	—	—	—	—	—
Vaginal prostaglandin .....	—	—	—	—	—	—
Sharp curettage .....	—	—	—	—	—	—
Other .....	—	—	—	—	—	—
Unknown .....	—	—	—	—	—	—
<b>Complications</b>						
None .....	3	—	—	1	2	—
Hemorrhage .....	—	—	—	—	—	—
Infection .....	—	—	—	—	—	—
Uterine perforation .....	—	—	—	—	—	—
Cervical laceration .....	—	—	—	—	—	—
Retained products .....	—	—	—	—	—	—
Failure of first method .....	—	—	—	—	—	—
Other .....	—	—	—	—	—	—
Multiple complications .....	—	—	—	—	—	—
Unknown .....	—	—	—	—	—	—

— Quantity is zero.

NOTE: Includes reported abortions to Oregon residents including those obtained out-of-state. Because some states (e.g., California) do not record data on residence for abortion patients, not all out-of-state abortions are included.