

Induced abortions by length of gestation, procedure type, complications, and age of patient, Oregon residents, 2014-2016

Gilliam County

Medical and health characteristics	Total	Patient's age				
		10-14	15-17	18-19	20+	N.S.
Total	3	*	*	*	1	—
Gestation (weeks)						
< 9	2	*	*	*	—	—
9-12	—	—	—	—	—	—
13-16	—	—	—	—	—	—
17-20	1	*	*	*	1	—
21-22	—	—	—	—	—	—
23+	—	—	—	—	—	—
Unknown	—	—	—	—	—	—
Contraceptive use ¹						
None	3	*	*	*	1	—
Birth control pills	—	—	—	—	—	—
Condoms	—	—	—	—	—	—
Other contraceptive	—	—	—	—	—	—
Unknown	—	—	—	—	—	—
Procedure type						
Suction curette	1	*	*	*	—	—
Medical (non-surgical)	—	—	—	—	—	—
Dilation & evacuation	2	*	*	*	1	—
Intra-uterine instillation	—	—	—	—	—	—
Vaginal prostaglandin	—	—	—	—	—	—
Sharp curettage	—	—	—	—	—	—
Other	—	—	—	—	—	—
Unknown	—	—	—	—	—	—
Complications ²						
None	3	*	*	*	1	—
Hemorrhage	—	—	—	—	—	—
Infection	—	—	—	—	—	—
Uterine perforation	—	—	—	—	—	—
Cervical laceration	—	—	—	—	—	—
Retained products	—	—	—	—	—	—
Failure of first method	—	—	—	—	—	—
Other	—	—	—	—	—	—
Multiple complications ³	—	—	—	—	—	—

— Quantity is zero.

¹ Multiple contraceptive methods may be reported for a single patient.

² Reported complications. Categorized as none if no specific complication was reported.

³ Patients having more than one complication are listed here. Their complications are not listed above.

* Detailed reporting of small numbers may breach confidentiality.

NOTE: Includes reported abortions to Oregon residents including those obtained out-of-state. Because some states (e.g., California) do not record data on residence for abortion patients, not all out-of-state abortions are included.