

Induced abortions by length of gestation, procedure type, complications, and age of patient, Oregon residents, 2014-2016

Malheur County

Medical and health characteristics	Total	Patient's age				
		10-14	15-17	18-19	20+	N.S.
Total	98	2	4	7	82	3
Gestation (weeks)						
< 9	59	1	3	5	48	2
9-12	23	1	—	—	21	1
13-16	10	—	—	1	9	—
17-20	3	—	1	1	1	—
21-22	2	—	—	—	2	—
23+	—	—	—	—	—	—
Unknown	1	—	—	—	1	—
Contraceptive use¹						
None	91	2	3	7	77	2
Birth control pills	1	—	1	—	—	—
Condoms	3	—	—	—	3	—
Other contraceptive	3	—	—	—	2	1
Unknown	—	—	—	—	—	—
Procedure type						
Suction curette	48	1	3	2	42	—
Medical (non-surgical)	41	1	—	4	34	2
Dilation & evacuation	8	—	1	1	5	1
Intra-uterine instillation	—	—	—	—	—	—
Vaginal prostaglandin	—	—	—	—	—	—
Sharp curettage	—	—	—	—	—	—
Other	1	—	—	—	1	—
Unknown	—	—	—	—	—	—
Complications²						
None	98	2	4	7	82	3
Hemorrhage	—	—	—	—	—	—
Infection	—	—	—	—	—	—
Uterine perforation	—	—	—	—	—	—
Cervical laceration	—	—	—	—	—	—
Retained products	—	—	—	—	—	—
Failure of first method	—	—	—	—	—	—
Other	—	—	—	—	—	—
Multiple complications ³	—	—	—	—	—	—

— Quantity is zero.

¹ Multiple contraceptive methods may be reported for a single patient.

² Reported complications. Categorized as none if no specific complication was reported.

³ Patients having more than one complication are listed here. Their complications are not listed above.

NOTE: Includes reported abortions to Oregon residents including those obtained out-of-state. Because some states (e.g., California) do not record data on residence for abortion patients, not all out-of-state abortions are included.