Induced abortions by length of gestation, procedure type, complications, and age of patient, Oregon residents, 2014-2016

Medical and health characteristics	T . (.)	Patient's age				
	Total	10-14	15-17	18-19	20+	N.S.
Total	4	*	*	*	3	_
Gestation (weeks)	-				0	
< 9	1	*	*	*	_	_
9-12	1	*	*	*	1	_
13-16	1	*	*	*	1	_
17-20	-	_	_	_	-	_
21-22	_	_	_	*	_	_
23+	1	*	*	*	1	_
Unknown	-	_	_	_	-	_
Contraceptive use ¹						
None	3	*	*	*	3	_
Birth control pills	-	_	_	_	-	_
Condoms	_	_	_	_	_	_
Other contraceptive	1	*	*	*	_	_
Unknown	-	_	_	_	_	_
Procedure type						
Suction curette	2	*	*	*	2	_
Medical (non-surgical)	1	*	*	*	-	_
Dilation & evacuation	1	*	*	*	1	_
Intra-uterine instillation	-	_	_	_	-	_
Vaginal prostaglandin	_	_	_	_	_	_
Sharp curettage	_	_	_	_	_	_
Other	_	_	_	_	_	_
Unknown	_	_	_	_	_	_
Complications ²						
None	4	*	*	*	3	_
Hemorrhage		_	_	_	-	_
Infection	_	_	_	_	_	_
Uterine perforation	_	_	_	_	_	_
Cervical laceration	_	_	_	_	_	_
Retained products	_	_	_	_	_	_
Failure of first method	_	_	_	_	_	_
Other	_	_	_	_	_	_
Multiple complications ³	_	-	_	_	_	_

Wheeler County

Quantity is zero.

Quantity is zero.
Multiple contraceptive methods may be reported for a single patient.
Reported complications. Categorized as none if no specific complication was reported.
Patients having more than one complication are listed here. Their complications are not listed above.

* Detailed reporting of small numbers may breach confidentiality.

NOTE: Includes reported abortions to Oregon residents including those obtained out-of-state. Because some states (e.g., California) do not record data on residence for abortion patients, not all out-of-state abortions are included.