


Effectiveness of Family Planning Methods



How to make your method most effective
 After procedure, little or nothing to do or remember.
Vasectomy and hysteroscopic sterilization: Use another method for first 3 months.

Injectable: Get repeat injections on time.
Pills: Take a pill each day.
Patch, Ring: Keep in place, change on time.
Diaphragm: Use correctly every time you have sex.

Condoms, sponge, withdrawal, spermicides: Use correctly every time you have sex.
Fertility awareness-based methods: Abstain or use condoms on fertile days. Newest methods (Standard Days Method and TwoDay Method) may be the easiest to use and consequently more effective.

CS 242797

CONDOMS SHOULD ALWAYS BE USED TO REDUCE THE RISK OF SEXUALLY TRANSMITTED INFECTIONS.

Other Methods of Contraception

Lactational Amenorrhea Method: LAM is a highly effective, temporary method of contraception.

Emergency Contraception: Emergency contraceptive pills or a copper IUD after unprotected intercourse substantially reduces risk of pregnancy.

Adapted from World Health Organization (WHO) Department of Reproductive Health and Research, Johns Hopkins Bloomberg School of Public Health/Center for Communication Programs (CCP). Knowledge for health project. Family planning: a global handbook for providers (2011 update). Baltimore, MD; Geneva, Switzerland: CCP and WHO; 2011; and Trussell J. Contraceptive failure in the United States. Contraception 2011;83:397-404.



U.S. Department of Health and Human Services
 Centers for Disease Control and Prevention