EXHIBIT 3

OAR Chapter 333, Division 200

OREGON HOSPITAL TRAUMA TEAM ACTIVATION CRITERIA

Systolic blood pressure <90 mmHg; or

Respiratory rate <10 or >29 breaths per minute (<20 in infant aged <1 year); or

Need for ventilatory support; or

Glasgow Coma Scale <9; or

All penetrating injuries to head, neck, torso, and extremities proximal to elbow or knee; or

Chest wall instability or deformity (e.g. flail chest); or

Two or more proximal long-bone fractures; or

Suspected spinal cord injury with motor sensory deficit; or

Transfers requiring blood transfusions; or

Emergency physician's discretion.

Glasgow Coma Scale of 9 - 13; or

Crushed, degloved, mangled, or pulseless extremity; or

Amputation proximal to wrist or ankle; or

Open or depressed skull fracture; or

Suspected pelvic fracture; or

Falls

- Adults: >20 feet (one story is equal to 10 feet); or
- Children: >10 feet or two or three times the height of the child;

High-risk auto crash

- Intrusion, including roof: >12 inches occupant site; >18 inches any site; or
- Ejection (partial or complete) from automobile; or
- Death in same passenger compartment; or
- Vehicle telemetry data consistent with a high risk of injury; or

Auto vs. pedestrian/bicyclist thrown, run over, or with significant (>20 mph) impact; or

Motorcycle or ATV crash >20 mph; or

EMS provider or receiving hospital judgment

YES



Activate **Full Trauma Team**

Full Trauma Team

- General Surgeon
- Emergency Physician
- Emergency Nurse(s)
- Laboratory
- Radiology
- Respiratory Therapist

Response times from patient arrival:

Level I and II – 15 min.

Level III and IV - 30 min

YES



Activate Modified Trauma Team

Modified Trauma Team

- Emergency Physician
- Emergency Nurse(s)
- Laboratory
- Radiology

CO-MORBID FACTORS

Older Adults

- Risk of injury/death increases after age 55 years; or
- SBP <110 might represent shock after age 65 years; or
- Low impact mechanisms (e.g. ground level falls) might result in severe injury; or

Children

Should be triaged preferentially to pediatric capable trauma centers; or

Pregnancy >20 weeks; or

- Without other trauma mechanism: triage to burn facility; or
- With trauma mechanism: triage to trauma center; or

Anticoagulants and bleeding disorders

 Patients with head injury are at high risk for rapid deterioration; or

Time sensitive extremity injury

Effective: January 1, 2016