**Maryland Emergency Medical Services** 

# Interhospital Transfer Resource Manual

Developed by the

Maryland Institute for Emergency Medical Services Systems

> Revised March 2021



Previous editions published:

January 1986

April 1994

January 2002

November 2009

October 2014

October 2019

November 2020

# Interhospital Transfer Resource Manual Table of Contents

Introduction	V										
The Maryland Emergency Medical System: Overview	vi										
Facility Acronyms											
Transportation Information											
How to Initiate a Referral and Transport	1										
Maryland Universal Interhospital Hand-Off Transfer Form and Instructions	2										
Transport Services	4										
Maryland EMS Clinician Descriptions	8										
Adult Trauma Centers and Guidelines											
List of Adult Trauma Referral Centers	9										
Map of Adult Trauma Referral Centers	11										
Adult Trauma Center Guidelines for Transfer	12										
Burn Injury (Adult)											
Eye Trauma											
Hyperbaric and Dive Medicine											
Hand and Upper Extremity Trauma											
Neurotrauma	25										
Special Pathogen/Highly Infectious Diseases											
Stroke Guidelines for Transfer											
Primary Stroke Centers											
List of Primary Stroke Centers	39										
Map of Primary Stroke Centers	42										
Comprehensive Stroke Centers											
List of Comprehensive Stroke Centers	43										
Map of Comprehensive Stroke Centers	44										
List of Endovascular Capable Centers in Maryland	45										
Map of Endovascular Capable Centers in Maryland	46										
Acute Ischemic Stroke Guidelines for Potential Endovascular	47										
Recanalization Therapy (ERT)											
Cardiac											
List of Cardiac Interventional Centers	51										
Map of Cardiac Interventional Centers	53										
ST-Elevation Myocardial Infarction (STEMI) Guidelines for Transfer	54										

Pediatric									
List of Pediatric Trauma Centers	57								
Map of Pediatric Trauma Centers	58								
Pediatric Trauma Centers' Guidelines	59								
List of Pediatric Burn Centers									
Map of Pediatric Burn Centers	63								
Indications for Transfer to a Pediatric Burn Center	64								
Perinatal									
List of Perinatal Referral Centers	65								
Map of Maryland Perinatal Referral Centers	66								
Maternal Referral/Consultation	67								
Neonatal Referral/Consultation	69								
Emergency Medical Treatment and Active Labor Act (EMTALA)									
Explanation of Regulations	74								
Sample Forms	81								
Transport of Stable Patients from a Trauma/Burn or Specialty									
Center to a Community Hospital or Rehabilitation/Long-Term Care Facility	84								
Interhospital Transfer Checklist	86								

# INTRODUCTION

This version of the Maryland Emergency Medical Services Interhospital Transfer Resource Manual replaces the manual dated October 2019. This updated manual reflects current practices and changes that have occurred in the Maryland Emergency Medical Services (EMS) System. The updated manual has been organized and tabbed to facilitate the quick retrieval of information.

The address and telephone number for each specialty center has been updated. Additionally, the information found in the section addressing EMTALA has been updated to reflect the most current language.

Direct communication needs to occur between the sending and receiving hospital staffs to arrange an interhospital transfer as specified in the manual. However, MIEMSS Emergency Medical Resource Centers (EMRCs) and System Communications (SYSCOM) are available to assist, if needed, to arrange transfers. SYSCOM, staffed by MIEMSS and the Maryland State Police (MSP), is the communications core for MSP helicopter dispatch, coordination, and monitoring.

Typically, tertiary care hospitals in and around Maryland are able to assist referring hospitals with the transfer of patients from community hospitals by both ground and air ambulances. In consultation with the referring physician, the receiving hospital is able to determine the appropriate transport method for specific patients. Interhospital air transportation is ideally accessed through the receiving hospital. However, if a referring physician or patient desires contact information for an alternate air clinician, he/she may call SYSCOM, which maintains a list of all available clinicians in the area.

To access SYSCOM, call 1-800-648-3001. SYSCOM remains the point of contact to arrange MSP helicopter transportation. (See page 1.)

For questions, or if you are having trouble contacting a specific trauma and/or specialty referral center, contact EMRC at 1-800-492-3805.

# THE MARYLAND EMERGENCY MEDICAL SYSTEM Overview

The Maryland Institute for Emergency Medical Services Systems (MIEMSS) is the state agency that coordinates the statewide system of EMS. MIEMSS oversees and coordinates all components of the statewide EMS system (including planning, operations, evaluations, and research), provides leadership and medical direction, conducts and/or supports EMS educational programs, operates and maintains a statewide communications system, designates trauma and specialty centers, licenses and regulates commercial ambulance services, licenses EMS clinicians, and participates in EMS-related public education and prevention programs.

Maryland EMS legislation, passed in 1993, mandated regulations for the designation of trauma centers and specialty centers. MIEMSS, with the approval of the EMS Board, defined and designated four levels of trauma centers.

The following types of specialty care centers have been designated in Maryland to provide specialized services to patients with certain types of illnesses or injuries, including Adult and Pediatric Burn, Eye Trauma, Hand and Upper Extremity Trauma, Neurotrauma, Perinatal, Stroke, Cardiac Interventional, and Hyperbaric and Dive Medicine.


In this coordinated system of emergency care, critically ill and injured patients are transported to the hospital that is best staffed, equipped, and experienced to treat their injuries or illness. This manual has been developed by MIEMSS, in collaboration with the trauma and specialty referral centers, to assist emergency department (ED) personnel in identifying those patients with specialty care needs who should be transferred to a trauma or specialty center.

The manual also provides reference information for access numbers and locations to the various specialty centers.

The Maryland State Police Aviation Command (MSPAC) medevac program provides helicopter transportation of critically ill or injured patients requiring time-critical transport to a higher level of care. Interhospital transports are provided by commercial ground and air ambulance companies throughout the state. This manual includes contact information for MSP medevac, commercial air ambulance, and commercial ground ambulance transportation.

# **FACILITY ACRONYMS**

JHM	Johns Hopkins Medicine
MedStar	MedStar Health
UM	University of Maryland



# **HOW TO INITIATE A REFERRAL AND TRANSPORT**

- 1. The referring physician should contact the receiving tertiary care hospital. The receiving physician must confirm that the transfer is accepted
  - Once acceptance of the patient is confirmed, the mode of transport is determined by the referring and receiving physicians based on the patient's medical needs during transport, as well as the need to minimize out-of-hospital transport time.
- 2. If helicopter transport is the indicated mode of transport:
  - a) The receiving referral center will arrange air transportation.
  - b) The following patient information should be provided:
    - Approximate weight and age
    - Suspected major injuries or medical condition
    - · Level of consciousness and airway status
    - Most recent vital signs
    - Ongoing therapies
    - Specialized equipment, (e.g., isolette)
- 3. If ambulance is the indicated mode of transport:
  - a) Either the referring or receiving hospital will contact an ambulance service of its choice that is capable of providing the level of care required, or
  - b) If the patient requires a critical level of care, or care outside the scope of practice of the Advanced Life Support (ALS) clinician, the hospital or the commercial ambulance service must provide for a supplemental clinician capable of providing the care required.
- 4. Information needed by the receiving center:
  - · Referring physician's name
  - Referring hospital
  - Location of patient within the hospital
  - Call-back number
  - Interhospital transportation documentation (eMEDS® Short Form)

#### Patient information needed:

- Name, age, and weight
- Needs an interpreter
- Mechanism of injury
- Type and extent of injury
- Treatment rendered
- Status of other family members injured and destination (if known)
- 5. A copy of all medical records must be sent with the patient, including:
  - Progress notes
  - Nursing notes
  - Medication and fluid records
  - · Radiologic studies
  - Laboratory results
  - Fact sheet that includes patient address, phone number, and emergency contact information
  - eMEDS® or Short Form
  - EKG
  - · Radio Consult Form

# PROMPT TRANSPORT

**Do not delay** transport while awaiting laboratory or radiology results. These can be communicated by phone and FAX as they become available.

# MARYLAND UNIVERSAL INTERHOSPITAL HAND-OFF TRANSFER FORM (Optional Use)

Patient Identification Label

		TIME-SPECIFIC																	PATIENT ASSESSMENT (most recent) SYSTEMS ASSESSMENT & PAST MEDICAL HISTORY PERTINENT TO TRANSFER													PATIENT INFORMATION														
Form Completed By Name:	Sending Facility Contact Name:	Sending Physician Name:	Social Concern: □Domestic Violence Suspicion, □Child/Elder; □ Suspicion □Neglect	Spinal Protection:   Backboard   C-Collar   Pelvic   Extremity:	Medicines in the first of the property of the	Mechanism of Injury:   MVC   MCC   Ded   ATV	TRAUMA: Date of Injury Time of Injury	NIHSS    NIHSS   NIHSS    NIHSS   NIHSS    NIHSS    NI	Patient Last Known v	Post Known Well: Date	ring EKG Date Time	CARDIAC: Acute Symptom Onset Date			wiedication	MEDICA IONS GIVEN PRIOR IO IRANSFER AI HOSPI AL: Antibiotics, Resuscitative, Sedative/Paralytic, Inrombolytics/I PA, Others	מי רואסיייייייייייייייייייייייייייייייייייי	ODERATIVE DROCEDIBES:	ומפט מפט		CACHONO	ISOI ATION BRECALLTIONS: DNo. DMBSA D	PRE-EXISTING MEDICATION: Anticoagulants   Y   N:	Thorax/Lungs "Respiratory":	Nervous/Neurological:	Musculoskeletal:	Integumentary:	HEENT	Genitourinary:	Gastrointestinal:	Cardiovascular:	Abnormal Assessment Findings	ALLERGIES:	GCS:Eye Wotor Verbal = I	ıy: □Natural □LMA □King □ ETT; Size		care Representative/Proxy/Legal Guardi	INFORMATION GIVEN TO CONTACT PERSON: _	Contact with Patient? □Y □N, Contact Notified of Transfer? □Y (Date	CONTACT PERSON: LAST NAME	MEDICAL ORDERS for LIFE SUSTAINING TREATMENT (MOLST): Documents □Y □N; Attached	PATIENT NAME: LAST	□Stroke Intervention □Adult Trauma □Pediatric Trauma	TRANSPORT MODE: DIAGNOSIS	TRANSFER DATE: TRANSFER TIME:	
			Child/Elder; ☐ Suspicion ☐Ne	Extremity;RULUI		/ Bicycle Beating Stabb		Dysphagia Screening Before Oral Intake □Y □N; □ Pass □Fail,			□EMS □ED, First Call CIC Date	Time Duration of Pain			Dosage	HUSPITAL: Antibiotics, Resu			Trop Myo Other	notine in Choose Turbo		ESBI TO D#	□N: Name:											lotal Pupil size: (mm) K	Depth	Kg. T   P	LAST NAME		of Transfer? □Y (Date	FIRST	TMENT (MOLST): Documents	FIRST	rauma □Specialty Referral; □Hand □Eye	DIAGNOSIS RELATED TO TRANSFER:	TIME:ETA:	
			□Abuse,			ing □Cut/Dierce □Eirearm □B	Death in same incident $\Box Y \Box N$ , Family Member $\Box Y \Box N$ , Was patient notified? $\Box Y \Box N$	ı			ate Time				Date	uscitative, Sedative/Paralytic,			-  -  -  -  -  -  -  -  -  -  -  -  -	ullis, Flatelets/ullis,	mits Blatslats/	Othor opposite	Other:										IMMUNIZATIONS/SCREENING DATES:		- 1	R			Time	RELATIONSHIP_	□Y □N; Attached □Y □N	DOB: AGE:			TRANSFER FROM:	
			Reported? 🗆 Y 🗆 N, To Whom?	Other		≀ırn □Eall □Drownin	Y  □N, Was patient n	Had stroke symptoms resolved by presentation □Y	, Stroke Activation.		. ED Discharge Date	ED Arrival Date			- IIII e	I hrombolytics/ I PA,	0111170	OTHERS	IMAGING: Films Sent   Y   N: Y-Rays		Othor Olle	0#5											DATES: Flu	Abnormal Knythm	; Tolerating	Pulse Ox	) -		) □N, Contact Enro	PHONE: Day			□Adult Burn □Pediatric Burn □Other			
Date: Ti	Receiving Facility Contact Name:	Receiving Physician Name:			Digot	g □Rlast □Other	otified? □Y □N	presentation □Y □N			)ate Time	Time			Medication	Others			טו טמוווא. וטנמו כו איזמווטומי		/ <del>;</del>			□Trach	□TIA □Stroke □Shunt □AVM				□Renal Failure		□CABG □CAD □AICD □MI □Pacemaker □Stent □Other	Past Medical History	_, Tetanus, Pneumonia _		ng □Y □N, OG Tube □Y □N, CPR □Y □N	nula			Contact Enroute to Receiving Hospital? □Y □N	ayNight		□M □F Pregnant? □Y □N LMP		REASON FOR TRANSFER: □Hi	TRANSFER TO:	
Time:	*								,						Dosage										Shunt □AVM □Hemorrhage □ Other						Pacemaker  S		, PPD_		SPR OY ON	L, Pain Score_			N	Cell				gher Level of C		
															Date				MBI	Data States					□ Other						Stent Other		; □Pos □Neg			□Numbers □Faces				Cellular				ANSFER: □Higher Level of Care; □Cardiac Intervention		
															ıme	1				Time of													ig			ices								rvention		

# THE MARYLAND UNIVERSAL INTERHOSPITAL HAND-OFF TRANSFER FORM Instructions

#### **Background:**

The Quality Improvement Committees for the specialty centers (Trauma, Stroke, Cardiac, and Perinatal) identify, address, and develop recommendations/solutions to issues/problems brought forth by the hospital specialty referral centers. One such issue identified by the centers is the disconnect between the report given from the sending hospital to the receiving hospital. Many receiving centers have indicated that reports issued by the sending hospital do not always provide a true representation of the patient received. The Quality Improvement Committees investigated this issue and together developed the "Maryland Universal Interhospital Hand-off Transfer Form". The form is focused on time-critical information and provides standardization in the report format.

#### **Purpose:**

The "Maryland Universal Interhospital Hand-off Transfer Form" was designed to communicate pertinent and accurate clinical patient care information at the time of transfer to the receiving hospital. This form does not negate, but is in addition to, the Hospital's EMTALA documentation, necessary medical record and test results needed for the transfer of a patient.

The Joint Commission requires an appropriate "Hand-off" between caregivers. The Committee on Trauma American College of Surgeons' Resources for the Optimal Care of the Injured Patient identifies interhospital transfer documentation requirements, which may include a mutually agreed upon transfer form. This form will meet the requirements of both the Joint Commission and the American College of Surgeons. Use of the "Maryland Universal Interhospital Hand-off Transfer Form" is optional.

## **Process:**

The sending hospital will complete any of the form sections pertinent to its patient. A telephone report to the receiving hospital will be done utilizing the form content for standardization. Additional information may be given as well.

The Hand-off Form will be given to the Transport Team to "Hand-off" to the receiving hospital representative upon arrival. EMS clinicians are not required to document on this form.

# TRANSPORT SERVICES

#### INTRODUCTION

Transportation of critically ill and injured patients from a residence or the scene is usually accomplished by a public service ambulance (i.e., EMS, fire department or rescue squad ambulance, or Maryland State Police (MSP) medevac helicopter). The interhospital transportation of patients is primarily carried out by Maryland-licensed commercial ambulances, commercial air medical transport, or MSP medevac helicopters.

The EMS clinicians on all of these ambulances and medevac helicopters are licensed or certified by MIEMSS at one of the following levels: Emergency Medical Responder (EMR), Emergency Medical Technician (EMT), Cardiac Rescue Technician (CRT), or Paramedic. The EMR and EMT are Basic Life Support (BLS) clinicians and the CRT and Paramedic are Advanced Life Support (ALS) clinicians. All EMS clinicians, those working in the public service arena, and those working for commercial ambulance services must provide care as defined in *The Maryland Medical Protocols for Emergency Medical Services*.

#### PUBLIC SERVICE AMBULANCES

The primary mission of public service ambulances is to respond to the scene or a 9-1-1 call for emergency medical assistance from a residence or the scene. Rarely are these ambulances available to hospitals for interhospital transport. However, if a commercial ambulance is not able to respond for a patient requiring emergency transport to a referral center and if the public service ambulance is available, along with the required staff, they may agree to assist with an interhospital transport.

# LICENSED COMMERCIAL AMBULANCE SERVICES Basic Life Support (BLS)

A licensed BLS commercial ambulance is staffed at a minimum with an EMR driver and an EMT attendant. BLS may transport patients within the scope of practice of an EMT, which includes patients who are stable with maintenance IVs. BLS licensed ambulances may not add a nurse or other health care clinician to staff the ambulance for the purpose of caring for a patient who requires care outside the scope of practice of the EMT. These patients must be transported by an ALS licensed ambulance.

## Advanced Life Support (ALS)

A licensed ALS commercial ambulance is staffed at a minimum with an EMT driver and a CRT or Paramedic. The ALS clinician may start IVs, as well as administer certain medications and perform certain procedures that are listed in *The Maryland Medical Protocols for Emergency Medical Services*, which defines the scope of practice for all Maryland EMS prehospital clinicians. Patients requiring care outside the scope of practice of the ALS clinician must be accompanied by a health care clinician authorized by law to provide the level of care required and in accordance with EMTALA (refer to the EMTALA section, page 73).

# **Specialty Care Transport (SCT)**

An SCT commercial ambulance is staffed at a minimum with an EMT driver and two additional clinicians, and must be a licensed ALS vehicle. The scope of practice of the SCT-credentialed paramedic requires additional training, including additional medications and procedures and is defined in *The Maryland Medical Protocols for Emergency Medical Services*. If the patient's care is outside of the scope of practice of an SCT-credentialed paramedic, the first of the two additional clinicians must be either a nurse or physician with critical care expertise. The second clinician may be either an SCT-credentialed paramedic or a paramedic who has been oriented to specialty care transports. If the patient's care is within the scope of practice of the SCT-credentialed paramedic, the first clinician may be either a nurse with critical care expertise<sup>1</sup> or the specially oriented SCT-credentialed paramedic. The second clinician may be either a CRT licensed after July 1, 2001, or a paramedic who has been oriented to specialty care transports.

# **Neonatal Transports**

Licensed neonatal commercial ambulances are specialized ambulances that are staffed and equipped to transport critically ill newborns from their hospital of birth to a tertiary care facility. Transport of critically ill newborns may be carried out only in a licensed neonatal ambulance.

## AIR MEDICAL TRANSPORT

Typically, tertiary care hospitals in and around Maryland are able to assist referring physicians with the transfer of patients from community hospitals by both ground and air ambulance. In consultation with the referring physician, the receiving hospital is able to determine the appropriate transport method for specific patients. Interhospital air transportation is usually accessed through the receiving facility. Nearly all interhospital air transports are provided by commercial air services that are under contract to the hospital and that typically bill third party payors and/or patients for transport services. Should a referring physician or patient desire contact information for an alternate air provider, he/she may call SYSCOM at 1-800-648-3001; SYSCOM maintains a list of all available clinicians in the area. State police helicopters play a secondary role in interhospital transports since the majority of their flights are from the scene of an incident.

#### **Utilization of Air Medical Transport for Interhospital Transfer**

Air medical transport should be considered for interhospital transfer of patients for whom time is critical. An air ambulance is best used when a patient requires time-critical interventions or when it is important to minimize time out of a hospital setting. In either case, air transport would result in clinically significant reduction in time over ground transport.

<sup>&</sup>lt;sup>1</sup>Nurses who provide care during an interhospital transfer with a Specialty Care Licensed ambulance service shall comply with COMAR 10.27.09.04. The transport nurse shall have competency with the ongoing care of the patient being transferred and either a national transport certification or two years of critical care experience within the past five years.

## **Time to Intervention Care**

Patients may need rapid evaluation and treatment that are not available at the sending hospital. Reducing the time to clinical intervention is clinically important, and air medical transport may be utilized in order to minimize the time to intervention. Examples of the type of patient that would fit in this category may include, but are not limited to:

- Trauma patients being transferred for emergent diagnosis and operative intervention
- STEMI patients being transferred for emergency intervention
- Stroke patients being transferred for emergency intervention
- Intracranial hemorrhage patients being transferred for emergent diagnosis and intervention
- Patients being transferred for aortic catastrophe (leaking aneurysm, dissection, or disruption)
- Patients with tenuous airways unable to be secured at the sending hospital
- Patients with unstable blood pressure
- Patients with unstable cardiac rhythm
- Patients with severe sepsis requiring tertiary care
- Perinatal patients with eclampsia

# Reduction of "Out-of-Hospital Time" or Provision of Appropriate Level of Care En Route

Frequently, patients transferred to tertiary care hospitals will require a high level of care; for these patients it is critical to minimize the time they are away from the hospital setting. As well, it may be that a referring hospital may not have timely access to SCT ground ambulances that would be able to provide the appropriate level of care. Examples of the type of patient that would fit into this category may include, but are not limited to:

- Patients with tenuous airways unable to be secured at the sending hospital
- Patients with unstable blood pressure or on vasopressor medication
- Patients with unstable cardiac rhythm or on cardioactive drips
- Patients on mechanical ventilation with poor oxygenation
- Patients on intraaortic balloon pumps (IABPs)
- Patients with ventricular assist devices (VADs)
- Neonatal and critical pediatric patients
- Perinatal patients receiving tocolysis
- Other unstable patients where time out-of-hospital needs to be minimized

# **Commercial Air Medical Transport Clinicians**

MIEMSS-licensed commercial helicopters are staffed and equipped to provide primarily interhospital critical care transport to a tertiary care facility.

You may contact the State Office of Commercial Licensing and Regulation at 410-706-8511 or <u>SOCALR@miemss.org</u> for the most current list or for information about arranging a commercial ground or air transport.

A current list of commercial services is also available online at <a href="www.miemss.org">www.miemss.org</a>; navigate to Commercial Ambulance Licensing & Regulation (under Departments and Programs) and click on List of Licensed Commercial Services.

# **Maryland State Police Medevac Helicopters**

The primary medical role for MSP helicopters is direct scene response. They are available for interhospital transports at a Paramedic level, primarily as a back-up to the commercial helicopter programs. Critical Care interhospital helicopter transports are performed by the commercial programs. MSP back-up for Critical Care interhospital transports is typically dependent upon crew augmentation by a transport nurse or physician from the tertiary receiving hospital, which may or may not be available at the time of such request.

To access SYSCOM, call 1-800-648-3001. SYSCOM remains the point of contact to arrange MSP helicopter transportation.













Emergency Medical Dispatch: Skills include: Call intake/call allocator; medical interrogation; medical pre-arrival and post-dispatch instructions; medical call prioritization; EMS resource allocation and management. Hours of training: 24 hours minimum from an EMS Board-approved agency. Licensure requirements: written exam; successful completion of a health care provider CPR course. Licensure period is for two years. Renewal requirements: Current health care provider CPR card; 24 hours continuing dispatch education.

Emergency Medical Responder: Skills include: Patient assessment; vital signs; bleeding control and bandaging; fracture management; emergency medical management; CPR and AED; airway management and optional O2 administration; optional self and buddy WMD auto-injectors. Hours of training: 51 hours minimum. Certification requirements: Psychomotor/practical exam and National Registry EMR cognitive exam. Certification period is for three years. Renewal requirements: 12 hours of approved continuing education (6 hours didactic and 6 hours of skills). Maryland grants legal recognition for Emergency Medical Responders from most states, National Registry, and EMS Board-approved certifying agencies.

Emergency Medical Technician: Skills include: Patient assessment; vital signs; bleeding control and bandaging; shock management; fracture management; CPR, AED, O2 administration; airway management; emergency medical management; patient-assisted medications; spinal immobilization; patient movement; transport. Hours of training: 165 hours minimum. Certification requirements: EMS field internship; psychomotor/practical exam and National Registry EMT cognitive exam; affiliation with a BLS EMS Operational Program. Certification period is for three years. Renewal requirements: 24 hours of approved continuing education (12 hours didactic and 12 hours of skills); a current NREMT certification at the time of Maryland renewal may be used. Maryland accepts reciprocity for Emergency Medical Technicians from most states and National Registry.

Cardiac Rescue Technician: Skills include: All skills listed under EMT plus additional advanced procedures such as intravenous and intraosseous cannulation; medication administration; EKG monitoring; electrical therapy. Hours of training: EMT certification plus approximately 400 additional hours. Licensure requirements: Psychomotor/practical exam and National Registry I-99 cognitive exam; successful completion of a Maryland ALS licensing protocol exam; affiliation with an ALS EMS Operational Program. Licensure period is for two years. Renewal requirements: 60 hours of approved continuing education; continued affiliation with an ALS EMS Operational Program.

Paramedic: Skills include: All skills listed under CRT plus additional skills and medication administration. Hours of training: EMT certification plus approximately 1,100 additional hours. Licensure requirements: NRP certification; successful completion of a Maryland ALS licensing protocol exam; affiliation with an ALS EMS Operational Program. Licensure period is for two years. Renewal requirements: Successful re-registration as an NRP; continued affiliation with an ALS EMS Operational Program.

# **ADULT TRAUMA REFERRAL CENTERS**

# PRIMARY ADULT CLINICAL RESOURCE CENTER (PARC)

R Adams Cowley Shock Trauma Center
University of Maryland Medical Center
22 South Greene Street, Baltimore, MD 21201
Maryland Express Care 410-328-1234
1-800-373-4111

# **LEVEL I**

The Johns Hopkins Hospital
Adult Trauma Center
1800 Orleans Street
Baltimore, MD 21287
Hopkins Access Line 410-955-9444
1-800-765-5447

# LEVEL II

Johns Hopkins Bayview Medical Center 4940 Eastern Avenue

4940 Eastern Avenue Baltimore, MD 21224 ine 410-955-9444

Hopkins Access Line 410-955-9444 1-800-765-5447

University of Maryland Prince George's

Hospital Center
3001 Hospital Drive
Cheverly, MD 20785-1189
301-618-3752

Sinai Hospital of Baltimore 2401 W. Belvedere Avenue Baltimore, MD 21215

Lifelink 410-601-5465 410-601-6161

Suburban Hospital - JHM 8600 Old Georgetown Road Bethesda, MD 20814 301-896-3880

# LEVEL III

University of Pittsburgh Medical Center (UPMC) Western Maryland 12500 Willowbrook Rd. Cumberland, MD 21502

240-964-7000

TidalHealth Peninsula Regional Medical Center 100 East Carroll Street Salisbury, Maryland 21801-5493

> 410-543-4722 866-614-4722

Meritus Medical Center 11116 Medical Campus Rd. Hagerstown, MD 21742 301-790-8000


# **TRAUMA CENTERS**

# **Out-of State\***

Christiana Hospital
4755 Ogletown-Stanton Road
Newark, DE 19718
1-800-428-0911
302-733-5555

MedStar Washington Hospital Center
110 Irving St, NW
Washington, DC 20010
MedStar Transport Center
202-877-7234
1-844-877-2424

<sup>\*</sup> These facilities are American College of Surgeons Level I Trauma Centers that each have a Memorandum of Understanding with the State of Maryland for trauma care.



# **ADULT TRAUMA CENTER GUIDELINES FOR TRANSFER**

Patients with severe multiple system injury from any location in the state are candidates for referral to one of the nine Maryland adult trauma centers or two out-of-state trauma centers. Transfer patients to the appropriate level trauma center based on specialty medical care needs and resources required for patients' injuries.

## INDICATIONS FOR TRANSFER

Adults with one or more of the following:

- A. Severe multiple injuries (two or more systems) or severe single system injury
- B. Cardiac or major vessel injuries
- C. Injuries with complications (e.g., shock, sepsis, respiratory failure, cardiac failure)
- D. Severe facial injuries
- E. Severe orthopaedic injuries
- F. Co-morbid factors (e.g., age > 55 years, cardiac or respiratory disease, insulin-dependent diabetes, morbid obesity)

# **ADULT BURN INJURY CENTERS**

Johns Hopkins Bayview Medical Center 4940 Eastern Avenue, Baltimore, MD 21224 Hopkins Access Line (HAL): 410-955-9444 1-800-765-5447 MedStar Washington Hospital Center 110 Irving St, NW, Washington, DC 20010 MedStar Transport Center: 1-844-877-2424 202-877-7234

# **BURN INJURY**

# INTRODUCTION

The decision about where to transport a burned patient is based on location of the patient and location of available beds.

The Johns Hopkins Burn Center for Adults is located at Johns Hopkins Bayview Medical Center in eastern Baltimore City.

The **Adult Burn Center at MedStar Washington Hospital Center** in the District of Columbia also participates in the Maryland Specialty Referral System.

#### BURN CENTER REFERRAL CRITERIA

- 1. Partial thickness burns greater than 10% total body surface area (TBSA) in all adult age groups
- 2. Burns that involve the face, hands, feet, genitalia, perineum, or major joints
- 3. Third-degree (full thickness) burns of any size in all adult age groups
- 4. Electrical burns, including lightning injuries
- 5. Chemical burn injuries
- 6. Inhalation injuries
- 7. Burn injuries in patients with pre-existing medical conditions that may complicate management, prolong recovery, or affect mortality
- 8. Any patient with burns and coinciding trauma in which the burn injury poses the greatest risk of morbidity and mortality
- 9. Burned children in hospitals without qualified personnel or equipment to care for children (transport to a Pediatric Burn Center; see page 62)
- 10. Burned patients who require special social, emotional, or long-term burn rehabilitation

**Note:** Adult Burn Centers receive patients who have reached their 15th birthday. Pediatric Burn Centers receive patients who have NOT reached their 15th birthday (see page 64).

# **EYE TRAUMA CENTER**

The Wilmer Eye Institute at
The Johns Hopkins Hospital
1800 Orleans Street, Baltimore, MD 21287
Hopkins Access Line (HAL): 410-955-9444
1-800-765-5447

# **EYE TRAUMA**

#### INTRODUCTON

The Wilmer Eye Institute at the Johns Hopkins Hospital serves as an Eye Trauma Center for patients who have sustained ophthalmic trauma.

#### REFERRAL CONTACT

For patients who are suspected of having an eye/ophthalmic injury in the setting of multiple system trauma, call the Hopkins Access Line (HAL) and ask for the Pediatric or Adult Trauma Clinician On-call, who will evaluate the patient for acceptance and assist to facilitate the transfer of care to the Eye Trauma Center at the Johns Hopkins Hospital.

For pediatric or adult patients with an isolated eye/ophthalmic injury, call the HAL number and ask for the Ophthalmology Resident On-call, who will evaluate the patient for acceptance and assist to facilitate the transfer of care to the Eye Trauma Center at the Johns Hopkins Hospital.

If any question or concerns should arise during the referral process, please call the Hopkins Access Line at 410-955-9444 to contact the Medical Director.

NOTE: Patients are to be directed to either the Pediatric Emergency Department or the Adult Emergency Department unless otherwise told to access another care area.

## INDICATIONS FOR EYE TRAUMA TRANSFER

- 1. Serious ophthalmic injury, including but not limited to:
  - A. Open globe (penetrating or rupture)
  - B. Chemical burns of the eye
  - C. Periorbital trauma
  - D. Intraocular foreign bodies (foreign material inside the eye, not on the surface)
- 2. Individualized consultations are available for any other eye injuries.
- 3. Patients with isolated eye injuries, who are medically stable; eye trauma patients with multisystem injury who require involvement of the adult or pediatric trauma teams to determine the appropriateness for transfer


NOTE: Patients with other significant trauma should be transported to the appropriate facility for stabilization before transfer to an Eye Trauma Center.

# STABILIZATION PROCEDURES/PREPARATION FOR TRANSPORT

- 1. Protect eye with a rigid eye shield ONLY.
- 2. DO NOT remove impaled objects or attempt to clean the eye or eyelids. Stabilize penetrating objects in place.
- 3. Chemical injuries should receive continuous irrigation (if strong alkaline or acid, attempt to determine initial pH of the eye):
  - A. Water, sterile water, or normal saline
  - B. Send specimen of chemical with patient.
- 4. Keep patient NPO.

# TRANSPORT PATIENT with:

- 1. Copy of medical record
  - A. Treatment rendered (including medications)
  - B. Laboratory and X-ray results if available
    - (1) Send copies of X-rays and CT scans, not reports, if obtained prior to transport.
    - (2) DO NOT delay transport awaiting results.
- 2. Eye shield
- 3. Specimens of chemical agent, if indicated



# **CENTER FOR HYPERBARIC AND DIVE MEDICINE**

R Adams Cowley Shock Trauma Center University of Maryland Medical Center 22 South Greene Street, Baltimore, MD 21201 Maryland Express Care: 410-328-1234 1-800-373-4111

# **HYPERBARIC TRANSFER GUIDELINES**

# DECOMPRESSION SICKNESS/AIR EMBOLUS GAS GANGRENE/SOFT TISSUE INFECTIONS SMOKE INHALATION/CARBON MONOXIDE POISONING

The Center for Hyperbaric Medicine at the R Adams Cowley Shock Trauma Center has the only multi-place chamber in the State of Maryland, accommodating up to 23 patients per dive. The hyperbaric chamber is staffed by a team of specially trained critical care nurses, physicians, and respiratory therapists enabling them to provide care to critically ill patients 24/7.

## INDICATIONS FOR TRANSFER

- Suspected decompression sickness\*
- · Compromised/crush injury
- · Diving Accidents\*
- Suspected air embolus\*
- Suspected gas gangrene/soft tissue infection (following consultation with Soft Tissue MD or designee)
- Central Retinal Artery Occlusion (CRAO)
- Other indications may be appropriately transferred after consultation with hyperbaric physician or designee

# Presence of any one symptom in smoke inhalation/CO poisoning:

- Loss of consciousness
- Change in mental status (e.g., confusion, stupor, combativeness)
- Carboxyhemoglobin level of 25% or higher, measured transcutaneously or by blood levels
- Symptoms suggestive of cardiac ischemia (e.g., chest pain, ST segment changes)
- Pregnancy

<sup>\*</sup> If air transport is chosen, helicopters must fly under 1,000 feet

# HAND AND UPPER EXTREMITY TRAUMA CENTER

MedStar Union Memoral Hospital 3333 North Calvert Street, Baltimore, MD 21218 Handline: 410-261-8100 1-855-540-HAND (4263)

# HAND AND UPPER EXTREMITY TRAUMA

#### INTRODUCTION

The Curtis National Hand Center at MedStar Union Memorial Hospital in Baltimore serves as a specialty referral center for patients experiencing hand and upper extremity trauma.

#### INDICATIONS FOR REFERRAL TO A HAND CENTER

- 1. Fractures and dislocations of the hand, wrist, forearm, and elbow (open and closed, with or without neurovascular compromise)
- 2. Complex lacerations or tissue loss (with or without nerve or tendon involvement)
- 3. Amputations (complete or partial from mid-humerus distally)
- 4. Thermal injuries isolated to the hand and upper extremity (burns and frostbite)
- 5. High-pressure injection injuries
- 6. Selected infections (complex suppurative processes at and below the level of the carpus)
- 7. Compartment syndrome of the forearm and hand
- 8. Nerve and vessel injuries of the arm
- 9. Crush or degloving injuries and other trauma resulting in loss or perfusion or suspected nerve injury

#### CONTRAINDICATIONS FOR TRANSFER

- 1. Patients with major and/or multiple system trauma
- 2. Patients with unstable or abnormal vital signs
- 3. Lower extremity amputation; lower extremity amputations should be directed to Pediatric or Adult Trauma Center: Patient may exhibit injuries to skeletal or soft tissue components with complete or incomplete amputation of ankle/foot lower extremity, complicated nerve, vessel, or compartment syndrome. Toe amputation (partial or complete).

## STABILIZATION PROCEDURES/PREPARATION FOR TRANSPORT

- 1. Total patient assessment
  - a. Assess for evidence of other trauma. (The Hand Trauma Center is not a multi-system trauma facility. It accepts only patients with isolated extremity trauma or extremity trauma with other minor injuries.) If the patient is stable, follow emergency care instructions below while consultation and preparation for transport are accomplished.

#### 2. Emergency care


- a. DO NOT wash, rinse, scrub, or apply antiseptic to extremity. Apply dry sterile dressing, wrap in Kling or Kerlix, apply pressure, elevate, and cool.
- b. DO NOT wash, rinse, scrub, or apply antiseptic solution to the severed part
  - Wrap in dry sterile gauze or towel (depending on size). Package amputated extremity in sealed plastic bag and place ON TOP OF coolant bags or sealed bag of ice in a container (Styrofoam).

# DO NOT FREEZE.

- ii. THE AMPUTATED PART MUST NOT BE SUBMERGED IN ICE WATER. If the ice melts, replace it with another bag of ice.
- c. For partial amputation:
  - i. Place severed part(s) in a functional position.
  - ii. Apply dry sterile dressing.
  - iii. Splint.
  - iv. Elevate extremity.
  - v. Apply coolant bags or ice bag to the outside of the dressing.
- d. If possible, control bleeding with pressure. If tourniquet is necessary, place it close to the amputation site.
- e. Consider appropriate pain medication.

#### TRANSPORT PATIENT WITH:

- 1. Copy of medical record including:
  - a. X-ray and laboratory results
    - DO NOT delay transport while awaiting results. X-rays and blood work can be obtained upon arrival to the Hand Center.
  - b. Documentation of medications given:
    - i. Tetanus prophylaxis
    - ii. Antibiotics
    - iii. Pain medications
- 2. Extremity and/or part:
  - a. Elevated and cooled
  - b. Splints, as necessary



# **NEUROTRAUMA CENTER**

R Adams Cowley Shock Trauma Center University of Maryland Medical Center 22 South Greene Street, Baltimore, MD 21201 Maryland Express Care: 410-328-1234 1-800-373-4111

# **NEUROTRAUMA TRANSFER GUIDELINES**

# **Head and Spinal Injuries**

#### INTRODUCTION

As the state's designated referral center for head and spinal injuries, the multidisciplinary team of clinical experts at the Neurotrauma Center at the R Adams Cowley Shock Trauma Center utilizes evidence-based treatment strategies to care for patients with traumatic brain injuries and spinal column and spinal cord injuries.

Those patients with severe brain injury receive a multisystem assessment with intracranial pressure and cerebral oxygenation parameters closely monitored so that factors that may cause secondary brain injury are rapidly recognized and treated, thus optimizing patient outcomes. Neurosurgeons are readily available to intervene if necessary and perform craniotomies for hematoma evacuation and gunshot wound debridement, elevation of depressed skull fractures, decompressive craniectomies, and cranioplasties. Surgical interventions for spinal column injuries include discectomies, laminectomies, arthrodesis, and open reduction internal fixations.

This section provides guidelines for the stabilization and transport of patients with head and spine injuries. Patients who are under 15 years of age should be transported to a pediatric trauma center.

#### INDICATIONS FOR HEAD INJURY TRANSFER


#### Presence of any one symptom below:

- 1. Patients with deterioration in level of consciousness
- 2. Patients with severe head injuries (Glasgow Coma Score ≤ 8)
- 3. Patients with focal or lateralizing signs such as hemiparesis
- 4. Patients with penetrating cranial injury, including gunshot wounds or depressed skull fractures
- 5. Patients with cerebrospinal fluid leak: rhinorrhea or otorrhea
- 6. Seizures within 48 hours of trauma
- 7. Inability to perform immediate rapid neurosurgical pre-operative studies, intracranial monitoring, or neurosurgical operation that is or is likely to be necessary in management of the patient
- 8. Patients with moderate head injuries who may require other procedures or prolonged anesthesia (Glasgow Coma Scale scores of 9 to 12-13)

## INDICATIONS FOR SPINE INJURY TRANSFER

# Presence of any one symptom below:

- 1. Adult spinal cord injuries
- 2. Patients with suspected spinal injury whose level of consciousness is deteriorating
- 3. Patients with possible spinal fracture or dislocations that are unstable or need stability evaluation
- 4. Patients with neurological deficits
- 5. Patients with penetrating spinal injury, including gunshot or stab wounds
- 6. Patients with documented stable or unstable spinal column injuries with or without neurologic deficit
- 7. Inability to rapidly reduce fractures compressing the spinal cord by closed and/or surgical techniques



# SPECIAL PATHOGEN/HIGHLY INFECTIOUS DISEASE ASSESSMENT AND TREATMENT NOTIFICATIONS

The potential transfer of a patient with a special pathogen should be coordinated with the assistance of the local health department and the Maryland Department of Health (MDH). When a patient with a special pathogen is identified, the first external point of contact should be the local health department. If unable to reach the local health department, notify MDH via one of the numbers below.

MARYLAND DEPARTMENT OF HEALTH
Infectious Disease Epidemiology and
Outbreak Response Bureau (IDEORB)
201 W. Preston St, Baltimore, MD 21201
410-767-6700 (daytime)
410-795-7365 (after hours)

MARYLAND PRIMARY EBOLA/SPECIAL PATHOGEN TREATMENT HOSPITAL

**Johns Hopkins Hospital** 

Biocontainment Unit

1800 Orleans Street, Baltimore, MD 21287

MARYLAND PRIMARY EBOLA/SPECIAL PATHOGEN TREATMENT HOSPITAL

University of Maryland Medical Center 22 S. Greene Street, Baltimore, MD 21201

MARYLAND PRIMARY EBOLA/SPECIAL PATHOGEN ASSESSMENT HOSPITALS

Anne Arundel Medical Center 2001 Medical Parkway

Annapolis, MD 21401

Frederick Health Hospital 400 West Seventh Street Frederick, MD 21707

Holy Cross Hospital 1500 Forest Glen Rd. Silver Spring, MD 20910

MedStar Southern Maryland Hospital Center 7503 Surratts Rd.

Clinton, MD 20735

TidalHealth Peninsula Regional Medical Center 100 E. Carroll St. Salisbury, MD 21801

# SPECIAL PATHOGEN/HIGHLY INFECTIOUS DISEASE ASSESSMENT AND INTERHOSPITAL TRANSFER

The Maryland Department of Health (MDH) has developed a three-tier hospital system for patients with symptoms of highly infectious special pathogens\*. These three tiers are frontline hospitals, assessment hospitals, and treatment centers.

Hospitals receiving a patient with symptoms of a special pathogen should do the following:

- Identify the signs and symptoms
- Isolate the patient and utilize appropriate personal protective equipment
- Inform notify your local health department (if unable to reach the local health department, contact the MDH epidemiologist on call at 410-767-6700 (during business hours) or 410-795-7365 after hours, weekends, and holidays).

The local health department and MDH will assist with pathogen identification and with coordinating transfer to an appropriate destination if the patient requires an interhospital transfer. MDH will notify MIEMSS to coordinate the appropriate transport unit for the patient.


Below is a brief description of the three-tier hospital criteria and roles and responsibilities of each:

Frontline hospitals: All hospitals that are not Assessment or Treatment Hospitals are considered Frontline Hospitals. These facilities are expected to promptly identify and isolate a Patient Under Investigation (PUI) for a special pathogen and to promptly inform the health department. Frontline hospitals are expected to screen and isolate the patient and provide care for 12-24 hours until a special pathogen can be ruled out, or a patient transfer can be arranged. All frontline hospitals must have the capability to support EMS clinicians that transport a PUI to their facility including: an appropriate location for doffing of PPE, shower facilities for decontamination of EMS clinicians, an area to decontaminate transport vehicles, and acceptance of waste associated with the transport of PUIs.

Assessment hospitals: These hospitals are facilities that are prepared to receive and isolate PUIs and to care for the patient until a diagnosis of a special pathogen can be confirmed or ruled out, or until discharge or transfer is completed (if special pathogen diagnosis is confirmed). These hospitals must have the ability to coordinate Ebola Virus Disease (EVD) and other special pathogen testing and be able to provide appropriate care for up to 96 hours (i.e., have sufficient staff training, PPE, and isolation facilities appropriate for 4-5 days of patient care). All assessment hospitals must have the capability to support EMS transport personnel as listed under frontline hospitals.

**Treatment Centers:** These centers are hospitals which have been assessed by CDC for EVD and special pathogen readiness and are prepared to care for and manage a patient with confirmed EVD for the duration of the patient's illness. At minimum, these hospitals need sufficient staff, PPE, and isolation facilities appropriate for at least 7 days of patient care. All treatment centers must have the capability to support EMS transport personnel as listed above.

\*Special pathogens include: emerging and/or highly infectious diseases such as Ebola, Lassa Fever, MERS, category A agents, BSL 4 pathogens, etc. When in doubt, contact your local health department for guidance (if unable to reach the local health department, contact the MDH epidemiologist on call at 410-767-6700 (during business hours) or 410-795-7365 after hours, weekends, and holidays).



### STROKE GUIDELINES FOR TRANSFER

### INTRODUCTION

Development of a statewide system for stroke care in Maryland includes three (3) levels of hospital care. Each designated hospital level of care is based on accessibility and availability of timely health care resources. The development of designated levels of care and inter-facility transfer guidelines will maximize quality of care, ensure patient safety, and promote effective use of health care resources.

- (1) Level I Comprehensive Stroke Center (CSC);
- (2) Level II Primary Stroke Center (PSC); and
- (3) Level III Acute Stroke Ready Center (ASRC)
- I. Level I Comprehensive Stroke Center: Defined as a facility with immediate 24-hour access to the necessary personnel, infrastructure, equipment, expertise, and programs to rapidly diagnose and treat stroke patients who require a high intensity of medical and surgical care, specialized tests, or interventional therapies.

### Timely Transfer of Patients to a Comprehensive Stroke Center:

Because of the potential for rapid clinical deterioration, patients who present with severe disease or who have the high likelihood of clinical deterioration should be considered for timely transfer to a CSC. The transfer process should be initiated as early as possible in the patient's course. Remember, "time is brain."

### Refer to Stabilization and Preparation for Transport Section on page 36

### Circumstances to consider for a transfer to a Comprehensive Stroke Center

- Non-traumatic Subarachnoid Hemorrhage
- Intracerebral Hemorrhage
  - Hemispheral /Supratentorial: (> 30 mL or > 3cm)
  - Cerebellar hemorrhage
  - Brain Stem hemorrhage
  - Intraventricular hemorrhage
  - Suspected underlying lesion by imaging (e.g., CTA reveals a possible AVM)
- s/p IV t-PA with Concerns
  - Potential for malignant cerebral edema
  - Potential need or benefit from intra-arterial recanalization interventions
  - Perceived higher risk for symptomatic intracranial hemorrhage (e.g., difficult to control hypertension, malignant hypertension)

- Consideration for Hemicraniectomy
  - Age dependent (considered especially for age less than 60)
  - Baseline Modified Rankin of 0 or 1 or baseline independent in activities of daily living (ADLs)
  - Potential for malignant cerebral edema (e.g., high NIH stroke scale)
- Consideration for Endovascular Recanalization Treatment (ERT) (Interventional Neuroradiology)

Consultation with tertiary facility initiated as soon as possible

There is a goal time of 90 minutes from arrival of the patient at the PSC to departure to the CSC or PSC with EVT capability.

Goal from acceptance of the patient for transfer by the CSC or PSC with EVT capability to arrival at the CSC or PSC with EVT capability is 90 minutes.

- A. Thrombectomy for M1 and ICA occlusions are of proven benefit and recommended by guidelines. Patients should be considered candidates for neurointervention if imaging demonstrates a large artery occlusion and:
  - a) Less than 4.5 hours from time of onset (time last known to be at neurologic baseline)
    - NIH stroke score greater than or equal to 6;
    - ASPECTS score greater than or equal to 6; and
    - functionally independent at baseline
  - b) Greater than 4.5 hours up to 20 hours from time of onset (time last known to be at neurologic baseline) and can be received in the tertiary facility by 22 hours of last known well (LKW).
    - ASPECTS score 8-10 or core measurements < 30 ml from CT Perfusion or MRI;
    - NIH stroke score greater than or equal to 6; and
    - functionally independent at baseline
- B. Other large artery occlusion patients can be considered based on the above criteria, however, though it is not a proven indication. However, it may be reasonable in certain select patients although the benefit is uncertain:
  - Basilar Artery
  - M2
  - M3
  - PCA
  - ACA

### Special Circumstances to consider transfer to a Comprehensive Stroke Center

- Pregnancy associated stroke
- Pediatric stroke (under age 18) refer to a MIEMSS designated Pediatric Trauma Center
- Any circumstance for which there is a perceived need for higher level of care
- Young adult (18-45 years of age) with ischemic stroke
- Large cerebellar infarct and anticipation for surgical decompression
- History of Sickle Cell Anemia
- II. Level II Primary Stroke Center: Defined as a facility with the immediate availability of necessary personnel, infrastructure, equipment, expertise and programs to rapidly diagnose, treat, and either admit the patient or transfer the acute stroke patient. Level II Primary Stroke Centers may be able to take referrals for some but not all conditions managed at a Comprehensive Stroke Center (e.g., Subarachnoid Hemorrhage).

### Circumstances to consider for a transfer to a Comprehensive Stroke Center

• Meeting circumstances as identified in §A of this guideline

### Circumstances to consider for keeping patient at a Primary Stroke Center

- Intracerebral Hemorrhage
  - a) Small volume (less than 30 ml or < 3cm)
  - b) No cerebellar / brain stem involvement
  - c) No intraventricular hemorrhage
  - d) An alert patient
  - e) No suspicion of an underlying lesion such as AVM / Aneurysm
- Patients not meeting circumstances as identified in §A under Comprehensive Center
- Discussion of goals of care is recommended when making decisions to transfer to a Comprehensive Stroke Center
  - Situations in which further interventions might be considered futile
  - Patients with advanced co-morbid disease
  - Patients with poor baseline level of independent function
  - Patients identified as DNR
- s/p IV t-PA without special concerns
- Unruptured and Asymptomatic Cerebral Aneurysm (consider outpatient clinic referral)

III. Level III - Acute Stroke Ready Center: Defined as a facility with limited access to the necessary personnel, infrastructure, equipment, expertise, and programs to treat the acute stroke patient. The Acute Stroke Ready Center does possess the means to deliver emergent stroke therapies and transfer the acute stroke patient to a Primary or Comprehensive Stroke Center based on the patient's immediate needs.

### Circumstances to consider for transfer to a Comprehensive Stroke Center

• Meeting circumstances as identified in §A of this guideline

### Circumstances to consider for transfer to a Primary Stroke Center

- Limited or no cranial neurosurgery coverage for patients s/p IV t-PA or hemorrhage
- s/p IV t-PA
- Intracerebral hemorrhage
  - Small volume (less than 30 ml or < 3 cm)
  - An alert patient
  - No midline shift

### **IV. Non-Stroke Centers:**

### Circumstances to consider for transfer to a Comprehensive Stroke Center

• Meeting circumstances as identified in §A of this guideline

### Circumstances to consider transfer to a Primary Stroke Center

- Limited or no cranial neurosurgery coverage
- Intracerebral Hemorrhage
  - Small volume (less than 30 ml or < 3cm)
  - An alert patient
  - No midline shift
- s/p IV t-PA

### STABILIZATION AND PREPARATION FOR TRANSPORT

 Upon identification of a patient who may require transfer, immediately contact the potential receiving Primary Stroke Center, Thrombectomy-capable Primary Stroke Center, or Comprehensive Stroke Center based on the above criteria. A list of centers and contact information is included on pages 39-45 of this manual.

Once transfer is recommended, the consulting facility will reply to the sending facility, within 15 minutes, whether or not a bed will be available for transfer. If a bed is not available, the consulting facility will advise the sending facility to contact an alternative Comprehensive Stroke Center, Thrombectomy-capable Primary Stroke Center, or Primary Stroke Center.

- 2. The transferring physician is responsible for contacting the accepting hospital and secur-ing an accepting physician at the receiving facility.
- 3. The accepting physician will determine the transfer location (e.g., directly to the unit, Interventional Radiology Lab, or the Emergency Department).

### ARRANGING FORTRANSPORTATION

When determining the mode of transport, the following factors should be considered:

1. How soon does the patient need to reach the referral center?

A complex stroke patient who might benefit from emergent neurosurgical or interventional neuroradiology treatment should have the transfer completed within 90 minutes of acceptance of patient at the Comprehensive Stroke Center.

Transfer times for all other cases will be determined by the receiving center based on the patient's diagnosis and clinical status.

The sending facility should inform the patient and family that the patient is being transferred for consideration for advanced treatment.


- However, upon arrival at the Comprehensive Stroke Center the patient will be re-evaluated.
- The appropriateness of advanced treatment will be determined by the receiving center after re-evaluation.
- Advanced treatment may include enrollment in a clinical trial as appropriate.
- 2. What are the weather/ground conditions that might inhibit air transport?
- 3. What are the transport times for ground versus air transport from the referring institution?
- 4. The transferring hospital physician should make an assessment as to whether the patient requires intubation for safe transport to the higher level of care.
- 5. Should a patient's clinical status change (for better or worse) prior to departure from the hospital, it is imperative that the transferring physician inform the receiving physician of the change in clinical status.
- 6. All reasonable efforts will be made to obtain a reliable cell phone number for the patient and for responsible family members.

The transportation decision should be made by the receiving physician in collaboration with the referring physician based on clinical judgment, with careful consideration given to the above questions. Please refer to the Transport Services section on page 4 for additional information on arranging transportation.

### TRANSPORT PATIENT WITH:

- Copy of Medical Record including treatment rendered;
- Signed consent to transfer patient to receiving facility;
- Documentation of medications given; and
- X-ray, neuroimaging, and laboratory results. Include a CD with any relevant imaging.

### DO NOT DELAY TRANSPORT WHILE AWAITING RESULT



### PRIMARY STROKE CENTERS

### **REGION I**

**UPMC Western Maryland** 

12500 Willowbrook Rd. Cumberland, MD 21502

Stroke center neurologist on call: 240-964-7000

### **REGION II**

Frederick Health Hospital

400 West Seventh Street Frederick, MD 21701

**Emergency Department:** 240-566-3500

**Meritus Medical Center** 

11116 Medical Campus Road Hagerstown, MD 21742

**MedStar Harbor Hospital** 

3001 S. Hanover St.

410-350-3200

Baltimore, MD 21225

**Emergency Department:** 301-790-8300

### **REGION III**

**Anne Arundel Medical Center** 

2001 Medical Parkway Annapolis, MD 21401 Patient Access 443-481-6818

Administrative Coordinators:

443-481-5909

**UM Baltimore Washington Medical Center** 

301 Hospital Dr. Glen Burnie, MD 21061

Stroke center physician on call:

410-787-4000

**UM Harford Memorial Hospital** 501 S. Union Avenue

Stroke Center neurologist on call:

Havre de Grace, MD 21078

443-843-5500

Administrative Coordinator:

443-843-6176

**Carroll Hospital Center** 

200 Memorial Ave. Westminster, MD 21157

Emergency Department -Stroke Physician:

410-871-6700

**Howard County General Hospital - JHM** 

5755 Cedar Lane

Columbia, MD 21044

**Emergency Department** 

Clinical Coordinator:

443-718-2100

MedStar Franklin Square Medical Center

9000 Franklin Square Dr. Baltimore, MD 21237

On Call Neurologist:

443-777-4663 (Mon-Fri until 4:30 pm)

or 443-777-7000 (to have paged)

**UM Midtown Campus** 827 Linden Ave.

Baltimore, MD 21201

Neurologist on call:

410-225-8000

**MedStar Good Samaritan Hospital** 

5601 Loch Raven Blvd. Baltimore, MD 21239

Stroke center physician on call:

443-444-8000

**Mercy Medical Center** 

301 St. Paul Place Baltimore, MD 21202

Stroke center neurologist on call:

410-332-9000

**Greater Baltimore Medical Center** 

6701 N. Charles St. Baltimore, MD 21204

Stroke center neurologist on call:

443-849-4145

**Northwest Hospital** 

5401 Old Court Road

Randallstown, MD 21133

**Emergency Department** 

Charge Nurse:

410-496-8709

### **REGION III (Continued)**

Sinai Hospital of Baltimore

2401 W. Belvedere Ave. Baltimore, MD 21215 Contact LifeLink: 410-601-5465

and press #2 for transfer

St. Agnes Ascension Health

900 S. Caton Ave. Baltimore, MD 21229

Stroke center neurologist on call: 667-234-600

UM St. Joseph Medical Center

7601 Osler Dr. Towson, MD 21204

Main Hospital: 410-337-1000 Emergency Dept.:

410-337-1226

**MedStar Union Memorial Hospital** 

201 E. University Parkway Baltimore, MD 21218

Stroke center physician on call: 410-554-2000

**UM Upper Chesapeake Medical Center** 

500 Upper Chesapeake Drive Bel Air, MD 21014

Emergency Dept.: 443-643-2000

Administrative Coordinator: 443-643-4099

### **REGION IV**

**Atlantic General Hospital** 

9733 Healthway Dr. Berlin, MD 21811

Stroke center physician on call:

410-641-1100

**UM Shore Medical Center at Easton** 

219 S. Washington St. Easton, MD 21601

Stroke center neurologist on call: 410-822-1000

TidalHealth Peninsula Regional Medical Center

100 E. Carroll St.

Salisbury, MD 21801 **Stroke center physician on call:** 

410-546-6400

ChristianaCare, Union Hospital

106 Bow St.

Elkton, MD 21921

House Supervisor/ Clinical Placement:

443-907-6136

### **REGION V**

Adventist HealthCare White Oak Medical Center

11890 Healing Way Silver Spring, MD 20904 **Emergency Department:** 301-240-5070

CalvertHealth Medical Center

100 Hospital Rd.
Prince Frederick, MD 20678
Emergency Department:
410-535-8344

**UM Charles Regional Medical Center** 

5 Garrett Ave. La Plata, MD 20640

Emergency Department: 301-609-4160

**Doctor's Community Hospital** 

8118 Good Luck Road Lanham, MD 20706

Emergency Department: 301-552-8665

**Holy Cross Germantown Hospital** 

19801 Observation Drive Germantown, MD 20876

Emergency Department: 301-557-6500

**Holy Cross Hospital** 

1500 Forest Glen Rd. Silver Spring, MD 20910 301-754-7205

Emergency Department: 301-754-7500

**MedStar Montgomery Medical Center** 

18101 Prince Philip Dr. Olney, MD 20832

Emergency Svcs.: 301-774-8900 Nsg. Coordinator: 301-774-8767

MedStar Southern Maryland Hospital Center

7503 Surratts Rd. Clinton, MD 20735

Admissions: 301-877-4290

ED: 301-877-4500

MedStar St. Mary's Hospital

25500 Point Lookout Rd. Leonardtown, MD 20650

Emergency Department: 301-475-6110

**UM Prince George's Hospital Center** 

3001 Hospital Drive. Cheverly, MD 20785

Emergency Department: 301-618-3162

Adventist HealthCare Shady Grove Medical Center

9901 Medical Center Dr.

Rockville, MD 20850


*Emergency Department:* 240-826-6596

Suburban Hospital - JHM

8600 Old Georgetown Rd. Bethesda, MD 20814

Stroke neurologist on call: 301-896-3100

# **Primary Stroke Centers**



### **COMPREHENSIVE STROKE CENTERS**

### **REGION III**

Johns Hopkins Bayview Medical Center

4940 Eastern Ave. Baltimore, MD 21224

Urgent admissions and transfers:

410-955-9444

**University of Maryland Medical Center** 

22 S. Greene St. Baltimore, MD 21201


Maryland Express Care: 410-328-1234

1-800-737-4111

The Johns Hopkins Hospital 1800 Orleans St.

Baltimore, MD 21287

Hopkins Access Line: 410-955-9444



### **ENDOVASCULAR CAPABLE CENTERS IN MARYLAND**

### **Johns Hopkins Bayview Medical Center**

4940 Eastern Ave. Baltimore, MD. 21224

Contact: call the HAL Line 410-955-9444 or

Toll-free 1-800-765-5447 and ask for the Stroke Attending

**Endovascular Availability: 24/7/365** 

### The Johns Hopkins Hospital

1800 Orleans St. Baltimore, MD 21287

Contact: call the HAL Line 410-955-9444 or Toll-free: 1-800-765-5447

**Endovascular Availability:** 24/7/365

### TidalHealth Peninsula Regional Medical Center

100 East Carroll St. Salisbury, MD. 21801

Contact: Radiology Charge Technician 410-543-7356 and they will contact

the Interventionist on-call

**Endovascular Availability: 24/7/365** 

### Sinai Hospital of Baltimore

2401 West Belvedere Ave. Baltimore, MD 21215

Contact: Lifelink 410-601-5465 and press #2 for stroke transfer

**Endovascular Availability: 24/7/365** 

### Suburban Hospital - JHM

8600 Old Georgetown Rd. Bethesda, MD 20814

Contact: call the HAL Line 410-955-9444 or Toll-free: 1-800-765-5447


**Endovascular Availability:** 24/7/365

### **University of Maryland Medical Center**

22 S. Greene St. Baltimore, MD 21201

Contact: Maryland Express Care 410-328-1234

**Endovascular Availability: 24/7/365** 



# ACUTE ISCHEMIC STROKE GUIDELINES FOR POTENTIAL ENDOVASCULAR RECANALIZATION THERAPY (ERT)

### **OVERVIEW**

- IV Alteplase (tPA) should be administered to acute ischemic stroke patients as soon as possible according to guidelines.
- Patients with a NIHSS score  $\geq 8$  should be considered for emergent endovascular recanalization therapy (ERT).
  - o There is evidence-based data supporting the benefit and safety of ERT for anterior circulation (carotid artery territory) acute ischemic stroke.
- If a patient is a potential candidate for ERT, contact an endovascular capable facility immediately to discuss patient management.

### **ELIGIBILITY CRITERIA**

- a) Age  $\geq 18$ 
  - a. Children with stroke symptoms who have not reached their 18th birthday shall be treated under the pediatric protocol. Consult with a local base station and a pediatric base station to arrange transport to a Maryland pediatric trauma center.
- b) Administer IV Alteplase per established guidelines as soon as possible— IV <u>Alteplase should NOT</u> be delayed for decisions about ERT (goal door to needle time is less than 60 minutes). <u>IV Alteplase does NOT preclude ERT</u>. IV Alteplase is the standard of care first-line treatment for patients within 4.5 hours of stroke onset.
- c) NIHSS score ≥ 8 or occlusion of large artery on vascular imaging such as CT Angiography or MRA.
- d) Non-contrast head CT without hemorrhage or hypodensity of greater than 1/3 of the MCA territory.
- e) Patients ineligible for IV Alteplase due to anticoagulant use or recent surgery can be considered for ERT on a case-by-case basis.
- f) Transfer procedures should be urgently initiated with a goal of patient arrival at the receiving facility within 6 hours from last seen well.
- g) Patients with basilar thrombosis/occlusion should be urgently considered if transfer can be initiated with a goal of patient arrival at the receiving facility within 12 hours from last seen well.

### **GENERAL COMMENTS**

- a) Discussion of goals of care is recommended in the following cases:
  - I. Situations in which further interventions might be considered futile:
    - i. Patients with advanced co-morbidities.
    - ii. Patients with poor baseline level of independent function.
    - iii. Patients identified as DNR.
- b) Patients not meeting the Eligibility Criteria will be considered on a case-by-case basis. For instance:
  - I. Time is greater than 6 hours from last seen well.
  - II. Contraindication to IV Alteplase other than established above or potential contraindication to ERT.

### COMMENTS ON TIMING AND TRANSFER STRATEGIES

- a) The decision about IV Alteplase should be independent of, and should not be delayed because of, decisions about ERT.
- b) Consent: Every attempt to identify family members to consent for transfer and advanced treatment should be made. The lack of an available person to provide consent should NOT preclude or delay discussion or transfer of a patient for ERT.
- c) If it is determined the patient is a candidate for transfer and evaluation for ERT then:
  - a. Contact with endovascular capable facility should be initiated as soon as possible without delaying administration of IV Alteplase.
  - b. Transfer should be initiated as soon as possible. There is no need to wait for the IV Alteplase infusion to be completed.
  - c. The sending facility should inform the patient and family that the patient is being transferred for consideration for advanced treatments, including ERT.
    - i. However, upon arrival at the endovascular-capable facility the patient will be re-evaluated to determine which management strategy is most appropriate.
    - ii. Advanced treatment may include enrollment in a clinical trial.

### TRANSFER RECOMMENDATIONS

For potential ERT patients, to decrease transfer time:

- a) Once transfer is accepted, set target time of 15 minutes for patient to be ready for transport.
- b) Avoid intubation if the patient is maintaining an airway, and is expected to maintain the airway during transport.
- c) Unless vascular imaging can be obtained immediately, and not delay transfer, avoid further brain or vessel imaging prior to transfer.
  - a. If vascular imaging is obtained, do not delay transfer while waiting for a Radiologist interpretation. Send images on a CD with the patient to the endovascular capable facility.
  - b. If time allows, for radiographic contrast allergic patients consider IV administration of Solumedrol 100 mg and Benadryl 50 mg.
  - c. <u>Do not delay transfer waiting for a discharge summary.</u> Sending facility should fax discharge summary to accepting facility as soon as stat discharge summary is completed.
  - d. Patients with wake-up stroke and patients whose clinical status is not included in these guidelines will be considered on a case-by-case basis.

THIS PAGE IS INTENTIONALLY BLANK.

### **CARDIAC INTERVENTIONAL CENTERS**

### **REGION I**

**UPMC Western Maryland** 

12500 Willowbrook Road Cumberland, MD 21502

ED transfer line: 240-964-1010

Fax: 240-964-1270

### **REGION II**

Frederick Health Hospital

400 W. 7th Street

Frederick, MD 21701

ED Charge RN: 240-566-3500

Fax: 240-566-3946

**Meritus Medical Center** 

11116 Medical Campus Road Hagerstown, MD 21742

Hospital Operator: 301-790-8000

Fax: 301-790-9437

### **REGION III**

**Anne Arundel Medical Center** 

2001 Medical Parkway Annapolis, MD 21401

STEMI Line 443-481-1122

Fax: 443-481-1299

The Johns Hopkins Hospital

1800 Orleans Street Baltimore, MD 21287

Hopkins Access Line: 410-955-9444

Fax: 1-877-884-8839

4940 Eastern Ave.

**UM Baltimore Washington Medical Center** 

301 Hospital Dr.

Glen Burnie, MD 21061

ED: 410-787-4312 Cath Lab: 410-787-4214

Fax: 410-595-1961

Hopkins Access Line: 410-955-9444

Johns Hopkins Bayview Medical Center

Fax: 410-550-7679

Baltimore, MD 21224

**Carroll Hospital Center** 

200 Memorial Avenue

Westminster, MD 21157

ED: 410-871-6700

ED Charge Nurse: 410-871-7686

Shift Coordinator: 410-871-6938

Fax: 410-871-7177

**St. Agnes Hospital** 900 S. Caton Avenue

Baltimore, MD 21229

Transfer Activation Line: 410-368-3480

Fax: 410-368-2009

MedStar Franklin Square Medical Center

9000 Franklin Square Drive

Rosedale, MD 21237

ED Charge Nurse 443-777-2712

Nursing Supervisor 443-777-2771

Fax: 443-777-7070

UM St. Joseph Medical Center

7601 Osler Drive

Towson, MD 21204

Cath Lab BAT line: 410-427-2170

Fax: 410-337-1118

**Howard County General Hospital - JHM** 

5755 Cedar Lane

Columbia, MD 21044

Hopkins Access Line: 410-955-9444 Fax:

410-740-7551

Sinai Hospital of Baltimore

2401 W. Belvedere Ave.

Baltimore, MD 21215

Heart Line: 1-800-900-HART (4278)

Fax: 410-601-6478

### **REGION III (continued)**

MedStar Union Memorial Hospital

201 E. University Parkway Baltimore, MD 21218

Heartline 410-554-2332 or 1-888-529-0200

Fax: 410-554-6544

**UM Upper Chesapeake Medical Center** 500 Upper Chesapeake Drive

Bel Air, MD 21014

Nursing Supervisor: 443-643-4099 ED Charge Nurse: 443-643-4042

Fax: 443-643-2019

### **University of Maryland Medical Center**

22 S. Greene Street (Adult ED located off of Lombard St.)

Baltimore, MD 21201

Maryland Express Care: 410-328-1234

Fax: 410-328-1717

### **REGION IV**

**Bayhealth Hospital** 

640 South State Street Dover, DE 19901

House Supervisor 302-744-7791

Fax: 302-744-6595

### **Christiana Hospital**

4755 Ogletown Stanton Road Newark, DE 19718

Transfer Telephone:

302-733-1430 or 302-733-5555, Access Center

Fax: 302-733-2108

### **UM Shore Medical Center - Easton**

219 South Washington St. Easton, MD 21601

Single-Call Access: 410-822-1000 #5433 (LIFE)

Fax: 410-820-8874

### TidalHealth Nanticoke

801 Middleford Road Seaford, DE 19973

302-629-6611, press 0 for operator Ask for cardiologist on call

Fax: 302-628-6320

### **TidalHealth Peninsula** Regional Medical Center

100 E. Carroll Street Salisbury, MD 21801 Peninsula Access Center Bed

Coordinators:

410-543-4722 Fax: 410-912-5757

### **REGION V**

### Adventist HealthCare White Oak **Medical Center**

11890 Healing Way Silver Spring, MD 20904

Patient Access: 1-866-684-8460

Fax: 301-240-6211

### **Holy Cross Hospital**

1500 Forest Glen Road Silver Spring, MD 20910-1484

Direct STEMI Line: 240-638-0300

ED Charge Nurse: 240-635-0301

Fax: 301-754-7504

### **UM Prince George's Hospital Center**

3001 Hospital Drive Cheverly, MD 20785

On-Call: 301-618-6000

Fax: 301-618-2987

### Adventist HealthCare Shady Grove Medical Center

9901 Medical Center Drive

Rockville, MD 20850

ED charge nurse and/or ED physician: 240-826-6596

Fax: 240-826-5206

### MedStar Southern Maryland Hospital Center

7503 Surratts Road Clinton, MD 20735

ED Physician: 1-866-724-3188

Fax: 301-877-4668

### Suburban Hospital - JHM

8600 Old Georgetown Road Bethesda, MD 20814

"BEAT" Line: 301-896-2328

Fax: 301-896-7195


### **MedStar Washington Hospital Center**

110 Irving Street, NW

Washington, DC 20010 MedStar Transfer Center: 1-800-824-6814

Fax: 202-877-7879

# Cardiac Interventional Centers



# ST-ELEVATION MYOCARDIAL INFARCTION (STEMI) GUIDELINES FOR TRANSFER

### INTRODUCTION

Time to reperfusion is one of the most important factors in the survival of STEMI patients. While thrombolytics may be beneficial if given within 30 minutes of ED arrival, there is a higher rate of complications than with primary percutaneous coronary intervention (pPCI), which is the treatment of choice if door to balloon times of less than 90 minutes can be achieved. Therefore, rapid identification and transfer of STEMI patients to Cardiac Interventional Centers where pPCI can be performed is critical to achieve optimal patient outcomes. When primary PCI is the treatment of choice, STEMI patients arriving at non-Cardiac Interventional Centers should be transferred out of the emergency department within 30 minutes of arrival.

### INDICATIONS FOR TRANSFER

Documented or suspected STEMI.

### STABILIZATION AND PREP FOR TRANSPORT:

- 1. Upon confirmation of a STEMI, immediately contact the receiving Cardiac Interventional Center. A list of centers and contact information starts on page 51 of this manual.
- 2. Notify commercial ground or air medical transport service that a STEMI patient needs to be transferred STAT. If a specialty care transport (SCT) unit and SCT paramedic or nurse is not available, a registered nurse or physician may need to accompany the patient to the Cardiac Interventional Center in an ALS-licensed unit depending on the condition of the patient and treatment that has been initiated at the transferring hospital. Please refer to the Transport Services section on page 4 for additional information on arranging transports.

### INFORMATION FOR CALL TO CIC CARDIOLOGIST:

Obtain the following history from patient or EMS if possible:

- Time of acute symptom onset
- Duration of pain
- Age and DNR status
- History of prior MI/stent/CABG/renal failure?
- CPR, intubation, or multiple defibrillations en route to ED?
- Send fax confirming STEMI EKG to CIC (see pages 51-52 for fax numbers)

3. Obtain labs and portable CXR <u>if time permits</u>. **Do not delay transport to obtain.** 


Consult with receiving CIC regarding administration of medications prior to transfer. Do not delay transport to administer medications unless otherwise advised to do so by the receiving CIC. AVOID ALL IVs/DRIPS IF POSSIBLE, BUT DO NOT COMPROMISE PATIENT CARE. DRIPS COMPLICATE AND MAY DELAY TRANSPORT.

- 4. All transport documentation and paperwork should be completed STAT:
  - Signed consent to transfer patient to other facility
  - Signed release of medical records form
  - Medical treatment and assessment forms and documentation
  - Data sheet with the following times:
    - o Time symptoms started
    - o Time of first qualifying ECG (prehospital or ED)
    - o Time patient arrived in ED
    - o Time of first call to CIC
    - o Time of administration of medications, if given
    - o Time patient left sending hospital

### TRANSPORT PATIENT WITH:

- Copy of Medical Record including treatment rendered (including medications) and lab and x-ray results if available
- Data Sheet
- Copy of EKG(s) obtained by EMS and/or ED documenting STEMI (fax in advance if possible)
- eMEDS®
- Prehospital Consultation/Interventions Radio Report Form

DO NOT DELAY TRANSPORT AWAITING LAB RESULTS OR OBTAINING TIMES. These may be faxed to the receiving CIC while patient is en route.



### **PEDIATRIC TRAUMA CENTERS**

The Johns Hopkins Hospital -Children's Center

1800 Orleans Street, Baltimore, MD 21287

HAL Line: 410-955-9444

Administrative Office: 410-614-1811


### **Children's National Medical Center**

111 Michigan Avenue, NW, Washington, DC 20010

ECIC: 202-476-5433

ECIC: 1-800-884-5433

Administrative Office: 202-476-6698



### PEDIATRIC TRAUMA CENTERS' GUIDELINES

Children who have not reached their 15<sup>th</sup> birthday should be transferred to a pediatric trauma center. There are two Level 1 Pediatric Trauma Centers serving the state of Maryland caring for children who have not reached their 15th birthday. Both the Johns Hopkins Children's Center and Children's National Medical Center are designated by MIEMSS as Pediatric Base Stations, Pediatric Trauma Centers, and Pediatric Burn Centers with specially trained physicians, nurses, and other health professionals, as well as specially adapted equipment, to meet the needs and problems unique to children and their families.

### The Johns Hopkins Children's Center, Baltimore, MD

ALS ground and air medical transport is available for children to be transferred to the hospital. The transport team is capable of performing invasive and noninvasive monitoring and is able to provide full ventilatory support for children. The **Hopkins Access Line (HAL)** provides telephone access to the pediatric transport team, pediatric critical care and emergency medicine physicians, pediatric trauma and burn service, and other subspecialty consultants. **To initiate a transport to the Johns Hopkins Children's Center, call the HAL at 410-955-9444.** 

### Children's National Medical Center, Washington, DC

ALS ground and air medical transport is available for children to be transferred to the hospital from other facilities. The transport team is capable of performing invasive and noninvasive monitoring and is able to provide full ventilatory support for children. The Emergency Communications Information Center (ECIC) allows for communications between hospitals and EMS agencies and access to the pediatric transport team, pediatric critical care and emergency medicine physicians, pediatric trauma and burn service, and other subspecialty consultants. To schedule a transport, call 1-800-884-5433 (pediatric transport team at the Children's National) and speak with the communication specialist who will connect you with an attending physician in the ED and trauma center.

It is recommended that you first attempt to contact the pediatric trauma center directly. If you are unable to contact the Johns Hopkins Children's Center through the HAL or the Children's National Medical Center through the ECIC, you may contact the EMRC at 1-800-492-3805 for assistance.

### REASONS FOR TRANSFER TO A PEDIATRIC TRAUMA CENTER

- 1. Trauma any of the following:
  - a. Multiple-system injury (two or more systems)
  - b. Blunt thoracic trauma
  - c. Blunt abdominal trauma
  - d. Penetrating wounds
    - (1) Head
    - (2) Chest
    - (3) Abdomen
    - (4) Extremity with neurovascular compromise
  - e. Cardiac or major vessel injury
  - f. Extremity injury
    - (1) Open fractures
    - (2) Major long bone fracture
    - (3) Neurovascular compromise
    - (4) Avulsion or amputation of upper or lower extremities
  - g. Massive maxillofacial trauma
  - h. Spinal injury with or without deficit
  - i. Estimated Injury Severity Score (ISS) greater than 13
  - j. Severe head injury
    - (1) Deteriorating Glascow Coma Scale (GCS) regardless of score (9-15)
    - (2) GCS less than or equal to 8
      - (3) Depressed skull fracture or open head injury
      - (4) CSF leak—otorrhea or rhinorrhea
      - (5) Focal or lateralizing signs
      - (6) Intracranial hemorrhage
- 2. Burns See list in Pediatric Burn Referral Center section on page 62.
- 3. Evidence of shock or respiratory compromise:
  - a. Hypotension
  - b. Hypoxia
  - c. Hypovolemia
  - d. Mottled, cold, pale extremities
  - e. Tachycardia
  - f. Thready pulse
  - g. Tachypnea
  - h. Decreased level of consciousness
  - i. Urine production less than 0.5 mL/kg/hr
  - j. Metabolic acidosis (pH less than 7.2)
- 4. Any seriously injured child who cannot be managed in the community hospital

### GENERAL GUIDELINES FOR TRANSPORT OF CHILDREN

NOTES: These steps are guidelines in the assessment and stabilization of a pediatric trauma patient. Not all of these steps need to be accomplished prior to transfer of a patient to a trauma center. Call the pediatric trauma center for consultation/transfer as early as possible after considering that a patient may need care in a trauma center.

For Newborn transports - refer to the Neonatal Transport Section on page 69.

- Children should receive 100% oxygen during transport unless contraindicated by preexisting condition.
- Children transported with an ETT will have both a gastric tube placed for decompression and exhaled CO2 monitoring along with a chest X-ray.
- Secure all lines and tubes.
- Children meeting trauma criteria should be immobilized with a collar and backboard appropriate for size.
- Children should be transported with a secured and patent IV/IO.
- Children receiving intravenous medications must have the IV rate regulated by an infusion pump.
- Children should be kept NPO.
- Children should have a baseline glucose taken and recorded.
- Children should be kept warm with blankets and heat; document initial and regular temperatures with vital signs.
- Transport service must be notified of the transport of any child with a potentially infectious disease.

When possible, the pediatric transport teams will bring one adult family member with the child; siblings will not be transported. Safety of all occupants of both ambulance and helicopter transports will be the primary consideration. Driving directions to the receiving hospital should be provided to the family, along with contact phone numbers.


### **PEDIATRIC BURN CENTERS**

The Johns Hopkins Hospital 1800 Orleans Street Baltimore, MD 21287 410-955-9444 (HAL line for referrals)

Administrative Office: 410-614-1811 for Pediatric Trauma and Burn Offices

Pediatric Burn Center
Children's National Medical Center
111 Michigan Avenue, NW
Washington, DC 20010
202-476-5433 or 1-800-884-5433
(ECIC for referrals and Pediatric Transport Team)

Administrative Office: 202-476-6698 for Pediatric Trauma and Burn Offices



## INDICATIONS FOR TRANSFER TO A PEDIATRIC BURN CENTER

- 1. Partial thickness burns greater than 10% total body surface area (TBSA) in all age groups
- 2. Burns that involve the face, hands, feet, genitalia, perineum, or major joints
- 3. Full thickness burns in any age group (formerly referred to as third-degree burns)
- 4. Electrical burns, including lightning injury
- 5. Chemical burns
- 6. Burns complicated by smoke inhalation
- 7. Circumferential burns
- 8. The two Pediatric Burn Centers are also Pediatric Trauma Centers; burns complicated by trauma should be transferred to these two centers
- 9. Burns in patients with serious pre-existing medical conditions
- 10. Burns with concern regarding intentionality of injury (child victimization)

When necessary, the referring physician, in consultation with one of the Pediatric Burn Centers, may request a Pediatric Transport Team from the receiving center. Both Pediatric Burn Centers have dedicated Pediatric Transport Teams with ground and air capabilities.

### **MARYLAND PERINATAL REFERRAL CENTERS**

### **University-Based Perinatal Referral Centers**

The Johns Hopkins Hospital 1800 Orleans St Baltimore, MD 21287

L&D: 410-955-5850 Fax: 410-614-7720 NICU: 410-955-5255 Fax: 410-614-8834 For Neonatal Transport: 1-888-540-6767

University of Maryland Medical Center 22 South Greene Street Baltimore, MD 21201-1595

L&D: 410-328-1234 Fax: 410-328-8622 NICU: 410-328-6716 Fax: 410-328-5273 For Neonatal Transport: 1-888-540-6767

### **Maryland Perinatal Referral Centers**

Anne Arundel Medical Center 2001 Medical Parkway Annapolis, MD 21401-2777

L&D: 443-481-6989 Fax: 443-481-6954 NICU: 443-481-6962 Fax: 443-481-6954

MedStar Franklin Square Medical Center 9000 Franklin Square Drive Baltimore, MD 21237-3998

L&D: 443-777-8149 Fax: 443-777-7447 NICU: 443-777-7050 <u>OR</u> 443-777-2096

Fax: 443-777-8409 OB In-House Attending: 443-777-2074

For Maternal Transportation: 844-777-2424

844-777-2424 Fax: 443-777-7447

Frederick Health Hospital 400 West Seventh Street Frederick, MD 21701 L&D: 240-566-3533

Fax: 240-566-3979 NICU: 240-566-3582 Fax: 240-566-3247

Greater Baltimore Medical

6701 North Charles Street Towson, MD 21204-6892 **L&D:** 443-849-2577

Fax: 443-849-3026 NICU: 410-849-2591

Fax: 443-849-3025

Holy Cross Hospital 1500 Forest Glen Road Silver Spring, MD 20910 L&D: 301-754-7591

Fax: 301-754-7594 NICU: 301-754-7600

Fax: 301-754-7604

Howard County General Hospital - JHM 5755 Cedar Lane

Columbia, MD 21044-2999 **L&D:** 410-740-7845

Fax: 410-740-7614 NICU: 410-740-7555 Fax: 410-740-7513

Johns Hopkins Bayview Medical Center 4940 Eastern Avenue Baltimore, MD 21224-1505

L&D: 410-550-0328 Fax: 410-550-1098 NICU: 410-550-0395

Fax: 410-550-0439
Mercy Medical Center

301 St. Paul Place Baltimore, MD 21202-2165 **L&D: 410-332-9325** 

Fax: 410-234-2591 NICU: 410-332-9568

Fax: 410-528-5320

Adventist HealthCare Shady Grove Medical Center 9901 Medical Center Drive Rockville, MD 20850-3395

L&D: 240-826-6386 Fax: 240-826-6200 NICU: 240-826-7495 240-826-5372

Sinai Hospital of Baltimore 2401 W. Belvedere Avenue Baltimore, MD 21215-5271

L&D: 410-601-5240 Fax: 410-601-6875 NICU: 410-601-6077 Fax: 410-601-5557

For Neonatal Transport: 410-601-5465 (LifeLink)

St. Agnes Hospital 900 Caton Avenue Baltimore, MD 21229-5299 **L&D: 667-234-2610** 


Fax: 667-234-3559 NICU: 667-234-2630 Fax: 667-234-3569

UM St. Joseph Medical Center 7601 Osler Drive Towson, MD 21204-7582

L&D: 410-337-1178 Fax: 410-337-1009 NICU: 410-337-1150 Fax: 410-337-1844

### **Out-of-State Neonatal Referral Centers**

Children's National Medical Center 111 Michigan Avenue, NW Washington, DC 20010 202-476-5433 1-800-884-5433



# PERINATAL REFERRAL: MATERNAL REFERRAL/CONSULTATION

### INTRODUCTION

The value of neonatal transport in reducing neonatal morbidity and mortality rates has been well documented in the medical literature. Current evidence supports the theory of maternal transport as a significant factor in the reduction of neonatal mortality rates.

### INDICATIONS FOR TRANSFER

1. Maternal status does not improve.

### Examples:

- Preterm labor
- Premature rupture of membranes (PROM)
- Hypertensive disorders
- Second trimester incompetent cervix
- Third trimester bleeding
- 2. Delivery will occur prior to 34 weeks of gestation.

### Examples:

- Preterm labor
- PROM
- 3. Newborn facilities are inadequate to support the infant should delivery occur within 24 hours.

### Examples:

- Suspected or known fetal anomalies
- Intrauterine growth retardation (IUGR)
- 4. The obstetrician or pediatrician feels that a mother, fetus, or newborn may require intensive care or special services available in the perinatal centers.

### Examples:

- Suspected or known fetal anomalies
- IUGR
- Pregnancies complicated by medical disorders, such as diabetes, cardiac disease, sickle cell disease, or thromboembolic disease

### HOW TO INITIATE A TRANSFER

The referring physician should:

- Contact any one of the perinatal referral centers, or
- If unable to make contact with the perinatal referral center, you may call the EMRC-to request a maternal transport and they will connect you with the institution of your choice. Toll-free: 1-800-492-3805

### ARRANGING FOR TRANSPORTATION

When determining the mode of transport, the following factors should be considered:

- 1. How soon does the patient need to reach the referral center?
- 2. What are the weather/ground conditions that might inhibit air transport?
- 3 What are the transport times for ground versus air transport from the referring institution?

The transportation decision should be made by the receiving physician in collaboration with the referring physician based on clinical judgment, with careful consideration given to the above questions.

For GROUND TRANSPORTATION, the referring hospital will arrange transportation through local or commercial ambulance services.

For AIR TRANSPORTATION IN MARYLAND, the receiving perinatal center will arrange air transport.

# PERINATAL REFERRAL: NEONATAL REFERRAL/CONSULTATION

#### INTRODUCTION

There are 15 Maryland hospitals with Neonatal Intensive Care Units (NICU) that are capable of caring for critically ill newborns. These same 15 hospitals also care for high-risk mothers. Two of these are university-based hospitals capable of caring for all types of newborns, including those requiring cardiac and complex surgical procedures as well as medical intensive care. The other 13 Maryland hospitals provide medical intensive care and some may accept certain types of surgical patients. There are at several other hospitals out-of-state with NICUs that may accept neonatal patients from Maryland hospitals within their surrounding geographic areas.

All Perinatal Referral Centers meet Level III or IV standards in their respective Neonatal Intensive Care Unit as set by MIEMSS. The Maryland Regional Neonatal Transport Program (MRNTP) transfers to hospitals associated with the Johns Hopkins Hospital or University of Maryland Medical System. In addition, Holy Cross Hospital, Sinai, and Shady Grove Adventist have their own neonatal transport teams.

### HOW TO INITIATE A TRANSFER

The referring physician should:

- · Contact any one of the hospital or transport programs directly, or
- If unable to make contact with the perinatal referral center, you may call EMRC to request a neonatal transport and they will connect you with the institution of your choice. Toll-free: 1-800-492-3805

When contacting the EMRC operator, be very clear that this is either a neonatal or maternal transport/consultation request so the call is appropriately handled.

### NEONATAL TRANSPORT

- Contact the desired Perinatal Referral Center to initiate the referral. Neonates with suspected cardiac or complex surgical problems should be referred to one of the university centers. Neonates with suspected surgical problems may also be referred to a perinatal referral center with surgical capabilities. The selection of the receiving Perinatal Referral Center must be in compliance with COMAR 30.08.12 (Guidelines for Levels of Perinatal Care).
- 2. The receiving Perinatal Referral Center, in consultation with the sending facility, will determine if ground or air transport is clinically most appropriate.

### **Ambulance Transports**

- The receiving Perinatal Referral Center is responsible for arranging appropriate ambulance transport in a timely manner.
- This transport must be carried out by a Maryland Licensed Neonatal Commercial Ambulance Service and in compliance with COMAR 30.09 (Commercial Ambulance Regulations).

### **Neonatal Assessment Prior to Transport**

- Assess/address the following
  - a. Airway positioning
  - b. Breathing bag mask ventilation if necessary place pulse ox and monitor
  - c. Circulation pink, warm
  - d. Sugar glucose >50
  - e. Temperature 36.5-37.5 axillary
  - f. Airway document respiratory rate (< 60) and heart rate (100-150 bpm) Pulse ox (92) every 15 minutes
  - g. Blood Pressure is the baby pink, well perfused, voiding and stooling
  - h. Any risk factors for sepsis
  - i. Parental disposition
  - j. Treat any of the above parameters based on the S.T.A.B.L.E. guidelines

### 2. Placement

Gestational age and weight - based on the perinatal guidelines

- a. Level I > 35 weeks
- b. Level II  $\geq$  32 weeks gestation and  $\geq$  1500 grams
- c. Level III & IV no gestational age or weight criterion

If born at a setting without Perinatal Services, needs to be taken to the closest appropriate facility with Peds/Perinatal Services. The Maryland Regional Neonatal Transport Program can be called to assist with the transport.

- 3. Items essential for transport
  - a. Maternal Care Records and Labor & Delivery medical record
  - b. Infant Care Records, nursery and Labor & Delivery records
  - c. Armband that matches the maternal armband
  - d. One tube each or cord blood and maternal blood, if available
  - e. Transport equipment as defined by COMAR 30.09.12.06 that is designed and developmentally appropriate for the age of the neonate.

### Commercial Air Medical Transport (See page 7)

### **Maryland State Police Helicopter Transports**


- 1. The attending pediatrician/neonatologist at the referring hospital should call the MRNTP at 1-888-540-6767 to request transport. The MRNTP transport nurse will contact the on-call neonatologist. In a three-way conversation, with the coordinating center monitoring and recording the call, the MRNTP nurse and neonatologist and the referring physician will determine whether air or ground transport is more appropriate.
- 2. The transport nurse will contact MSP to request a helicopter transport and to obtain an estimated time of arrival of the MSP helicopter at the MRNTP hospital to pick up the nurse and the air sled.

- 3. Approval for helicopter utilization will be made by the attending neonatologist on-call for the MRNTP based on the patient's need to get to a higher level of care in a time-critical manner. Patients must either have a need to receive a specific intervention quickly or be unstable, such that air transport is needed to minimize their out-of-hospital time:
  - a. Any disagreements regarding approval for the MSP helicopter utilization will be resolved via an immediate conference call between the requesting physician, the MRNTP neonatologist, and the State Aeromedical Director; and
  - b. Neonatal centers without helipads immediate accessible to the hospital (i.e., an intermediate ambulance transport is required) will be utilized only when no appropriate center with an accessible helipad is available.
- 4. Once the MRNTP agrees to provide the neonatal transport nurse for the mission, online medical direction, including direction to the neonatal transport nurse, will be overseen by the MRNTP neonatologist until the patient arrives at the receiving Perinatal Referral Center.

# TRANSPORT OF STABLE NEWBORNS FROM A PERINATAL CENTER TO A COMMUNITY OR CONVALESCENT HOSPITAL

The following refers to stable infants being transferred for convalescent care who do not need the same level of care during transport as newborns being transferred to a Perinatal Referral Center or those being transferred who still require intensive care.

- 1. These elective transports must be prearranged between the referring Perinatal Referral Center and the receiving hospitals.
- Transports must be carried out by a licensed Neonatal Ambulance Service and in compliance with COMAR 30.09. Neonatal Commercial Ambulance Services may also transport stable infants; however, since these patients do not need the same level of care, they may also be transported in an Advanced Life Support (ALS) Commercial Ambulance or Specialty Care Transport (SCT).
- 3. When an ALS Commercial Ambulance is utilized for these transports, it must have:
  - a. One neonatal transport incubator powered by internal batteries as well as by alternating current power. The incubator must be secured with litter fasteners that meet the U.S. General Services Administration standard for ambulance litter fasteners and anchorages;
  - b. A registered nurse with a current Neonatal Resuscitation Program (NRP) certification from the American Academy of Pediatrics must accompany the infant.



# **EMTALA**

# THE EMERGENCY MEDICAL TREATMENT AND ACTIVE LABOR ACT

## **EMTALA - EXPLANATION OF REGULATIONS**

The Emergency Medical Treatment and Active Labor Act (EMTALA) became effective in 1986 as a federal law as part of the Consolidated Omnibus Budget Reconciliation Act of 1985 (COBRA). The purpose of EMTALA is to ensure that hospitals that receive Medicare funding and maintain a "dedicated ED" assess and stabilize or transfer patients with an emergency medical condition without consideration of ability to pay. It is sometimes referred to as the "antidumping" law.

If an individual's emergency medical condition has not been stabilized prior to transferring the individual to another hospital, the sending hospital must comply with certain EMTALA requirements in order to make an appropriate transfer.<sup>1</sup>

At the outset, it is important to remember that duties imposed by EMTALA are in addition to traditional state law requirements that patients be transferred in accordance with the standard of medical care applicable in a given situation. State regulations regarding patient transfer between hospitals are set forth in COMAR 10.07.01.23. Furthermore, a patient must always be transferred under the conditions that a reasonably prudent physician of like skill and training would require.

The general principles of EMTALA, as of March 2019, are briefly outlined below. Legal counsel should be consulted and the full statutory and regulatory materials reviewed to understand how EMTALA might impact a particular situation at any given time.

### WHAT HOSPITALS DOES EMTALA APPLY TO?

Under regulations promulgated in 2003, EMTALA applies to hospitals that have a "dedicated emergency department," which is defined as "any department or facility of the hospital, regardless of whether it is located on or off the main hospital campus, that meets at least one of the following requirements:

- (1) It is licensed by the State in which it is located under applicable State law as an emergency room or ED;
- (2) It is held out to the public (by name, posted signs, advertising, or other means) as a place that provides care for emergency medical conditions on an urgent basis without requiring a previously scheduled appointment; or

<sup>1</sup> Appendix V, Interpretive Guidelines - Responsibilities of Medicare Participating Hospitals in Emergency Cases (Rev. 60, 07-16-10), Tag A-2409/C-2409, § 489.24 (e)(2)(i-iv). (Available online at <a href="http://www.cms.gov/Regulations-and-Guidance/Guidance/Manuals/downloads/som107ap\_v\_emerg.pdf">http://www.cms.gov/Regulations-and-Guidance/Manuals/downloads/som107ap\_v\_emerg.pdf</a>.)

(3) During the calendar year immediately preceding the calendar year in which a determination under this section is being made, based on a representative sample of patient visits that occurred during that calendar year, it provides at least one-third of all of its outpatient visits for the treatment of emergency medical conditions on an urgent basis without requiring a previously scheduled appointment.<sup>2</sup>

### WHERE DOES EMTALA APPLY?

EMTALA applies everywhere on "hospital property," but regulations promulgated in 2003 make somewhat of a distinction between individuals presenting at a dedicated ED and individuals presenting elsewhere on hospital property.

Hospital property is defined as the physical area immediately adjacent to the main buildings as well as other areas within 250 yards of the main buildings and any other areas determined on an individual case basis by the Centers for Medicare and Medicaid Services (CMS) regional office. Hospital property includes the hospital's parking lots, sidewalks, and driveways, plus certain facilities located off campus.<sup>3</sup>

Hospital property does not include "other areas or structures of the hospital's main building that are not part of the hospital, such as physician offices, rural health centers, skilled nursing facilities, or other entities that participate separately under Medicare or restaurants, shops, or other nonmedical facilities."

#### WHEN DOES EMTALA APPLY?

The medical screening examination/stabilization requirements of EMTALA apply when an individual presents to a "dedicated ED" and requests examination or treatment for a **medical condition.**<sup>5</sup>

For individuals presenting on hospital property at locations other than the dedicated ED, EMTALA applies when the individual requests examination or treatment for an **emergency medical condition** (as opposed to requesting examination or treatment for a **medical condition**).<sup>6</sup>

A request for examination or treatment is considered to exist if a prudent layperson observer would believe, based on the individual's appearance or behavior, that the individual needs examination or treatment for a medical condition (if in a dedicated ED) or needs emergency examination or treatment (if elsewhere on hospital property).<sup>7</sup>

<sup>2 42</sup> CFR § 489.24(b)

<sup>3</sup> Appendix V Interpretive Guidelines, above n.1, Tag 406, § 489.24 (a)(1)(i)

<sup>4 42</sup> CFR § 489.24(b)

<sup>5</sup> The statute speaks in terms of "comes to the emergency department." 14 USC §1395dd (a). The regulations refine that concept. 42 CFR § 489.24(b) (definition of "comes to the emergency department")

<sup>6 42</sup> CFR § 489.24(b) (definition of "comes to the emergency department")

<sup>7</sup> Ibid.

In general, once an ambulance is on hospital property, EMTALA applies if the hospital is subject to EMTALA.<sup>8</sup>

However, EMS entry on hospital property with a patient to rendezvous with an air medical transport does not trigger EMTALA if the hospital is not the recipient hospital, unless a request is made by EMS personnel, the patient, or a legally responsible person acting on the individual's behalf for the examination or treatment of an emergency medical condition.

EMTALA also applies when an individual is in a ground or air ambulance owned or operated by a hospital to which EMTALA applies even if the ambulance is not on hospital property unless the ambulance is operated under community-wide EMS protocols directing transport locations.<sup>10</sup>

EMTALA does not apply to patients who have begun to receive outpatient services as part of an encounter, 11 and EMTALA does not apply to an individual admitted as an inpatient. Inpatient means an individual who is admitted to a hospital (including the ED) for bed occupancy for purposes of receiving inpatient hospital services as described in 42 CFR § 409.10(a) with the expectation that he or she will remain at least overnight and occupy a bed even though the situation later develops that the individual can be discharged or transferred to another hospital and does not actually use a hospital bed overnight. 12

Receiving hospitals are required to report violations of the EMTALA transfer provisions to CMS.<sup>13</sup> Failure to do so is itself a violation of EMTALA.

#### **SCREENING EXAMINATION**

When EMTALA applies, the hospital must provide a screening examination to the individual who has presented. The examination must be within the hospital's capabilities and conducted by individuals determined qualified in the hospital's by-laws or rules and regulations and who meet the emergency services requirements of hospitals participating in Medicare.

"Depending on the individual's presenting symptoms, the ... [medical screening examination] represents a spectrum ranging from a simple process involving only a brief history and physical examination to a complex process that also involves performing ancillary studies and procedures such as (but not limited to) lumbar punctures, clinical laboratory tests, CT scans, and/or diagnostic tests and procedures." The medical screening examination is an ongoing process, not an isolated event.

"Triaging is not equivalent to a medical screening examination. Triaging merely determines the 'order' in which patients will be seen, not the presence or absence of an emergency medical condition." <sup>15</sup>

<sup>8</sup> Ibid.

<sup>9</sup> Appendix V Interpretive Guidelines, above n.1, Tag A-2406/C-2411, § 489.24 (a)(1)(i)

<sup>10 42</sup> CFR § 489.24(b).

<sup>11</sup> Encounter means a direct personal contact between a patient and a physician, or other person who is authorized by State licensure law and, if applicable, by hospital or ... [Critical Access Hospital] staff bylaws, to order or furnish hospital services for diagnosis or treatment of the patient. 42 CFR 410.2; Torretti v. Main Line Hosps., Inc., 580 F.3d 168, 174-176 (3d Cir. 2009)

<sup>12 42</sup> CFR § 489.24

<sup>13 42</sup> CFR § 489.20(m)

<sup>14</sup> Appendix V Interpretive Guidelines, above n.1, Tag A-2406/C-2406, § 489.24 (a)(1)(i).

<sup>15</sup> Ibid.

### ABILITY TO PAY MUST NOT INTERFERE

At no time should any effort be made to determine the patient's ability to pay for or cover by insurance the costs of the EMTALA requirements. CMS and the Office of Inspector General advise that the hospital should employ properly trained staff members to respond to patient inquiries about costs in an effort to make certain the patient realizes the extent to which EMTALA procedures are available without cost. <sup>16</sup>

However, "It is not impermissible under EMTALA for a hospital to follow normal registration procedures for individuals who come to the ED. For example, a hospital may ask the individual for an insurance card, so long as doing so does not delay the medical screening examination. In addition, the hospital may seek other information (not payment) from the individual's health plan about the individual such as medical history. And, in the case of an individual with an emergency medical condition, once the hospital has conducted the medical screening examination and has initiated stabilizing treatment, it may seek authorization for all services from the plan, again, as long as doing so does not delay the implementation of the required ...[medical screening examination] and stabilizing treatment. A hospital that is not in a managed care plan's network of designated clinicians cannot refuse to screen and treat (or appropriately transfer, if the medical benefits of the transfer outweigh the risks or if the individual requests the transfer) individuals who are enrolled in the plan who come to the hospital if that hospital participates in the Medicare program." <sup>17</sup>

### EMTALA STABILIZATION REQUIREMENT

If it is determined that an emergency medical condition exists, either by means of a screening examination or otherwise, the hospital must either provide treatment within the capabilities of the staff and facilities available at the hospital to stabilize the condition or transfer the patient to another medical facility which can and has agreed to provide appropriate care.

If a patient refuses treatment or transfer, EMTALA provides specific requirements for documenting the circumstances of a refusal and the fact that the patient was properly informed of the risks and benefits. Samples of such documentation follow. Before any forms are implemented, the proposed procedure for using such forms should be reviewed with counsel to ensure appropriateness in a given situation.

<sup>16 64</sup> Fed. Reg. 217 at 61358 (1999)

<sup>17</sup> Appendix V Interpretive Guidelines, above n.1, Tag A-2406/C-2406, § 489.24 (a)(1)(i)

<sup>18</sup> See sample forms on pages 75-77.

### EMTALA PRECONDITIONS FOR A PATIENT TRANSFER

In general, if a hospital is aware that a patient is experiencing an emergency medical condition, the patient cannot be transferred until:

- A. The emergency medical condition has been stabilized as required under EMTALA; or
- B. The following conditions are met:
  - (1) The transfer is requested in writing by the patient<sup>19</sup> or a legally responsible person acting on the patient's behalf after being informed of the hospital's obligations under EMTALA. The request must state the reasons for the request and indicate that the person making the request is aware of the risks and benefits of the transfer; and
  - (2) A physician has signed a certification<sup>20</sup> that, based upon the information available at the time of the transfer, the medical benefits reasonably expected from the provision of appropriate medical treatment at another medical facility outweigh the increased risks to the individual or, in the case of a woman in labor, to the woman or the unborn child, from being transferred. The certification must contain a summary of the risks and benefits upon which it is based; or, if a physician is not physically present in the emergency department at the time an individual is transferred, a qualified medical person (as determined by the hospital in its by-laws or rules and regulations) has signed the certification after consultation with a physician who agrees with the certification and countersigns the certification, which contains a summary of the risks and benefits upon which it is based.

## EMTALA REQUIREMENTS FOR A PROPER PATIENT TRANSFER TO ANOTHER MEDICAL FACILITY

EMTALA requires that transfer of a patient who is not stabilized meets four requirements<sup>21</sup>:

- 1. The sending hospital must provide medical treatment within its capacity that minimizes the risks to the individual's health and, in the case of a woman in labor, the health of the unborn child;
- 2. The receiving facility must have available space and qualified personnel for the treatment of the individual and must have agreed to accept transfer of the individual and to provide appropriate medical treatment;
- 3. The sending hospital must send to the receiving facility all medical records (or copies) related to the emergency condition available at the time of transfer, including:
  - Available history;
  - Records related to the individual's emergency condition;
  - Observations of signs or symptoms;
  - Preliminary diagnosis;
  - Results of diagnostic studies or telephone reports of the studies;
  - Treatment provided;
  - Results of any tests;
  - The informed written consent or certification (or a copy) required for the transfer; and
  - Name and address of any on-call physician who has refused or failed to appear within a reasonable time to provide necessary stabilizing treatment.

<sup>19</sup> See sample form on page 76.

<sup>20</sup> See sample form on page 77.

<sup>21</sup> Appendix V Interpretive Guidelines, above n.1, Tag A-2409/C-2409, § 489.24(e)(2)(i-iv).

Other records, such as test results not yet available or historical records not readily available from the hospital's files, must be sent as soon as practicable after transfer; and

4. The transfer must be effected through qualified personnel and transportation equipment, as required, including the use of necessary and medically appropriate life support measures during the transfer.

Under EMTALA, EMS clinicians may not always be qualified to provide the level of care for certain patients being transferred. A patient's condition may make the presence of a physician or some other specialist mandatory. Under current CMS guidelines, the physician at the sending hospital (not at the receiving hospital)<sup>22</sup> has responsibility to determine the appropriate mode, equipment, and attendants for transfer.

A Medicare-participating hospital that has specialized capabilities or facilities (including burn units, shock-trauma units, and neonatal intensive care units) may not refuse to accept from a referring hospital an appropriate transfer of an individual who requires such specialized capabilities or facilities if the receiving hospital has the capacity to treat the individual. This requirement applies regardless of whether the hospital has a dedicated emergency department.<sup>23</sup>

Between 2008 and 2012, the CMS explored requiring a hospital with specialized capabilities to continue to have EMTALA obligations despite the fact that a patient had been admitted to another hospital. On February 2, 2012, CMS published notice that it would not change the EMTALA responsibilities of a hospital with specialized capacities, and that the EMTALA obligations of a hospital with specialized capabilities would cease with respect to a patient when the patient is admitted to another hospital.<sup>24</sup>

<sup>22</sup> Appendix V Interpretive Guidelines, above n.1, Tag 409 §489.24(e)(2)(iv)

<sup>23 42</sup> CFR § 24(f)

<sup>24 77</sup> FR 5217

### **ENFORCEMENT**

EMTALA is enforced by CMS and by the Department of Health and Human Services' Office of the Inspector General.

Investigations are based on complaints, and the limitations period is two years after the date of the violation with respect to which the action is brought.

Possible penalties for violations are:

Termination of a hospital's Medicare clinician agreement;

A maximum civil money penality for a hospital or responsible physician

- (a) Hospital less than 100 beds \$52,484
- (b) Hospital more than 100 beds \$106,965

The exclusion of a physician from Medicare and Medicaid programs;

Civil suit by a patient for damages; and

A suit by a receiving facility that suffered loss because of another hospital's violation of EMTALA.

## Sample Form. Review with counsel before using.

## Refusal of Examination, Treatment, or Transfer

I understand that [insert name of hospital] (Hospital) must provide me a medical screening examination to determine whether I have an emergency medical condition, and if I do, to either stabilize the condition or transfer me in an appropriate manner to another facility.

I further understand that the medical screening and stabilization or transfer in connection with an emergency condition must be performed by Hospital without regard for whether I am able to pay or whether I have insurance that will pay part or all of the costs of the examination, treatment, or transfer.

Hospital p	roposes to	perform	the	following	examination,	treatment,	or	transfer:

Hospital has informed me of the following risks and benefits of this proposed examination, treatment, or transfer:

I refuse the examination, treatment, or transfer set forth above for the following reasons:

I understand my refusal is against medical advice and that my refusal may result in serious harm to me, including death.

Date:	
Patient Signature:	
Patient Printed Name:	
Date of Birth:	
Address:	
Witness signature:	
Witness printed name:	
Witness address:	

# Sample Form. Review with counsel before using. Patient Request for Transfer

I understand that [insert name of hospital] (Hospital) must provide me a medical screening examination to determine whether I have an emergency medical condition, and if I do, to either stabilize the condition or transfer me in an appropriate manner to another facility.

I further understand that the medical screening and stabilization or transfer in connection with an emergency condition must be performed by Hospital without regard for whether I am able to pay or whether I have insurance that will pay part or all of the costs of the examination, treatment, and/or transfer.

eatment, and/or transfer.	mountained that will puly pairt of all of the ex	, sub of the chairmant.
I understand these obligati	ons of Hospital, and I request a transfer to	:
The reasons for my reques	et for a transfer are:	
Hospital has informed me	that the transfer that I request exposes me	to the following risks:
Date: Patient Signature: Patient Printed Name: Date of Birth: Address:		· · · ·
Witness signature: Witness printed name: Witness address:		· · ·

## Sample Form. Review with counsel before using.

# **Certification of Transfer**

Patient Name:
It is hereby certified that, based upon the information available at the time of transfer, the medical benefits to this patient reasonably expected from the provision of appropriate medical treatment at another medical facility outweigh the increased risks to the individual, and, in the case of labor, to the unborn child, from being transferred.
This certification is based on the following risks and benefits.
Risks:
Benefits:
Name of Certifying Physician: *
Signature of Certifying Physician:
Date:
*If a physician is not physically present in the emergency department at the time of transfer, a qualified medical person (as determined by the hospital in its by-laws or rules and regulations) must consult with a physician and sign the certification below. The physician mus subsequently countersign above:
Name of qualified medical person:
Signature of qualified medical person:
Name of physician consulted:
At (time) on (date)

### TRANSPORT OF STABLE PATIENTS FROM A TRAUMA/BURN OR SPECIALTY CENTER TO A COMMUNITY HOSPITAL OR REHABILITATION/LONG-TERM CARE FACILITY

The following refers to stable patients (pediatric and adult) being transferred for specialty or convalescent care who do not need to stay at the tertiary care centers and have non-emergent transport care needs. Examples include: back transfer to referring hospital closer to home, transfer to out-of-state hospital closer to patient's home, rehabilitation or long-term care, another acute care facility for consultation, or a psychiatric facility.

- 1. These elective transports must be prearranged between the referring Specialty Care Referral Center and the receiving hospital.
- 2. Transports must be carried out by a currently licensed Basic or Advanced Life Support Commercial Ambulance (see COMAR 30.09).
- 3. When a BLS or ALS Commercial Ambulance is utilized for these transports, it must have:
  - Stretcher that can be secured in accordance with the U.S. General Services Administration standard for ambulance AND the appropriate child restraint or five-point adult stretcher straps appropriate for the patient's height and weight;
  - Method to secure a family member in an appropriate seat belt if accompanying the patient;
  - Method to secure patient care equipment and personal items;
  - Equipment required by SOCALR licensure;
  - Additional equipment required for the specific needs of the patient being transferred.
- 4. The transferring attending physician will contact the attending physician at the receiving hospital to determine if the back transfer or transfer for additional care is appropriate based upon the patient's condition and will determine the composition of the transport team needed in transport. (Refer to INTERHOSPITAL TRANSFER CHECKLIST on pages 86-87 as a guideline.)
- 5. On the day of transfer, verbal patient care report will be given nurse to nurse from the sending facility to the receiving facility, providing updated information in addition to the written discharge summaries in the transfer records.

- 6. The sending and receiving facility discharge planners, case manager, and/or social workers will provide for the continuity of care to include:
  - a) Verify availability of patient care bed at the appropriate level of care
  - b) Time and date of transfer
  - c) Arrange for appropriate transport via a licensed Maryland BLS or ALS ambulance
  - d) Verify consent for transfer by the patient, parent, or guardian
  - e) Patient belongings to be transported by family and those needed in ambulance transport
- 7. Transfer summary of the patient's care and copies of pertinent part of the patient's chart to include but not limited to:
  - Physician orders for care during transfer
  - Primary family member and contact information (e.g., parents, guardian, spouse, significant other, legal guardian)
  - Medical history and history from acute admission
  - Patient primary care clinician prior to the acute admission, if known
  - Physician and Nursing discharge summaries
  - Discharge teaching provided to the patient and family
  - Results from most recent laboratory studies and diagnostic studies
  - Results from most recent radiology studies
  - Rehabilitation, PT, OT, speech progress notes, and plan of care

## Sample Form. Review with counsel before using.

## INTERHOSPITAL TRANSFER CHECKLIST

The reason for transfer: higher level of care for specialty care patient request directed by payor other (please specify)
Attending physician written order for transfer on chart yes no
Reason for transfer has been discussed with patient and/or family yes no
Consent for transfer has been signed by patient and/or responsible family member yes no
Medical screening exam provided by:
Attending physician has contacted receiving physician yes no
Name of accepting physician: Contact phone numbers:
Name of receiving hospital:
Report Given MD to MD: time Names: Names:
Mode of transport:  Specialty Care Ambulance  Neonatal Care Ambulance  Advanced Life Support Ambulance  Basic Life Support Ambulance  Ground Ambulance  Air Medical Transport  Other
Mode of transport: ambulance air medical transport private car
Level of care needed during transport BLS ALS RN MD Other: (please specify)
Equipment needed for support of patient during transport is available on transport unit yes no
Medications and IV fluids needed during transport are with patient yes no
Patient's airway and ventilation is being controlled with

are being provided to the receiving hospital at the time of the patient's arrival:
<ol> <li>Prehospital care record</li> <li>ED record of care</li> <li>Medical history, if available</li> <li>Results of laboratory studies and diagnostic studies</li> <li>Copies of radiographs</li> <li>Nursing care records, including I &amp; O documentation and vital signs</li> <li>Doctor's orders for care during transfer</li> <li>Transfer consent form</li> <li>Patient's belongings</li> <li>Family contact information/notification of transfer</li> </ol>
PROMPT TRANSPORT  Do not delay transport while awaiting laboratory or radiology results. These can be communicated by phone as they become available.  Name and number from referring hospital:

The following copies of the medical records related to the patient's emergency condition



Maryland Institute for Emergency Medical Services Systems