

2017 Benchmark Selection: Staff Recommendation

| Measure (page #) | 2015 Performance | 2015 Benchmark | 2016 Benchmark | 2017 Benchmark Staff Recommendation |
|---|---|---|---|--|
| Access to Care (CAHPS) (39) | State: 83.8% High CCO: 87.7% Low CCO: 78.1% | 87.2%, 2014 National Medicaid 75 th percentile; average of adult & child rates. MN method with 2 percentage point floor | 89.1%, 2015 national Medicaid 75 th percentile; weighted average of adult and child rates MN method with 2 percentage point floor | TBD 2016 National Medicaid 75 th percentile; weighted average of adult and child rates – pending NCQA MN method with 2 percentage point floor |
| Adolescent Well Care Visits (42) | State: 37.5% High CCO: 46.2% Low CCO: 22.3% | 62.0%, 2014 National Medicaid 75 th percentile (admin data) MN method with 3 percentage point floor | 61.9%, 2015 National Medicaid 75 th percentile (admin data) MN method with 3 percentage point floor | TBD 2016 National Medicaid 75 th percentile (admin data only) – pending NCQA MN method with 3 percentage point floor |
| Alcohol and substance misuse (SBIRT) (44) | State: 12.7% High CCO: 23.7% Low CCO: 3.3% | 12.0%, Committee consensus; weighted to include adolescents MN method with 3 percentage point floor | 12%, Committee consensus; weighted to include adolescents MN method with 3 percentage point floor | Increased percentage TBD Suggested 17.2% (90th percentile of 2015 CCOs) MN method with 3 percentage point floor |

| Measure (page #) | 2015 Performance | 2015 Benchmark | 2016 Benchmark | 2017 Benchmark Staff Recommendation |
|--|--|--|---|---|
| Ambulatory care: emergency department utilization (52) | State: 43.1 High CCO: 29.2 Low CCO: 61.8 <i>Lower is better</i> | 39.4/1,000 member months 2014 National Medicaid 90 th percentile MN method | 39.8/1,000 member months 2015 National Medicaid 90 th percentile MN method with 3 percent floor | TBD 2016 National Medicaid 90 th percentile – pending NCQA MN method with 3 percent floor |
| Assessments for children in DHS Custody (126) | State: 58.4% High CCO: 76.7% Low CCO: 40.3% | 90%, Committee consensus MN method with 3 percentage point floor | 90%, Committee consensus MN method with 3 percentage point floor | 90%, Committee consensus MN method with 3 percentage point floor |
| Childhood immunization status (69) | State: 70.7% High CCO: 83.5% Low CCO: 55.5% | N/A | 82.0% 2015 National Medicaid 75 th percentile MN method with 3 percentage point floor | TBD 2016 National Medicaid 75 th percentile – pending NCQA MN method with 3 percentage point floor |
| Cigarette smoking prevalence (163) | <i>Baseline data from EHRs not yet available, however CAHPS survey data indicate statewide prevalence of 30.1%, ranging from 39.2% - 21.7%</i> | N/A | 25%, Committee consensus N/A – benchmark only | 25%, Committee consensus MN method with 1 percentage point floor |

| Measure (page #) | 2015 Performance | 2015 Benchmark | 2016 Benchmark | 2017 Benchmark Staff Recommendation |
|----------------------------------|---|--|---|---|
| Colorectal cancer screening (73) | State: 46.6% High CCO: 52.7% Low CCO: 36.0% | 47%, Committee consensus MN method with 3 percentage point floor | 47%, Committee consensus MN method with 3 percentage point floor | Increased percentage TBD -90 th percentile of Oregon CCOs (50.8%) -National Commercial 75 th percentile (~66%) MN method with 3 percentage point floor |
| Controlling hypertension (79) | State: 64.7% High CCO: 71.3% Low CCO: 59.1% | 64.0%, 2014 National Medicaid 75 th percentile MN method with 3 percentage point floor | 69.0% 2015 National Medicaid 90 th percentile MN method with 3 percentage point floor | TBD 2016 National Medicaid 90 th percentile – pending NCCA MN method with 3 percentage point floor |
| Dental sealants (81) | State: 18.5% High CCO: 24.3% Low CCO: 11.3% | 20.0%, Committee consensus MN method with 3 percentage point floor | 20.0%, Committee consensus MN method with 3 percentage point floor | 20%, Committee consensus MN method with 3 percentage point floor |
| Depression screening (87) | State: 37.4% High CCO: 62.8% Low CCO: 0.5% | 25.0%, Committee consensus MN method with 3 percentage point floor. | 25.0%, Committee consensus MN method with 3 percentage point floor | Increased percentage TBD -75 th percentile of Oregon CCOs (52.9%) - MN method with 3 percentage point floor |

| Measure (page #) | 2015 Performance | 2015 Benchmark | 2016 Benchmark | 2017 Benchmark Staff Recommendation |
|--|---|--|--|---|
| Developmental screening (89) | State: 54.7% High CCO: 84.1% Low CCO: 36.2% | 50.0%, Committee consensus MN method | 50.0%, Committee consensus MN method with 3 percentage point floor | Increased percentage TBD -75 th percentile of Oregon CCOs (60.1%) - ? MN method with 3 percentage point floor |
| Diabetes: HbA1c poor control (91) | State: 26.7% High CCO: 11.4% Low CCO: 33.9% <i>Lower is better</i> | 34.0%, 2014 National Medicaid 75 th percentile MN method with 3 percentage point floor | 19.0%, 2015 National Commercial 90 th percentile MN method with 3 percentage point floor | 19.0%, 2015 National Commercial 90 th percentile MN method with 3 percentage point floor |
| Effective contraceptive use (95) | State: 36.3% High CCO: 45.4% Low CCO: 31.9% | 50.0%, Committee consensus MN method with 3 percentage point floor | 50.0%, Committee consensus MN method with 3 percentage point floor | 50.0%, Committee consensus MN method with 3 percentage point floor |
| Follow-up after hospitalization for mental illness (103) | State: 75.3% High CCO: 90.0% Low CCO: 60.0% | 70.0%, 2014 National Medicaid 90 th percentile MN method with 3 percentage point floor | 79.9%, 2014 CCO 90 th percentile MN method with 3 percentage point floor | 86.8%, 2015 CCO 90 th percentile MN method with 3 percentage point floor |

| Measure (page #) | 2015 Performance | 2015 Benchmark | 2016 Benchmark | 2017 Benchmark Staff Recommendation |
|---|---|---|---|---|
| Patient Centered Primary Care Home enrollment (129) | State: 87.5% High CCO: 99.9% Low CCO: 73.5% | N/A – 60% threshold | N/A – 60% threshold | N/A – 60% threshold |
| Satisfaction with care (CAHPS) (151) | State: 85.4% High CCO: 89.1% Low CCO: 80.2% | 89.6%, 2014 National Medicaid 75 th percentile, weighted average of adult and child rates MN method with 2 percentage point floor | 89.2%, 2015 National Medicaid 75 th percentile; weighted average of adult and child rates MN method with 2 percentage point floor | TBD, 2016 National Medicaid 75 th percentile; weighted average of adult and child rates – pending NCQA MN method with 2 percentage point floor |
| Timeliness of prenatal care (146) | State: 84.7% High CCO: 92.3% Low CCO: 72.3% | 90.0%, 2014 National Medicaid 75 th percentile MN method | 93.0%, 2015 National Medicaid 90 th percentile MN method with 3 percentage point floor | TBD, 2016 National Medicaid 90 th percentile – pending NCQA MN method with 3 percentage point floor |