

Percent of Oregon adults who reported engaging in heavy alcohol consumption in the
last month, by ethnicity/race
Behavioral Risk Factor Surveillance System (BRFSS), Oregon, 2010 - 2011

Ethnicity/Race	N	Unadjusted		Age-Adjusted	
		Est. (%)	SE	Est. (%)	SE
Non-Hispanic White	9,923	8.1	0.4	8.4	0.4
Non-Hispanic Black/African American	351	4.7†	2.1	4.5†	1.7
Non-Hispanic Asian/Pacific Islander	416	5.1†	2.7	4.7†	2.4
Non-Hispanic American Indian/Alaska Native	380	6.7†	2.4	6.8†	2.4
Hispanic	345	4.4†	1.8	4.1†	1.6
<i>TOTAL</i>	<i>11,415</i>	<i>7.6</i>	<i>0.4</i>	<i>7.7</i>	<i>0.4</i>

Source: Oregon BRFSS 2010-2011 Race Oversample
Number of respondents (N) is unweighted; percentages are weighted

See technical notes for calculations used to determine total N for each ethnicity/race

Age-Adjusted to standard U.S. 2000 population - 3 groups (18-34, 35-54, 55+)

* = an estimate that significantly differs from non-Hispanic Whites ($p < .05$); calculated for age-adjusted values only

† = an estimate with a relative standard error > 30%, and should be considered unreliable

Heavy drinking is defined as 2 or more alcoholic drinks per day in the past 30 days for men, and 1 or more alcoholic drinks per day in the past 30 days for women

Date: March 17, 2014