

Oregon Behavioral Risk Factor Surveillance System (BRFSS), 1996

Do you currently use any smokeless tobacco products such as chewing tobacco or snuff?
(Weighted Column Percents)

| | | Total | Age Groups | | | | | 65+ |
|----------------------|----------------------|-------|------------|-------|-------|-------|-------|------|
| | | | 18-24 | 25-34 | 35-44 | 45-54 | 55-64 | |
| Yes, chewing tobacco | | 2.8 | 8.5 | 5.5 | 3.0 | 1.3 | .9 | .2 |
| Yes, snuff | | .7 | .4 | .6 | 1.3 | .7 | .3 | .2 |
| Yes, both | | .3 | .8 | .2 | | .7 | | .2 |
| No, neither | | 96.0 | 90.3 | 93.8 | 95.0 | 97.3 | 98.8 | 99.3 |
| Refused | | .2 | | | .7 | | | .2 |
| Male | Yes, chewing tobacco | 6.7 | 17.8 | 13.1 | 7.1 | 2.7 | 2.2 | .5 |
| | Yes, snuff | 1.5 | .8 | 1.4 | 2.7 | 1.5 | .7 | .5 |
| | Yes, both | .6 | 1.7 | .5 | | 1.5 | | .5 |
| | No, neither | 90.9 | 79.7 | 85.1 | 88.8 | 94.3 | 97.1 | 98.5 |
| | Refused | .3 | | | 1.4 | | | |
| Female | Yes, chewing tobacco | .1 | .7 | | | | | |
| | Yes, snuff | .1 | | | .2 | | | |
| | No, neither | 99.8 | 99.3 | 100.0 | 99.5 | 100.0 | 100.0 | 99.7 |
| | Refused | .1 | | | .2 | | | .3 |

Denominator is: All survey respondents

Do you currently use any smokeless tobacco products such as chewing tobacco or snuff?
(Unweighted Frequencies)

| | | Total | Age Groups | | | | | 65+ |
|----------------------|----------------------|-------|------------|-------|-------|-------|-------|-----|
| | | | 18-24 | 25-34 | 35-44 | 45-54 | 55-64 | |
| Total | | 2915 | 258 | 529 | 698 | 546 | 327 | 557 |
| Yes, chewing tobacco | | 83 | 22 | 29 | 21 | 7 | 3 | 1 |
| Yes, snuff | | 19 | 1 | 3 | 9 | 4 | 1 | 1 |
| Yes, both | | 8 | 2 | 1 | | 4 | | 1 |
| No, neither | | 2799 | 233 | 496 | 663 | 531 | 323 | 553 |
| Refused | | 6 | | | 5 | | | 1 |
| Male | Total | 1233 | 118 | 222 | 294 | 262 | 136 | 201 |
| | Yes, chewing tobacco | 82 | 21 | 29 | 21 | 7 | 3 | 1 |
| | Yes, snuff | 18 | 1 | 3 | 8 | 4 | 1 | 1 |
| | Yes, both | 8 | 2 | 1 | | 4 | | 1 |
| | No, neither | 1121 | 94 | 189 | 261 | 247 | 132 | 198 |
| | Refused | 4 | | | 4 | | | |
| Female | Total | 1682 | 140 | 307 | 404 | 284 | 191 | 356 |
| | Yes, chewing tobacco | 1 | 1 | | | | | |
| | Yes, snuff | 1 | | | 1 | | | |
| | No, neither | 1678 | 139 | 307 | 402 | 284 | 191 | 355 |

| | | | | | |
|--|---|--|---|--|---|
| Refused | 2 | | 1 | | 1 |
| Denominator is: All survey respondents | | | | | |

Oregon Behavioral Risk Factor Surveillance System (BRFSS), 1996

On how many of the past 30 days did you use smokeless tobacco?
(Weighted Column Percents)

| | | Total | Age Groups | | | | | |
|---------------|-------------|-------|------------|-------|-------|-------|-------|------|
| | | | 18-24 | 25-34 | 35-44 | 45-54 | 55-64 | 65+ |
| 1 | | 2.7 | 12.0 | | | | | |
| 2 | | 4.5 | 4.0 | 6.1 | 3.3 | 6.7 | | |
| 3 | | .9 | 4.0 | | | | | |
| 4 | | 2.7 | 4.0 | 3.0 | 3.3 | | | |
| 5 | | 4.5 | 8.0 | 3.0 | 3.3 | 6.7 | | |
| 6 | | .9 | 4.0 | | | | | |
| 7 | | .9 | 4.0 | | | | | |
| 8 | | 1.8 | 4.0 | 3.0 | | | | |
| 10 | | .9 | | | | 6.7 | | |
| 15 | | 7.3 | 4.0 | 6.1 | 3.3 | 20.0 | | 33.3 |
| 20 | | 4.5 | 4.0 | 9.1 | 3.3 | | | |
| 25 | | 2.7 | 4.0 | 3.0 | 3.3 | | | |
| 28 | | 1.8 | | | 6.7 | | | |
| 29 | | .9 | | | 3.3 | | | |
| 30 | | 57.3 | 44.0 | 60.6 | 60.0 | 60.0 | 75.0 | 66.7 |
| None | | 5.5 | | 6.1 | 10.0 | | 25.0 | |
| Male | 1 | 1.9 | 8.3 | | | | | |
| | 2 | 4.6 | 4.2 | 6.1 | 3.4 | 6.7 | | |
| | 3 | .9 | 4.2 | | | | | |
| | 4 | 2.8 | 4.2 | 3.0 | 3.4 | | | |
| | 5 | 4.6 | 8.3 | 3.0 | 3.4 | 6.7 | | |
| | 6 | .9 | 4.2 | | | | | |
| | 7 | .9 | 4.2 | | | | | |
| | 8 | 1.9 | 4.2 | 3.0 | | | | |
| | 10 | .9 | | | | 6.7 | | |
| | 15 | 7.4 | 4.2 | 6.1 | 3.4 | 20.0 | | 33.3 |
| | 20 | 4.6 | 4.2 | 9.1 | 3.4 | | | |
| | 25 | 2.8 | 4.2 | 3.0 | 3.4 | | | |
| | 28 | 1.9 | | | | 6.9 | | |
| | 29 | .9 | | | | 3.4 | | |
| 30 | 58.3 | 45.8 | 60.6 | 62.1 | 60.0 | 75.0 | 66.7 | |
| None | 4.6 | | 6.1 | 6.9 | | 25.0 | | |
| Female | 1 | 50.0 | 100.0 | | | | | |
| | None | 50.0 | | | 100.0 | | | |

Denominator is: All survey respondents who use smokeless tobacco products

Oregon Behavioral Risk Factor Surveillance System (BRFSS), 1996

On how many of the past 30 days did you use smokeless tobacco?
(Unweighted Frequencies)

| | | Total | Age Groups | | | | | |
|--------------|--------------|-------|------------|-------|-------|-------|-------|-----|
| | | | 18-24 | 25-34 | 35-44 | 45-54 | 55-64 | 65+ |
| Total | | 110 | 25 | 33 | 30 | 15 | 4 | 3 |
| 1 | | 3 | 3 | | | | | |
| 2 | | 5 | 1 | 2 | 1 | 1 | | |
| 3 | | 1 | 1 | | | | | |
| 4 | | 3 | 1 | 1 | 1 | | | |
| 5 | | 5 | 2 | 1 | 1 | 1 | | |
| 6 | | 1 | 1 | | | | | |
| 7 | | 1 | 1 | | | | | |
| 8 | | 2 | 1 | 1 | | | | |
| 10 | | 1 | | | | 1 | | |
| 15 | | 8 | 1 | 2 | 1 | 3 | | 1 |
| 20 | | 5 | 1 | 3 | 1 | | | |
| 25 | | 3 | 1 | 1 | 1 | | | |
| 28 | | 2 | | | 2 | | | |
| 29 | | 1 | | | 1 | | | |
| 30 | | 63 | 11 | 20 | 18 | 9 | 3 | 2 |
| None | | 6 | | 2 | 3 | | 1 | |
| Male | Total | 108 | 24 | 33 | 29 | 15 | 4 | 3 |
| | 1 | 2 | 2 | | | | | |
| | 2 | 5 | 1 | 2 | 1 | 1 | | |
| | 3 | 1 | 1 | | | | | |
| | 4 | 3 | 1 | 1 | 1 | | | |
| | 5 | 5 | 2 | 1 | 1 | 1 | | |
| | 6 | 1 | 1 | | | | | |
| | 7 | 1 | 1 | | | | | |
| | 8 | 2 | 1 | 1 | | | | |
| | 10 | 1 | | | | 1 | | |
| | 15 | 8 | 1 | 2 | 1 | 3 | | 1 |
| | 20 | 5 | 1 | 3 | 1 | | | |
| | 25 | 3 | 1 | 1 | 1 | | | |
| | 28 | 2 | | | 2 | | | |
| | 29 | 1 | | | 1 | | | |
| | 30 | 63 | 11 | 20 | 18 | 9 | 3 | 2 |
| | None | 5 | | 2 | 2 | | 1 | |
| | Total | 2 | 1 | | 1 | | | |

| | | | | | | | |
|---|-------------|---|---|--|---|--|--|
| Female | 1 | 1 | 1 | | | | |
| | None | 1 | | | 1 | | |
| Denominator is: All survey respondents who use smokeless tobacco products | | | | | | | |

