	sted to the 2000 U.S. Census Population.
County	% who consumed at least 5 serving of fruits
	and vegetables per day
OREGON	25.8%
Baker	27.9%
Benton	30.1%
Clackamas	25.2%
Clatsop	22.6%
Columbia	26.0%
Coos	22.1%
Crook	23.9%
Curry	33.2%
Deschutes	29.5%
Douglas	19.5%**
Grant	18.0%†
Harney	22.3%†
Hood River	30.1%
Jackson	25.6%
Jefferson	32.5%
Josephine	23.3%
Klamath	25.8%
Lake	19.6%†
Lane	23.6%
Lincoln	23.5%
Linn	22.9%
Malheur	23.0%
Marion	24.6%
Morrow	13.1%
Multnomah	28.1%
Polk	25.0%
Tillamook	28.3%
Umatilla	20.3%
Union	21.6%
Wallowa	26.1%
Washington	29.8%**
Yamhill	23.5%
Gilliam/Wheeler	26.0%†
Sherman/Wasco	· · · · · · · · · · · · · · · · · · ·
<u> </u>	

Age-adjusted Prevalence of Selected Chronic Diseases and Related Risk Factors Among Adults by County, Oregon 2002-2005

Tables were created using combined BRFSS survey results from 2002 through 2005. Two numbers are provided for each condition or disease: an unadjusted rate and an age-adjusted rate.

Unadjusted rates, provided in separate tables, give a description of the absolute burden of a disease or risk factor for an individual county.

Age-adjusted rates supplied in this table allow you to compare rates for a county to the State's rate. They adjust for differences that could result from a county's population being, on average, older or younger than the State's. When comparing counties with Oregon or with each other, use these age-adjusted rates.

† Rate based on less than 50 respondents overall or fewer than 12 persons in one age group; may not accurately reflect behavior of entire county and results should be interpreted with caution. Estimates based on 10 or fewer respondents are not reported.

** Statistically significant difference compared to Oregon.

Three age groups were used for age adjusting: 18-34yrs., 35-54yrs, and 55+yrs.

Source: Oregon Behavioral Risk Factor Surveillance System - healthoregon.org/brfss