

Table I: Age-Adjusted and Unadjusted Prevalence of Selected Chronic Conditions among Adults, by County, Oregon 2006-2009

| County | Arthritis | | Asthma | | Heart Attack | | Angina | |
|-----------------------|------------|--------------|------------|--------------|--------------|--------------|------------|--------------|
| | unadjusted | age-adjusted | unadjusted | age-adjusted | unadjusted | age-adjusted | unadjusted | age-adjusted |
| OREGON | | 25.8% | | 9.7% | | 3.3% | | 3.4% |
| Baker | 31.6% | 26.5% | 7.9% | 6.9% | 3.5%† | 2.7%† | 6.2%† | 4.9% |
| Benton | 26.0% | 27.9% | 8.8% | 9.4% | 3.0% | 3.3% | 2.8% | 3.0% |
| Clackamas | 23.7% | 22.4% | 9.6% | 9.7% | 2.6% | 2.5% | 3.1% | 2.9% |
| Clatsop | 39.7% | 39.1%* | 7.9% | 7.5% | 4.8% | 3.9% | 4.0% | 3.1% |
| Columbia | 24.4% | 22.4% | 10.3% | 9.8% | 4.4%† | 4.1%† | 2.5%† | 2.1%† |
| Coos | 34.3% | 28.4% | 12.3% | 13.1% | 7.7%† | 7.3%† | 8.2% | 7.7%† |
| Crook | 25.2% | 22.4% | 5.7%† | 5.6%† | – | 2.0%† | – | – |
| Curry | 31.5% | 34.1%† | 11.7% | 12.6% | 8.4% | 4.4% | 10.1% | 6.0%* |
| Deschutes | 29.9% | 28.0% | 9.0% | 9.2% | 3.0% | 2.7% | 2.8% | 2.4% |
| Douglas | 40.8% | 36.8%* | 12.6% | 12.7% | 5.7% | 4.4% | 5.2% | 4.0% |
| Grant | 18.3%† | 11.2%† | 12.0%† | 13.6%† | 4.2%† | 2.9%† | 3.6%† | 2.5%† |
| Harney | 30.6%† | 26.8%† | 12.9%† | 13.3%† | 7.5%† | 6.0%† | – | – |
| Hood River | 25.2%† | 24.2% | 2.8%† | – | 6.2%† | 5.8%† | 3.9%† | 3.6%† |
| Jackson | 28.6% | 25.4% | 8.5% | 8.3% | 3.5% | 2.9% | 3.5% | 2.9% |
| Jefferson | 48.0% | 47.5% | 7.1%† | 6.6%† | 3.7%† | 3.0%† | 3.5%† | 2.8%† |
| Josephine | 31.2% | 26.4% | 11.3% | 11.6% | 4.9% | 3.4% | 5.7% | 4.0% |
| Klamath | 36.2% | 32.5% | 11.3% | 11.5% | 6.9% | 6.0%* | 4.7% | 3.7% |
| Lake | 29.8%† | 25.2%† | 6.3%† | 7.2%† | – | – | – | – |
| Lane | 28.3% | 27.1% | 10.5% | 10.6% | 3.4% | 3.1% | 3.8% | 3.5% |
| Lincoln | 31.8% | 27.9% | 10.7% | 11.2% | 5.6% | 3.9% | 3.4% | 2.5%† |
| Linn | 32.6% | 29.5% | 10.4% | 10.5% | 5.4% | 4.5% | 5.0% | 4.2% |
| Malheur | 27.3% | 27.1% | 8.4% | 8.4% | 3.7%† | 3.4%† | 2.0%† | 1.7%† |
| Marion | 25.1% | 25.0% | 11.4% | 11.3% | 4.1% | 4.0% | 3.9% | 3.8% |
| Morrow | 27.9%† | 24.3%† | 7.9%† | 8.2%† | – | – | – | 6.5%† |
| Multnomah | 23.9% | 24.3% | 9.2% | 9.2% | 2.8% | 2.9% | 2.9% | 3.0% |
| Polk | 26.5% | 25.6% | 9.1% | 9.1% | 5.3% | 5.2% | 4.7% | 4.4% |
| Tillamook | 32.9% | 28.7% | 8.1%† | 8.2%† | 5.8%† | 4.8%† | 4.8%† | 3.9%† |
| Umatilla | 28.9% | 27.8% | 7.6% | 7.5% | 3.4% | 3.0% | 3.7% | 3.4% |
| Union | 31.9% | 31.0% | 12.8% | 13.3% | 4.3%† | 4.0%† | 6.2% | 5.3% |
| Wallowa | 45.8% | 38.2% | 8.0%† | 6.2%† | – | – | 8.7%† | 5.4%† |
| Washington | 22.2% | 23.1% | 9.0% | 9.0% | 2.4% | 2.5% | 2.9% | 3.1% |
| Wheeler | – | – | – | – | – | – | – | – |
| Yamhill | 24.6% | 24.9% | 11.5% | 11.5% | 3.2% | 3.1% | 3.2% | 3.2% |
| Gilliam/Sherman/Wasco | 21.7% | 17.5% | 5.9% | 6.5%† | 6.0% | 4.6% | 6.3% | 5.0% |

* Statistically significant difference compared with Oregon.

† This number may be statistically unreliable and should be interpreted with caution.

– This number is suppressed because it is statistically unreliable.

Age-adjusted estimates are adjusted to the 2000 Standard Population using three age groups (18-34, 35-54, and 55+).

Source: Oregon BRFSS County Combined Dataset 2006-2009

Table I (continued): Age-Adjusted and Unadjusted Prevalence of Selected Chronic Conditions among Adults, by County, Oregon 2006-2009

| County | Stroke | | Diabetes | | High Blood Pressure | | High Blood Cholesterol | |
|-----------------------|------------|--------------|------------|--------------|---------------------|--------------|------------------------|--------------|
| | unadjusted | age-adjusted | unadjusted | age-adjusted | unadjusted | age-adjusted | unadjusted | age-adjusted |
| OREGON | | 2.3% | | 6.8% | | 25.8% | | 33.0% |
| Baker | 3.1%† | 2.4%† | 12.8% | 11.4%* | 39.0% | 31.3% | 56.2% | 59.9%* |
| Benton | 1.5% | 1.6% | 5.5% | 6.3% | 20.3% | 22.9% | 28.4% | 25.8% |
| Clackamas | 2.8% | 2.6% | 7.0% | 6.6% | 26.8% | 25.6% | 37.6% | 33.1% |
| Clatsop | 3.6% | 2.8% | 8.5% | 7.1% | 36.9% | 33.9% | 40.6% | 28.3% |
| Columbia | 3.1%† | 2.8%† | 7.3% | 6.6% | 34.0% | 31.0% | 41.6% | 30.4% |
| Coos | 5.4%† | 5.7%† | 12.6% | 11.0% | 34.1% | 28.5% | 45.5% | 41.8% |
| Crook | – | – | 9.9% | 9.0% | 48.0% | 46.2%* | 44.1% | 41.8% |
| Curry | 2.9%† | 1.5%† | 8.2% | 6.1%† | 40.3% | 25.8% | 46.9% | 32.6% |
| Deschutes | 1.4% | 1.2%* | 5.6% | 5.0% | 23.6% | 20.6%* | 36.0% | 32.1% |
| Douglas | 4.5% | 3.8%* | 11.4% | 10.1%* | 35.0% | 31.4% | 40.2% | 35.1% |
| Grant | 3.5%† | 2.4%† | 9.3%† | 6.7% | 33.4% | 21.7% | 49.2% | 42.1%† |
| Harney | – | – | 9.5%† | 8.1%† | 28.8% | 22.2% | 48.1% | 37.3% |
| Hood River | – | – | 4.9%† | 4.4%† | 25.5% | 23.2% | 17.7%† | 13.4%† |
| Jackson | 2.6% | 2.2% | 7.2% | 6.4% | 29.2% | 26.8% | 41.5% | 35.5% |
| Jefferson | 2.3%† | 1.9%† | 7.3% | 6.5% | 20.5% | 16.9%* | 29.4% | 20.2%† |
| Josephine | 3.9% | 3.3% | 8.2% | 5.8% | 38.0% | 34.3% | 45.5% | 43.7% |
| Klamath | 2.5%† | 2.3%† | 7.9% | 7.3% | 32.0% | 29.4% | 38.9% | 34.3% |
| Lake | – | – | – | – | 39.6% | 30.2% | 58.6% | 45.1% |
| Lane | 2.4% | 2.2% | 6.5% | 6.3% | 28.7% | 27.3% | 39.2% | 33.6% |
| Lincoln | 3.4%† | 2.8%† | 11.3% | 8.9% | 39.8% | 36.2% | 54.7% | 41.0% |
| Linn | 4.0% | 3.7% | 9.0% | 7.9% | 29.7% | 27.7% | 34.4% | 29.6% |
| Malheur | 2.7%† | 2.6%† | 8.0% | 7.6% | 30.2% | 27.8% | 41.2% | 37.5% |
| Marion | 2.4% | 2.3% | 8.3% | 8.3% | 28.1% | 27.9% | 32.1% | 29.1% |
| Morrow | – | – | 7.1%† | 6.8%† | 19.9%† | 16.1% | 33.8%† | 23.1%† |
| Multnomah | 1.8% | 1.8% | 6.0% | 6.2% | 22.8% | 23.0% | 36.4% | 33.7% |
| Polk | 2.5% | 2.2% | 6.6% | 6.0% | 25.9% | 24.4% | 34.7% | 31.0% |
| Tillamook | 2.8%† | 2.0%† | 10.0% | 8.3% | 35.9% | 32.2% | 40.9% | 29.6% |
| Umatilla | 2.9% | 2.7% | 9.6% | 9.3% | 32.1% | 30.9% | 46.1% | 39.7% |
| Union | 4.4%† | 3.9%† | 6.5% | 6.5% | 25.5% | 22.6% | 41.6% | 36.1% |
| Wallowa | – | – | 4.0%† | 2.4%† | 30.3% | 18.8% | 44.4% | 32.4% |
| Washington | 1.8% | 1.9% | 5.6% | 5.9% | 21.8% | 22.9% | 32.5% | 30.2% |
| Wheeler | – | – | – | – | – | – | – | – |
| Yamhill | 2.2% | 2.2% | 7.5% | 7.5% | 27.2% | 26.8% | 36.1% | 34.9% |
| Gilliam/Sherman/Wasco | 2.5%† | 1.8%† | 6.7% | 5.8% | 31.5% | 27.3% | 33.1% | 25.7% |

* Statistically significant difference compared with Oregon.

† This number may be statistically unreliable and should be interpreted with caution.

– This number is suppressed because it is statistically unreliable.

Age-adjusted estimates are usually adjusted to the 2000 Standard Population using three age groups (18-34, 35-54, and 55+), however, the Grant, Jefferson and Morrow high blood cholesterol estimates were based on collapsed age groups (18-54 and 55+) due to small numbers.

Source: Oregon BRFSS County Combined Dataset 2006-2009