Oregon Adults 65+ Who Had a Pneumonia Shot Within the Past Year, Oregon,
2014-2017

Adults 18 years & older	Unadjusted		
	%	95% Confidence Interval ¹	
		L. limit %	U. limit %
STATEWIDE	77.2	75.9	78.4
BAKER	76.8	62.1	86.9
BENTON	83.2	74.9	89.2
CLACKAMAS	77.5	73.1	81.5
CLATSOP	63.3 ~	49.6	75.1
COLUMBIA	73.5	62.1	82.5
coos	68.0 ~	59.2	75.7
CROOK	77.3	64.2	86.7
CURRY	69.0	54.7	80.4
DESCHUTES	80.8	75.4	85.3
DOUGLAS	82.9	77.3	87.3
GRANT	86.2	65.6	95.4
HARNEY	*	*	*
HOOD RIVER	94.6 ~	83.6	98.3
JACKSON	76.7	71.6	81.2
JEFFERSON	74.2	57.3	86.0
JOSEPHINE	73.6	66.4	79.7
KLAMATH	65.1 ~	54.0	74.9
LAKE	51.1 ~	28.5	73.3
LANE	78.0	74.1	81.4
LINCOLN	69.5	60.1	77.5
LINN	73.7	66.8	79.5
MALHEUR	65.2	50.4	77.5
MARION	85.9 ~	82.2	88.8
MORROW	52.5	20.6	82.6
MULTNOMAH	79.6	76.3	82.5
POLK	78.4	69.3	85.3
TILLAMOOK	75.9	61.1	86.3
UMATILLA	61.2 ~	50.0	71.2
UNION	62.3	46.5	75.9
WALLOWA	76.1	57.0	88.4
WASHINGTON	80.8	76.9	84.2
WHEELER	*	*	*
YAMHILL	84.5	77.1	89.9
GILLIAM/WASCO/SHERMAN			

^{*} This number is suppressed because it is statistically unreliable.

Source: Oregon Behavioral Risk Factors Surveillance System 2014-2017 county combined

NOTE: Age adjustment is based on three age groups: 18-34; 35-54; and 55+, per U.S.

[~] Statistically Significant: significantly higher or lower than the statewide rate.

¹ Confidence intervals provide a measure of how much an estimate might vary due to chance. We present 95% confidence intervals, which means that if we conducted this survey over and over, 95% of the confidence intervals calculated would contain the true population value. Wider intervals suggest the estimate is more unreliable and should be interpreted with caution, however it is the best estimate available.