

TABLE 3-5. Contraceptive Use, Number of Previous Abortions, and Number of Living Children by Age of Patient, Oregon Occurrence, 2002

Contraceptive Used, Previous Abortions, and Number of Living Children	Total	Age Groups								N.S.
		< 15	15-19	20-24	25-29	30-34	35-39	40-44	45+	
Total	13,172	46	2,242	4,462	2,914	1,971	1,098	409	25	5
None Used	8,620	32	1,631	2,916	1,865	1,208	688	261	16	3
No Previous Abortion	4,865	31	1,304	1,728	892	511	272	115	10	2
One	2,232	1	255	767	546	359	221	78	4	1
Two	874	—	50	274	247	175	95	32	1	—
Three	360	—	13	95	97	85	48	21	1	—
Four or More	266	—	6	45	79	75	47	14	—	—
Pills Used	1,437	—	207	576	343	198	87	24	1	1
No Previous Abortion	751	—	157	319	147	78	38	11	1	—
One	412	—	38	179	94	62	32	6	—	1
Two	184	—	11	58	68	34	10	3	—	—
Three	52	—	—	16	16	12	6	2	—	—
Four or More	35	—	1	3	16	12	1	2	—	—
Condoms Used	2,439	14	371	814	524	410	232	68	5	1
No Previous Abortion	1,284	13	307	475	230	153	78	25	2	1
One	704	1	54	227	162	152	82	25	1	—
Two	263	—	8	73	83	53	36	8	2	—
Three	107	—	2	22	32	27	16	8	—	—
Four or More	75	—	—	17	16	23	18	1	—	—
Other Contraceptive	886	—	56	227	232	186	113	68	4	—
No Previous Abortion	431	—	44	125	96	80	52	32	2	—
One	259	—	11	68	78	53	28	21	—	—
Two	115	—	—	18	39	28	19	9	2	—
Three	50	—	1	9	13	16	9	2	—	—
Four or More	28	—	—	6	6	9	5	2	—	—
Contraceptive Use Unknown	12	—	4	4	2	1	1	—	—	—
No Previous Abortion	8	—	3	3	1	—	1	—	—	—
One	3	—	—	1	1	1	—	—	—	—
Two	1	—	1	—	—	—	—	—	—	—
Three	—	—	—	—	—	—	—	—	—	—
Four or More	—	—	—	—	—	—	—	—	—	—
Previous Abortions Unknown ...	4	—	—	—	—	—	3	1	—	—
Number of Living Children										
No Children	5,709	44	1,791	2,210	961	426	205	64	6	2
Total with Children	7,459	2	451	2,250	1,952	1,545	893	344	19	3
One	3,435	1	379	1,393	781	516	272	88	3	2
Two	2,626	1	66	687	758	609	352	140	13	—
Three	949	—	4	142	306	276	148	70	2	1
Four	309	—	1	22	85	99	74	28	—	—
Five or More	140	—	1	6	22	45	47	18	1	—

— Quantity is zero.

NOTE: Contraceptive totals include abortions where the number of previous abortions is unknown. Multiple contraceptive methods may be reported for a single patient.