

TABLE 3-5. Contraceptive Use, Number of Previous Abortions, and Number of Living Children by Age of Patient, Oregon Occurrence, 2007

Contraceptive Used, Previous Abortions, and Number of Living Children	Total	Age Groups								N.S.
		< 15	15-19	20-24	25-29	30-34	35-39	40-44	45+	
Total	11,883	49	2,006	3,720	2,889	1,630	1,072	357	46	114
None Used	7,751	35	1,404	2,443	1,839	1,033	658	227	31	81
No Previous Abortion	4,498	34	1,171	1,492	891	458	280	108	15	49
One	1,843	1	186	613	504	279	186	60	6	8
Two	797	-	38	222	247	154	93	29	5	9
Three	291	-	3	62	98	70	42	12	1	3
Four or More	264	-	2	39	83	64	53	17	4	2
Pills Used	1,391	2	242	500	358	145	100	32	3	9
No Previous Abortion	794	2	197	318	150	57	48	14	2	6
One	372	-	38	119	124	48	26	15	-	2
Two	136	-	5	43	46	21	16	3	1	1
Three	37	-	-	12	14	6	5	-	-	-
Four or More	45	-	-	5	23	12	5	-	-	-
Condoms Used	2,025	11	295	578	514	321	210	70	7	19
No Previous Abortion	1,059	11	248	351	205	115	89	28	4	8
One	548	-	37	151	166	107	64	17	1	5
Two	240	-	8	48	72	58	33	16	1	4
Three	93	-	-	17	41	18	11	4	1	1
Four or More	72	-	-	5	29	22	12	4	-	-
Other Contraceptive	868	1	84	240	228	155	115	33	5	7
No Previous Abortion	403	1	62	128	97	56	41	14	3	1
One	239	-	17	60	70	46	39	4	1	2
Two	125	-	4	27	31	33	17	10	1	2
Three	48	-	1	12	17	9	6	3	-	-
Four or More	42	-	-	7	10	11	11	2	-	1
Contraceptive Use Unknown	25	1	3	4	3	4	-	2	-	8
No Previous Abortion	9	1	2	2	1	-	-	1	-	2
One	4	-	-	1	-	2	-	1	-	-
Two	4	-	-	1	2	1	-	-	-	-
Three	-	-	-	-	-	-	-	-	-	-
Four or More	-	-	-	-	-	-	-	-	-	-
Previous Abortions Unknown ...	19	-	2	8	2	3	2	1	-	1
Number of Living Children										
No Children	5,714	47	1,688	2,153	1,047	451	215	65	7	41
Total with Children	6,135	2	314	1,559	1,833	1,175	856	292	39	65
One	2,746	1	272	994	762	351	250	75	12	29
Two	2,140	1	38	433	702	455	343	133	15	20
Three	851	-	4	104	283	225	161	56	8	10
Four	289	-	-	22	67	109	68	15	2	6
Five or More	109	-	-	6	19	35	34	13	2	-

- Quantity is zero.

NOTE: Contraceptive totals include abortions where the number of previous abortions is unknown. Multiple contraceptive methods may be reported for a single patient.