

Table 2-39: Maternal characteristics by planned place of birth, Oregon occurrence, 2015

| Selected maternal characteristics | Total births ¹ | Planned hospital birth | | | Planned out-of-hospital birth | | |
|--|---------------------------|--------------------------------|--------|-------|-------------------------------|-------|-----|
| | | Clinical estimate of gestation | | | | | |
| | | <37 | 37-40 | 41+ | <37 | 37-40 | 41+ |
| Total births | 46,102 | 3,454 | 35,433 | 5,054 | 33 | 1,459 | 540 |
| Mother's age | | | | | | | |
| <20 | 2,327 | 186 | 1,837 | 273 | 1 | 18 | 7 |
| 20-24 | 8,987 | 637 | 7,028 | 1,047 | 6 | 172 | 67 |
| 25-29 | 13,360 | 932 | 10,253 | 1,517 | 11 | 453 | 157 |
| 30-34 | 13,247 | 946 | 9,995 | 1,554 | 12 | 522 | 186 |
| 35-39 | 6,716 | 567 | 5,162 | 615 | 2 | 246 | 106 |
| 40+ | 1,463 | 186 | 1,157 | 48 | 1 | 48 | 17 |
| Single mention race² | | | | | | | |
| White | 31,576 | 2,303 | 23,770 | 3,647 | 27 | 1,276 | 469 |
| African American | 1,044 | 96 | 777 | 144 | – | 14 | 7 |
| American Indian | 462 | 42 | 366 | 42 | – | 7 | 5 |
| Asian/Hawaiian/Pacific Islander | 2,613 | 199 | 2,155 | 219 | 3 | 20 | 10 |
| Other/multiple races | 1,871 | 141 | 1,453 | 193 | 1 | 50 | 21 |
| Hispanic | 8,536 | 673 | 6,912 | 809 | 2 | 92 | 28 |
| Marital status | | | | | | | |
| Married | 29,520 | 2,062 | 22,502 | 3,287 | 22 | 1,151 | 430 |
| Unmarried | 16,578 | 1,392 | 12,929 | 1,767 | 11 | 307 | 110 |
| Mother's education | | | | | | | |
| 8th grade or less | 1,406 | 115 | 1,149 | 116 | – | 16 | 5 |
| Some high school | 4,913 | 456 | 3,924 | 454 | 1 | 51 | 12 |
| High school graduate/GED | 10,118 | 803 | 7,941 | 1,003 | 5 | 238 | 93 |
| Some college | 11,468 | 881 | 8,822 | 1,174 | 9 | 392 | 163 |
| Associate's degree | 3,840 | 275 | 2,994 | 400 | 4 | 123 | 35 |
| Bachelor's degree | 8,800 | 573 | 6,447 | 1,176 | 9 | 421 | 153 |
| Postbaccalaureate | 5,337 | 309 | 4,017 | 710 | 5 | 209 | 76 |
| Source of payment³ | | | | | | | |
| Medicaid/Oregon Health Plan | 20,915 | 1,660 | 16,441 | 2,072 | 6 | 508 | 171 |
| Private insurance | 23,877 | 1,716 | 18,354 | 2,890 | 23 | 621 | 227 |
| Self-pay | 710 | 34 | 193 | 26 | 4 | 304 | 134 |
| Other coverage | 531 | 37 | 408 | 58 | – | 21 | 6 |
| Birth order | | | | | | | |
| 1st | 18,224 | 1,332 | 13,092 | 3,009 | 20 | 510 | 233 |
| 2nd | 14,805 | 987 | 11,903 | 1,218 | 6 | 507 | 146 |
| 3rd | 7,409 | 568 | 6,034 | 480 | 3 | 213 | 84 |
| 4th + | 5,664 | 567 | 4,404 | 347 | 4 | 229 | 77 |
| Pre-pregnancy body mass index | | | | | | | |
| Underweight (< 18.5) | 1,421 | 117 | 1,104 | 124 | 2 | 53 | 15 |
| Normal (18.5 - 24.9) | 21,404 | 1,436 | 16,149 | 2,524 | 20 | 915 | 297 |
| Overweight (25.0 - 29.9) | 11,367 | 854 | 8,773 | 1,290 | 8 | 277 | 141 |
| Obese (> 30.0) | 11,374 | 983 | 9,035 | 1,064 | 3 | 189 | 80 |
| Maternal tobacco use | | | | | | | |
| Tobacco use | 4,579 | 509 | 3,581 | 414 | 2 | 36 | 11 |
| No tobacco use | 41,382 | 2,928 | 31,756 | 4,630 | 31 | 1,413 | 526 |
| Initiation of care | | | | | | | |
| 1st trimester | 36,181 | 2,740 | 28,244 | 3,817 | 21 | 960 | 337 |
| 2nd trimester | 7,576 | 503 | 5,558 | 952 | 11 | 373 | 158 |
| 3rd trimester | 1,733 | 90 | 1,281 | 248 | 1 | 81 | 27 |
| No care | 328 | 81 | 164 | 12 | – | 23 | 13 |
| Prenatal care⁴ | | | | | | | |
| Adequate | 42,903 | 3,031 | 33,246 | 4,709 | 31 | 1,327 | 480 |
| Inadequate | 2,607 | 356 | 1,756 | 295 | 2 | 108 | 46 |

– Quantity is zero.

¹ Total includes 111 births that occurred en route, were unplanned home deliveries, or other out-of-hospital births not otherwise classified.

Total also includes 26 births with unknown gestation.

² Non-Hispanic single mention race. The Hispanic category may include any mention of race.

³ Expected principal method of payment for delivery. Actual method of payment may differ.

⁴ Adequate care: Care that began in the first or second trimester and included at least five visits.

Inadequate care: No care, or care that began in the third trimester or fewer than five visits.