

TABLE 3-5. Contraceptive use, number of previous abortions, and number of living children by age of patient, Oregon occurrence, 2015

Contraceptive used, previous abortions, and number of living children	Total	Age groups								
		<15	15-19	20-24	25-29	30-34	35-39	40-44	45+	N.S.
Total	8,610	21	901	2,622	2,297	1,551	871	318	26	3
None used	5,413	19	600	1,660	1,443	957	512	199	20	3
No previous abortion	3,221	19	520	1,152	766	439	219	94	11	1
One	1,243	–	64	345	353	266	154	53	7	1
Two	525	–	6	101	191	133	63	29	1	1
Three	175	–	2	25	63	46	32	7	–	–
Four or more	190	–	1	22	54	62	35	15	1	–
Pills used	861	–	111	301	231	136	67	15	–	–
No previous abortion	504	–	91	187	125	59	34	8	–	–
One	231	–	13	82	71	44	17	4	–	–
Two	78	–	3	20	26	18	10	1	–	–
Three	22	–	–	7	6	5	2	2	–	–
Four or more	16	–	–	4	3	5	4	–	–	–
Condoms used	1,108	2	102	307	271	221	154	47	4	–
No previous abortion	609	2	85	191	143	102	67	16	3	–
One	300	–	14	84	73	71	41	17	–	–
Two	108	–	1	21	30	30	19	6	1	–
Three	39	–	–	4	10	7	14	4	–	–
Four or more	41	–	–	4	13	10	10	4	–	–
Other contraceptive	902	–	63	255	263	178	103	39	1	–
No previous abortion	509	–	54	163	150	76	50	15	1	–
One	230	–	6	64	70	47	29	14	–	–
Two	104	–	3	22	25	32	15	7	–	–
Three	29	–	–	1	12	12	3	1	–	–
Four or more	21	–	–	2	4	8	5	2	–	–
Contraceptive use unknown ..	417	–	33	116	110	92	47	18	1	–
No previous abortion	251	–	30	80	60	44	26	11	–	–
One	97	–	3	25	31	24	11	3	–	–
Two	38	–	–	6	9	15	5	2	1	–
Three	13	–	–	2	4	4	2	1	–	–
Four or more	12	–	–	2	3	3	3	1	–	–
Number of living children										
No children ¹	4,229	21	754	1,618	1,037	518	216	59	5	1
Total with children	4,345	–	142	994	1,250	1,027	651	259	20	2
One	1,951	–	122	657	533	361	190	82	5	1
Two	1,438	–	15	264	462	367	243	76	10	1
Three	616	–	4	64	177	192	115	60	4	–
Four	222	–	1	7	66	63	58	27	–	–
Five or more	118	–	–	2	12	44	45	14	1	–

– Quantity is zero.
N.S. = Not stated.

¹ Rows will not add to total due to some patients having an unknown number of children.

NOTE: Contraceptive totals include abortions where the NUMBER of previous abortions is unknown. Multiple contraceptive methods may be reported for a single patient.