

TABLE 7-17. Neonatal, Postneonatal, and Infant Death Rates by Mother's Risk Factors, Oregon Residents, Birth Cohort 2000

Risk Factor	Neonatal ¹		Post-Neonatal ²		Infant ³	
	No.	Rate	No.	Rate	No.	Rate
Total	163	3.6	97	2.1	260	5.7
Marital Status						
Married	102	3.2	43	1.3	145	4.5
Unmarried	61	4.4	54	3.9	115	8.3
Mother's Age						
10-14	1	–	1	–	2	–
15-19	23	4.5	20	3.9	43	8.4
20-24	45	3.7	31	2.5	76	6.2
25-29	41	3.2	27	2.1	68	5.4
30-34	33	3.3	7	0.7	40	4.0
35-39	16	3.4	7	1.5	23	4.9
40-44	3	–	2	–	5	5.0
45+	–	–	1	–	1	–
Non-Hispanic						
White	114	3.3	76	2.2	190	5.5
African American	7	7.0	3	–	10	10.1
American Indian	2	–	3	–	5	7.4
Asian ⁴	4	–	4	–	8	3.5
Total Hispanic	36	4.9	11	1.5	47	6.4
Mother's Education						
8 th Grade or Less	10	3.6	8	2.9	18	6.5
Some High School	32	5.1	26	4.2	58	9.3
HS diploma/GED	56	3.8	33	2.2	89	6.0
More than High School	51	2.4	26	1.2	77	3.7
Start of Prenatal Care						
1 st Trimester	120	3.2	68	1.8	188	5.1
2 nd Trimester	27	4.0	26	3.8	53	7.8
3 rd Trimester	4	–	2	–	6	4.4
No Care	12	23.5	1	–	13	25.4
Tobacco Use						
Yes	21	3.4	35	5.7	56	9.2
No	126	3.2	61	1.6	187	4.8
Alcohol Use						
Yes	2	–	3	–	5	7.7
No	145	3.3	90	2.0	235	5.3
Multiple Birth						
Yes	22	16.7	6	4.5	28	21.2
No	132	3.0	91	2.0	223	5.0

– Quantity is zero or rate is based on fewer than five occurrences.

NOTE: Because of unreported items, the sum of all categories may not equal the total.

All rates per 1,000 live births.

¹ Neonatal deaths include infant deaths of less than 28 days.

² Postneonatal deaths occur from day 28 through 364 after birth.

³ Infant death is the death of a child prior to its first birthday.

⁴ Includes Chinese, Japanese, Filipino, and Other Asian & Pacific Islander.