

**TABLE 7-10. Postneonatal Deaths by Weeks of Gestation and Weight,
Oregon Residents, Birth Cohort 2003**

Birthweight (In Grams)	Total	Weeks of Gestation									
		<20	20-23	24-27	28-31	32-35	36	37-39	40	41+	N.S.
Total	80	—	—	3	5	8	4	28	18	14	—
001-349	—	—	—	—	—	—	—	—	—	—	—
350-499	—	—	—	—	—	—	—	—	—	—	—
<500	—	—	—	—	—	—	—	—	—	—	—
500-749	1	—	—	1	—	—	—	—	—	—	—
750-999	2	—	—	2	—	—	—	—	—	—	—
1000-1249	1	—	—	—	1	—	—	—	—	—	—
1250-1499	3	—	—	—	2	1	—	—	—	—	—
1500-1999	7	—	—	—	2	4	—	1	—	—	—
2000-2499	4	—	—	—	—	1	1	2	—	—	—
<2500	18	—	—	3	5	6	1	3	—	—	—
2500-2999	19	—	—	—	—	2	2	10	4	1	—
3000-3499	23	—	—	—	—	—	1	10	6	6	—
3500-3999	14	—	—	—	—	—	—	4	4	6	—
4000-4499	5	—	—	—	—	—	—	1	4	—	—
4500-4999	—	—	—	—	—	—	—	—	—	—	—
Unknown	1	—	—	—	—	—	—	—	—	1	—

— Quantity is zero.

Postneonatal deaths occur from day 28 through 364 after birth.

NOTE: Calculated gestation from reported date of last menses. If calculated gestation is unknown, the clinical estimate of gestation is used.