

TABLE 6-13. Years of Potential Life Lost before Age 75 from the Leading Causes of Death, by Year, Oregon Residents, 1998-2012

| Year | Total | Cancer | Unintended Injury | Heart Disease | Suicide | Alcohol-induced ¹ | Perinatal Conditions | Diabetes | CLRD ² |
|------------|---------|--------|-------------------|---------------|---------|------------------------------|----------------------|----------|-------------------|
| 1998 | 215,510 | 51,412 | 36,953 | 29,253 | 16,486 | 6,976 | 5,918 | 5,362 | 5,417 |
| 1999 | 207,575 | 48,693 | 29,370 | 30,546 | 13,915 | 5,645 | 8,396 | 5,546 | 5,878 |
| 2000 | 206,973 | 49,688 | 31,398 | 27,487 | 14,317 | 6,778 | 7,856 | 4,839 | 5,487 |
| 2001 | 211,233 | 51,244 | 30,249 | 27,225 | 15,023 | 7,821 | 8,396 | 5,852 | 5,567 |
| 2002 | 222,274 | 52,637 | 31,185 | 28,489 | 14,455 | 8,125 | 8,966 | 5,929 | 5,802 |
| 2003 | 225,545 | 50,810 | 34,383 | 28,869 | 15,585 | 10,033 | 8,591 | 7,237 | 6,493 |
| 2004 | 221,453 | 50,892 | 34,830 | 26,449 | 15,294 | 9,877 | 8,396 | 7,497 | 5,848 |
| 2005 | 224,868 | 53,166 | 31,845 | 26,721 | 14,874 | 9,553 | 10,131 | 7,585 | 6,543 |
| 2006 | 231,592 | 52,025 | 36,529 | 26,871 | 16,158 | 9,082 | 9,067 | 7,590 | 6,807 |
| 2007 | 234,443 | 51,747 | 36,820 | 27,845 | 16,266 | 10,168 | 10,311 | 7,551 | 7,307 |
| 2008 | 231,750 | 51,479 | 38,621 | 27,793 | 16,342 | 10,362 | 8,994 | 6,621 | 7,598 |
| 2009 | 230,153 | 53,568 | 34,029 | 25,605 | 17,158 | 10,686 | 8,323 | 7,530 | 7,341 |
| 2010 | 224,366 | 54,941 | 30,199 | 23,929 | 17,963 | 10,666 | 7,891 | 7,292 | 7,799 |
| 2011 | 230,525 | 55,353 | 33,117 | 24,368 | 18,023 | 11,984 | 8,201 | 7,831 | 7,604 |
| 2012 | 228,909 | 54,352 | 31,236 | 24,889 | 19,481 | 11,856 | 8,473 | 7,273 | 7,141 |

| Year | Congenital Anomalies | Cerebro-vascular Disease | Homicide ³ | Viral Hepatitis | Undetermined External Causes | Sudden Infant Death Syndrome | Pneumonia & Influenza | HIV Disease | Septicemia |
|------------|----------------------|--------------------------|-----------------------|-----------------|------------------------------|------------------------------|-----------------------|-------------|------------|
| 1998 | 7,491 | 6,015 | 5,511 | 1,561 | 1,854 | 3,353 | 2,372 | 2,372 | 1,153 |
| 1999 | 7,846 | 5,629 | 4,804 | 989 | 2,146 | 1,939 | 1,519 | 2,420 | 1,656 |
| 2000 | 6,556 | 5,276 | 3,798 | 1,713 | 2,040 | 3,802 | 1,301 | 2,040 | 1,446 |
| 2001 | 6,844 | 6,011 | 3,887 | 1,681 | 2,663 | 2,162 | 1,873 | 2,050 | 1,240 |
| 2002 | 7,439 | 6,012 | 4,728 | 2,560 | 3,592 | 2,310 | 2,344 | 2,691 | 1,423 |
| 2003 | 6,313 | 6,108 | 3,522 | 2,050 | 3,575 | 1,714 | 1,985 | 2,675 | 1,309 |
| 2004 | 6,720 | 6,221 | 4,502 | 2,105 | 3,284 | 1,416 | 1,671 | 1,902 | 1,481 |
| 2005 | 5,695 | 6,274 | 4,078 | 1,717 | 3,370 | 1,491 | 2,421 | 1,729 | 1,658 |
| 2006 | 6,918 | 5,737 | 4,429 | 1,817 | 3,390 | 2,236 | 1,578 | 1,478 | 1,429 |
| 2007 | 6,293 | 6,339 | 3,147 | 3,536 | 3,691 | 2,833 | 1,684 | 1,518 | 1,709 |
| 2008 | 6,271 | 5,135 | 3,949 | 2,860 | 2,693 | 1,492 | 2,236 | 1,045 | 1,839 |
| 2009 | 4,264 | 5,714 | 3,684 | 3,276 | 3,004 | 2,163 | 3,822 | 1,076 | 2,096 |
| 2010 | 5,688 | 5,206 | 4,080 | 3,197 | 3,432 | 2,385 | 1,760 | 1,130 | 1,660 |
| 2011 | 5,831 | 5,709 | 4,235 | 3,177 | 2,437 | 2,087 | 1,786 | 859 | 1,581 |
| 2012 | 5,405 | 5,171 | 4,159 | 2,597 | 2,379 | 1,865 | 1,482 | 1,359 | 1,253 |

¹ See Table 6-6, footnotes 38-39, for a list of included conditions and their ICD codes. Prior to 1999, figures do not include deaths due to alcohol poisoning.

² Chronic Lower Respiratory Disease.

³ Excludes legal intervention.