

**TABLE 7-10. Postneonatal deaths¹ by weeks of gestation and weight
Oregon residents, birth cohort 2012**

Birthweight (in grams)	Total	Weeks of gestation									
		<20	20-23	24-27	28-31	32-35	36	37-39	40	41+	N.S.
Total ²	79	—	—	2	2	8	5	37	17	8	—
001-349	—	—	—	—	—	—	—	—	—	—	—
350-499	—	—	—	—	—	—	—	—	—	—	—
<500	—	—	—	—	—	—	—	—	—	—	—
500-749	1	—	—	—	1	—	—	—	—	—	—
750-999	1	—	—	1	—	—	—	—	—	—	—
1000-1249	1	—	—	1	—	—	—	—	—	—	—
1250-1499	2	—	—	—	1	1	—	—	—	—	—
1500-1999	2	—	—	—	—	2	—	—	—	—	—
2000-2499	13	—	—	—	—	4	2	7	—	—	—
<2500	20	—	—	2	2	7	2	7	—	—	—
2500+	59	—	—	—	—	1	3	30	17	8	—
2500-2999	18	—	—	—	—	1	2	13	—	2	—
3000-3499	23	—	—	—	—	—	1	10	9	3	—
3500-3999	13	—	—	—	—	—	—	5	5	3	—
4000-4499	5	—	—	—	—	—	—	2	3	—	—
4500+	—	—	—	—	—	—	—	—	—	—	—

¹ Postneonatal deaths occur from day 28 through 364 after birth.

² Includes unknown weight.

— Quantity is zero.

NOTE: Calculated gestation from reported date of last menses. If calculated gestation is unknown, the clinical estimate of gestation is used. Total includes reports with unknown birthweight and/or gestation.