


OREGON NOROVIRUS WATCH, APRIL 2015. This issue of *Oregon Norovirus Watch* includes norovirus outbreak data from the 2014–2015 norovirus season. In April 2015, 13 norovirus or norovirus-like outbreaks were reported, of which 8/13 (62%) were in Nursing homes or Assisted-living facilities and 2/13 (15%) were restaurant-associated (see Fig. 1).




Since August 2014, 42/87 (48%) lab-confirmed and norovirus-like outbreaks occurred in Nursing homes or Assisted-living facilities. The most common norovirus genogroup subtype since August 2014 is GII.4 “Sydney” with 24/48 (50%) lab-confirmed outbreaks, followed by GII.4 “Miami,” which has been lab-confirmed in 6/48 (13%) outbreaks (see Fig. 2).

Figure 1. Laboratory-confirmed norovirus & norovirus-like outbreaks by setting, Oregon Aug 2014 – Apr 2015 (n=87)



*"Other" includes hotels, workplaces, private homes, caterers.

Figure 2. Laboratory-confirmed norovirus genotypes by month, Oregon, Aug 2014 – Apr 2015 (n=48)*



*48/87 (55%) suspect norovirus outbreaks have been lab-confirmed thus far.

Please send any comments or data requests to Keenan Williamson, MPH (email keenan.j.williamson@state.or.us) or Dr. Emilio DeBess (phone 971-673-1111).

Oregon Norovirus Watch monthly reports are also found on our calicivirus web site: <http://public.health.oregon.gov/DiseasesConditions/DiseasesAZ/Pages/disease.aspx?did=110>.