

**State of Oregon
West Nile Virus Summary Report
2013**

May 7, 2014

Emilio DeBess, DVM, MPVM

Acute and Communicable Disease Prevention

800 NE Oregon St., Ste. 772

Portland, OR 97232

Phone: (971) 673-1111

Fax: (971) 673-1100

E-mail: Emilio.E.DeBess@state.or.us

Table of Contents

2013 Program Highlights	3
Introduction	4
WNV Surveillance and Related Activities	4
Human Surveillance	4-5
Veterinary Surveillance	4-6
Avian Surveillance	6
Sentinel Chicken Surveillance	8
Mosquito Surveillance	9-10
Vector Control Districts in Oregon	12
References and Acknowledgment	13

List of Tables

Table 1	Confirmed WNV infections in Oregon, 2004–2013	3
Table 2	Trend data for Oregon residents who contracted WNV in Oregon, 2004–2013	6
Table 3	Equine WNV tests and positive test results for Oregon counties, 2013	6
Table 4	Avian WNV tests results for Oregon counties, 2004–2013	7
Table 5	WNV positive mosquito pools, Oregon 2013	8
Table 6	Female mosquitoes collected for testing by Oregon VCDs, 2013	9
Table 7	Trend data, WNV Positive Mosquito Pools, Oregon 2004–2013	10

List of Figures

Figure 1	Positive WNV tests for Oregon in 2013.	3
Figure 2	Map of Oregon with shaded counties reporting WNV in 2013.	5
Figure 3	Oregon vectors of WNV based on laboratory vector competence studies (2)	11
Figure 4	Oregon counties with participating vector control districts (VCDs)	12

2013 Program Highlights

Some of the principal findings and accomplishments of Oregon’s surveillance, education, and planning programs for West Nile virus (WNV) in 2013 include the following:

- Continued statewide surveillance of mosquitoes, humans, birds, sentinel chickens, and horses.
- 16 human cases of WNV reported.
- 6 equine cases
- 2 corvids (birds) positive for WNV.

Figure 1. Number of positive WNV tests, Oregon, 2013.




Table 1. Confirmed WNV infections in Oregon, 2004–2013.

Group	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
Human	5	8	73	27	16	12	0	0	12	16
Horses	32	46	35	16	0	5	0	2	2	6
Birds	23	15	25	52	2	16	0	0	2	2
Mosquito Pools	0	11	22	28	16	262	4	3	71	88
Sentinel Chickens	0	15	0	11	0	0	0	0	0	0

Introduction

Oregon's surveillance program for West Nile Virus (WNV) was launched in 2001. It first appeared in Oregon in 2004 when the first human, avian, and equine WNV cases were diagnosed.

In 2013, 16 Oregonians, 2 birds, 6 horses and 88 mosquito pools were diagnosed with WNV.


Twelve Vector Control Districts (VCDs) and one county health department perform mosquito surveillance in Oregon (Figure 4).

The VCDs collect mosquitoes and dead birds, identify them, and prepare them for testing. Some VCDs conduct initial WNV tests for mosquito pools and dead birds using RAMP (Rapid Analyte Measurement Platform). In counties without VCDs, this work may be conducted by the local health department or the Oregon Department of Fish and Wildlife (ODFW). Confirmatory testing of WNV for humans is performed by the Oregon State Public Health Laboratory (OSPHL). Oregon State University's (OSU's) Veterinary Diagnostic Laboratory performs WNV testing of mosquitoes, dead birds, horses, and other mammals.

The Oregon WNV surveillance findings for humans, horses, birds, and mosquitoes in 2013 are summarized in the sections that follow.

Figure 2. Map of Oregon with shaded counties reporting WNV, 2013.

WEST NILE VIRUS OREGON MAP 2013



County	Human	Chicken	Birds/Raptor	Horses	Mosquitoes
Harney	2				
Malheur	11		2	1	60
Baker					13
Benton	1*				
Jackson					4
Klamath				1	4
Lane	1*				
Linn	1*				
Morrow					2
Umatilla				3	2
Union				1	3
Total	16		2	6	88
	* Acquired out-of-county/state				

Updated: December 2013



WNV Surveillance and Related Activities

Human Surveillance

In 2013, 16 Oregon residents tested positive for WNV by IgM antibody

Table 2. Trend data for Oregon residents who contracted WNV in Oregon, 2004–2013.

Year	All Cases	Neuroinvasive	Deaths
2004	5	0	0
2005	8	1	0
2006	73	13	1
2007	27	7	1
2008	15	3	0
2009	8	0	0
2010	0	0	0
2011	0	0	0
2012	12	1	0
2013	17	8	0

Veterinary Surveillance

Surveillance for WNV in Oregon's equine population resulted in 6 positive tests while 14 other equine tests were negative for WNV. Positive test results for Oregon counties in 2013 are summarized in Table 3. No other mammals tested positive for WNV in 2013.

Table 3. Positive equine WNV test results, Oregon 2013.

County	Number of Positive Test Results
Klamath	1
Malheur	1
Umatilla	3
Union	1
Total	6

Avian Surveillance

Surveillance for WNV in Oregon’s avian population resulted in two positive test results out of 35 birds tested by OSU’s Veterinary Diagnostic Laboratory and the VCDs. Of the 35 birds that were collected, 20 were of the family Corvidae (a.k.a. corvids) while the remaining 15 were American species other than corvid. Table 4 shows the avian species collection totals in Oregon by county for 2013. Trend data for avian WNV testing and positive test results for Oregon counties for the years 2004–2013 are presented in Table 5.

Table 4. Avian WNV test results for Oregon counties, 2013.

Avian Species Collection Totals by County			
County	Total Corvid Tested	Total Other Species Tested	Total Positives
Coos	1	0	0
Jackson	2	0	0
Malheur	2	0	2
Marion	2	0	0
Morrow	0	3	0
Multnomah	0	9	0
Umatilla	0	1	0
Washington	1	1	0
TOTAL	8	14	2

Table 5. Avian WNV tests and trend of positive test results for Oregon counties, 2004–2013.

Year	Number Tested	Number Positive	% Positive
2004	448	23	5%
2005	298	15	5%
2006	212	25	12%
2007	246	55	22%
2008	117	2	2%
2009	90	16	18%
2010	24	0	0%
2011	20	0	0%
2012	35	2	6%
2013	22	2	9%

Sentinel Chicken Surveillance

None tested in 2013

Mosquito Surveillance

In 2013, the VCDs conducted surveillance for WNV in Oregon's mosquito population. Figure 4 (page 12) shows the counties with participating VCDs and their activities. Statewide, 107,994 mosquitoes or 2700 mosquito pools were collected and tested for WNV. The mosquitoes submitted represent 15 mosquito species. PCR testing for WNV was conducted by OSPHL and RAMP was performed by some VCDs. Table 6 displays the number of mosquito pools per species that tested positive for WNV in Oregon in 2013 only. Table 7 displays the mosquito species and the number of individual female mosquitoes that VCDs collected for testing in Oregon in 2013. Table 8 displays the mosquito species in Oregon between 2004 through 2013 found positive for WNV. Figure 3 indicates the efficiency of vector transmission for various mosquito species (information obtained from the Centers for Disease Control and Prevention).

Table 6. WNV Positive Mosquito Pools, Oregon 2013.

VCD	Mosquito Species	Number of Positive Mosquito Pools	Collection Date
Baker	<i>Culex tarsalis</i>	13	7/25-8/30
Jackson	<i>Culex tarsalis</i>	4	8/18-9/13
Klamath	<i>Culex tarsalis</i>	3	8/18-9/13
Klamath	<i>Anopheles freeborni</i>	1	8/30
Malheur	<i>Culex tarsalis</i>	56	6/12-9/10
Malheur	<i>Culex pipiens</i>	4	9/10
Morrow	<i>Culex pipiens</i>	1	9/11
Morrow	<i>Culex tarsalis</i>	1	9/18
Umatilla	<i>Culex pipiens</i>	1	9/5
Umatilla	<i>Culex tarsalis</i>	1	8/20
Union	<i>Culex pipiens</i>	1	8/30
Union	<i>Culex tarsalis</i>	2	8/19-9/12

Table 7. Female mosquitoes collected for testing by Oregon VCDs, 2013.

<i>County</i>	<i>Aedes dorsalis</i>	<i>Aedes increpitus</i>	<i>Aedes vexans</i>	<i>Anopheles freeborni</i>	<i>Anopheles punctipennis</i>	<i>Class Insecta</i>	<i>Coquillettidia perturbans</i>	<i>Culex erythrothorax</i>	<i>Culex pipiens</i>	<i>Culex tarsalis</i>	<i>Culiseta incidens</i>	<i>Culiseta inornata</i>	<i>Genus Ochlerotatus</i>	<i>Ochlerotatus melanimon</i>	<i>Other Species/Unknown</i>	
Baker										1976						
Columbia		32	7292		235		188			247			2316		11707	
Deschutes			670	650					830	3220		50				
Jackson			2130					41	4853	1784						
Klamath	4200		1461	1932		50			851	8107		4759		705		
Malheur									358	2600						
Morrow		549	1362	749	10				7371	6424		349				
Multnomah			1456		552		280		658	2648	164					528
Umatilla									4514	1639						
Union			162						2746	1574						
Washington				87	1067		1500		4014	4297						
Total	4200	581	14533	3418	1864	50	1968	41	26195	34516	164	5158	2316	705	12235	107994

Table 8. Trend data, WNV Positive Mosquito Pools, Oregon 2004–2013.

Year	Mosquito Species	Number of Positives
2004	-	-
2005	<i>Culex tarsalis</i> <i>Culex stigmatosoma</i> <i>Culex pipiens</i>	11 pools*
2006	<i>Culex tarsalis</i>	22 pools
2007	<i>Aedes vexans</i> <i>Culex pipiens</i> <i>Culex tarsalis</i>	8 pools 2 pools 23 pools
2008	<i>Aedes vexans</i> <i>Culex pipiens</i> <i>Culex tarsalis</i>	5 pools 3 pools 8 pools
2009	<i>Aedes vexans</i> <i>Anopheles freeborni</i> <i>Anopheles punctipennis</i> <i>Coquillettidia perturbans</i> <i>Culex pipiens</i> <i>Culex tarsalis</i> <i>Culex sp.</i>	1 pool 1 pool 1 pool 1 pool 75 pools 131 pools 52 pools
2010	<i>Culex pipiens</i> <i>Culex tarsalis</i> <i>Culex sp.</i>	1 pool 2 pools 1 pool
2011	<i>Culex sp.</i>	3 pools
2012	<i>Culex pipiens</i> <i>Culex tarsalis</i> <i>Culex sp.</i>	53 pools 3 pools 15 pools
2013	<i>Culex pipiens</i> <i>Culex tarsalis</i> <i>Anopheles freeborni</i>	14 pools 73 pools 1 pool

*1 pool ≈ 40 mosquitoes

**Figure 3. Potential Oregon vectors of WNV based on laboratory vector competence studies (1).
Posted with permission.**

Table 3. Potential for selected North American mosquitoes to transmit WNV based on bionomics, vector competence, virus isolations, and involvement with other arboviruses

Species	Association with other viruses ^a	Host preference	Activity time	Flight range	Vector competence for WNV ^b	Field isolations of WNV ^c	Potential to serve as a	
							Enzootic vector ^d	Bridge vector ^e
<i>Ae. aegypti</i>		Mammals	Crepuscular/day	200 m	+++ , 3	+	0	+
<i>Ae. albopictus</i>	EEE	Opportunistic	Crepuscular/day	200 m	++++ , 3, 6	+	+	++++
<i>Ae. vexans</i>	EEE, WEE, SLE	Mammals	Crepuscular/night	>25 km	++ 1, 5, 8	+++	0	++
<i>Cq. perturbans</i>	EEE	Opportunistic	Crepuscular/night	5 km	+, 4	+	+	+
<i>Cs. melanura</i>	EEE	Birds	Crepuscular/night	9 km	+, 8	++	++	0
<i>Cs. inornata</i>	WEE	Mammals	Crepuscular/night	2 km	+++ , 5	+	+	++
<i>Cx. stigmatosoma</i>	SLE	Birds	Night	1 km	+++ , 5	0	+++	+
<i>Cx. erythrothorax</i>	WEE	Opportunistic	Crepuscular/day	<2 km	++++ , 5	0	++	+++
<i>Cx. nigripalpus</i>	EEE, SLE	Opportunistic ^f	Crepuscular	5 km	++ , 4	+++	+++	++
<i>Cx. pipiens</i>	SLE	Birds	Crepuscular/night	2 km	+++ , 1, 3, 5	++++	+++++	++
<i>Cx. quinquefasciatus</i>	SLE	Birds	Crepuscular/night	2 km	+++ , 4, 5	0	++++	++
<i>Cx. restuans</i>	SLE	Birds	Crepuscular/night	2 km	++++ , 4	+++	+++++	++
<i>Cx. salinarius</i>	EEE, SLE	Opportunistic	Crepuscular/night	10 km	++++ , 4	+++	+++	++++
<i>Cx. tarsalis</i>	WEE, SLE	Opportunistic ^f	Crepuscular/night	>6 km	++++ , 5, 7	++++	++++	+++
<i>Oc. atropalpus</i>		Mammals	Day and night	1 km	++++ , 3	+	+	++
<i>Oc. canadensis</i>	EEE	Mammals	Day	2 km	++ , 8	+	0	++
<i>Oc. cantator</i>	EEE	Mammals	Day	>10 km	++ , 8	+	0	++
<i>Oc. dorsalis</i>	WEE	Mammals	Day and night	5 km	+++ , 5	+	0	++
<i>Oc. japonicus</i>	JE?	Mammals	Crepuscular/day	unk	++++ , 2, 3	+++	+	++++
<i>Oc. melanimon</i>	WEE	Mammals	Day and night	>10 km	+++ , 5	0	0	++
<i>Oc. sierrensis</i>		Mammals	Crepuscular/day	1 km	+, 5	0	0	+
<i>Oc. sollicitans</i>	EEE	Mammals	Crepuscular/night	>25 km	++ , 1, 3	+	0	+
<i>Oc. taeniorhynchus</i>	EEE	Mammals	Day and night	>25 km	+, 1, 3	+	0	+
<i>Oc. triseriatus</i>		Mammals	Day	200 m	+++ , 8	++	0	+++
<i>Ps. ferox</i>	SLE	Mammals	Day	2 km	0, 8	+	0	0

Distribution and bionomics based on and generalized from information in Carpenter and LaCasse (1955), Darsie and Ward (1981), and Moore et al. (1993).

^a Known association with other viruses with a similar transmission cycle. EEE, eastern equine encephalomyelitis virus; JE; Japanese encephalitis virus; SLE; St. Louis encephalitis virus; WEE; western equine encephalomyelitis virus. Based on Karabatsos (1985).

^b Efficiency with which this species is able to transmit WNV in the laboratory. 0, incompetent; +, inefficient; +++++, extremely efficient vector. Based on 1 (Turell et al. 2000), 2 (Sardelis and Turell 2001), 3 (Turell et al. 2001), 4 (Sardelis et al. 2001), 5 (Goddard et al. 2002), 6 (Sardelis et al. 2002), 7 (Turell et al. 2003), or 8 (present study).

^c Relative number of WNV-positive pools detected. 0, none; +, few; +++++, many.


^d Potential for this species to be an enzootic or maintenance vector based on virus isolations from the field, vector competence, feeding behavior, etc. 0, little to no risk; +++++, this species may play a major role.

^e Potential for this species to be an epizootic or bridge vector based on virus isolations from the field, vector competence, feeding behavior, etc. 0, little to no risk; +++++, this species may play a major role.

^f Feeds primarily on avian hosts in spring and early summer and mixed between avian and mammalian hosts in late summer and fall.

Vector Control Districts

Figure 4. Oregon counties with participating vector control districts (VCDs) and their activities.



References

1. Turell, MD, et al. "An Update on the Potential of North American Mosquitoes (*Diptera: Culicidae*) to Transmit West Nile Virus. *J. Med. Entomol.* 42(1): 57-62 (2005)

Acknowledgment:

To all the Oregon Vector Control Districts and the Oregon State University Veterinary Diagnostic Laboratory, without your input, admirable and hard work, none of this would be possible.