

Disability Oregon— PRAMS, 2019–2020



Background

Disability, as defined by the World Health Organization, encompasses impairments, activity limitations, and participation restrictions a person may experience in their daily lives.¹ There are many types of disability including: significant cognitive or emotional limitations; impairment of vision, hearing, or physical function; and inability to complete activities of daily living.²

Approximately 12-18% of reproductive-aged women have a disability.³ Compared with women without a disability, women with a disability are as likely to desire a future pregnancy, be sexually active, and experience pregnancy.⁴⁻⁶ However, women with a disability are less likely to report utilization of reproductive health care and more likely to experience pregnancy complications and adverse birth outcomes.⁵⁻⁸

Pregnancy Risk Assessment Monitoring System (PRAMS)

[PRAMS](#), which includes forty-six states, the District of Columbia, New York City, Northern Mariana Islands, and Puerto Rico, representing approximately 81% of all U.S. live births, collects site-specific, population-based data on maternal attitudes, behaviors, and experiences before, during, and shortly after pregnancy.

A supplemental questionnaire on disability was added by selected PRAMS sites and data collection began in 2019 to address the gap in population-based data on disability among women with a recent live birth. This supplemental questionnaire is made up of the Washington Group Short Set of Questions on Disability ([WG-Short Set](#)), which address six functional types of disability: remembering, seeing, hearing, communicating, walking/climbing stairs, and self-care.⁹

Prevalence* of Any Disability† and Disability Type‡ Among Women with a Recent Live Birth, 2019–2020

Type of Disability	Oregon % (95% CI)*		20 PRAMS Sites [¶] % (95% CI)*
	2019	2020	2020
Any disability [†]	7.7 (5.6-10.6)	7.5 (5.7-9.8)	6.7 (6.1-7.3)
Remembering [‡]	4.2 (2.6-6.7)	3.9 (2.6-5.6)	3.9 (3.5-4.4)
Seeing [‡]	2.5 (1.4-4.2)	1.5 (0.8-2.7)	1.8 (1.5-2.1)
Hearing [‡]	0.2 (0.1-0.5)	1.5 (0.7-3.0)	0.8 (0.6-1.0)
Communicating [‡]	0.8 (0.3-2.3)	0.8 (0.4-1.6)	0.4 (0.3-0.6)
Walking/Climbing stairs [‡]	1.1 (0.4-2.7)	0.6 (0.2-1.9)	0.6 (0.5-0.9)
Self-care [‡]	0.1 (0.0-0.3)	0.4 (0.2-1.1)	0.3 (0.2-0.5)

Abbreviation: CI, Confidence Interval

* Weighted prevalence presented.

† Any disability was defined as having “a lot of difficulty” or “cannot do at all” on one or more of the WG-Short Set.

‡ Type of disability was defined as having “a lot of difficulty” or “cannot do at all” on each of the individual WG-Short Set.

¶ 20 PRAMS sites that collected disability data and met or exceeded the 50% response rate threshold for 2020: Colorado, District of Columbia, Georgia, Kansas, Louisiana, Maryland, Massachusetts, Michigan, Mississippi, Missouri, Montana, Nebraska, New Hampshire, New Mexico, North Dakota, Oregon, South Dakota, Tennessee, Vermont, and Virginia.

Prevalence* of Any Disability† Among Women with a Recent Live Birth by Select Characteristics, 2020

Characteristic	Oregon % (95% CI)*	20 PRAMS Sites [†] % (95% CI)*
Age (years)		
≤19	6.9 (2.9-15.8) [^]	10.9 (7.6-15.3)
20-24	10.5 (5.9-18.0)	9.0 (7.5-10.7)
25-34	7.0 (4.8-10.0)	6.0 (5.3-6.8)
≥35	6.2 (3.2-11.9)	5.4 (4.2-6.8)
Race/Ethnicity		
Non-Hispanic		
American Indian or Alaska Native	7.9 (3.7-16.0)	10.8 (6.5-17.5)
Asian or Pacific Islander	4.8 (3.0-7.6)	4.1 (2.5-6.7)
Black	6.2 (3.5-10.7)	6.6 (5.4-8.0)
Another or Mixed Race	9.0 (6.2-13.0)	6.5 (4.2-9.8)
White	7.5 (5.0-11.1)	6.8 (6.1-7.7)
Hispanic	8.1 (5.9-10.9)	6.4 (5.1-7.9)
Education		
Less than high school	9.8 (4.8-18.9)	9.7 (7.8-12.0)
High school diploma or GED	12.3 (7.9-18.6)	9.2 (7.9-10.7)
Some college or higher	5.3 (3.5-7.9)	5.0 (4.4-5.7)

Abbreviations: CI, Confidence Interval; GED, General Education Development

*.†. See footnotes on first page

<30 respondents; not reported

[^] <60 respondents; may not be reliable

SUMMARY OF ALL PRAMS SITES[†]

Based on results from the 2020 data for 20 PRAMS sites, among women with a recent live birth:

- 6.7% reported experiencing any disability.
- 3.9% reported difficulty remembering, 1.8% reported difficulty seeing, 0.4% reported difficulty communicating, 0.6% reported difficulty walking/climbing stairs, 0.8% reported difficulty hearing, and 0.3% reported difficulty with self-care.

[†] See footnotes on first page

RESOURCES

Disability and Health: <https://www.cdc.gov/ncbddd/disabilityandhealth/women.html>

Center for Research on Women with Disabilities: <https://www.bcm.edu/research/research-centers/center-for-research-on-women-with-disabilities>

References

1. World Health Organization, International Classification of Functioning, Disability and Health (ICF). <https://www.who.int/classifications/international-classification-of-functioning-disability-and-health>. Geneva: 2001, WHO.
2. Centers for Disease Control and Prevention. Disability and health overview. <https://www.cdc.gov/ncbddd/disabilityandhealth/disability.html>.
3. Okoro CA, Hollis ND, Cyrus AC, Griffin-Blake S. Prevalence of disabilities and health care access by disability status and type among adults — United States, 2016. *MMWR Morb Mortal Wkly Rep.* 2018;67:882–887. doi:10.15585/mmwr.mm6732a3.
4. Bloom TL, Mosher W, Alhusen J, Lantos H, Hughes RB. Fertility desires and intentions among U.S. women by disability status: Findings from the 2011-2013 National Survey of Family Growth. *Matern Child Health J.* 2017;21(8):1606–1615. doi:10.1007/s10995-016-2250-3.
5. Mitra M, Clements KM, Zhang J, Smith LD. Disparities in adverse preconception risk factors between women with and without disabilities. *Matern Child Health J.* 2016;20(3):507-15. doi:10.1007/s10995-015-1848-1.
6. Mitra M, Clements KM, Zhang J, Iezzoni LI, Smeltzer SC, Long-Bellil LM. Maternal characteristics, pregnancy complications, and adverse birth outcomes among women with disabilities. *Med Care.* 2015;53(12):1027-32. doi:10.1097/MLR.0000000000000427.
7. Sonalkar, S., et al. (2020). "Gynecologic care for women with physical disabilities: a qualitative study of patients and providers." *Womens Health Issues.* 30(2):136-141. doi:10.1016/j.whi.2019.10.002
8. Holt L, C. M., Duncanson L, Hazen C, Kumar A, McKeon BA, Woodard L (2021). "Perceived barriers to gynecologic care by women who use wheelchairs." *Cureus.* 13(6). doi:10.1097/01.AOG.00000663120.90198.f2
9. Madans JH, Loeb ME, Altman BM. Measuring disability and monitoring the UN Convention on the Rights of Persons with Disabilities: the work of the Washington Group on disability statistics. *BMC Public Health.* 2011;11 Suppl 4:S4. doi:10.1186/1471-2458-11-S4-S4.

To learn more about PRAMS and for data availability by site and year visit: <https://www.cdc.gov/prams>



THE BEST SOURCE OF DATA ON MOTHERS AND BABIES