

Communicable Disease Control


Clostridioides difficile incidence

Clostridioides difficile (formerly *Clostridium difficile*) is a toxin-producing bacterium that can cause symptoms ranging from diarrhea to life-threatening inflammation of the colon. *C. difficile* infections (CDI) are often linked to medical care and individuals taking antibiotics. Nationally, CDI is the most common healthcare-associated infection. Population-based surveillance and mandatory reporting through the National Healthcare Safety Network (NHSN) allow us to identify populations at higher risk of acquiring CDI in the state of Oregon.

In 2017, we observed a decrease in Oregon’s healthcare-onset, laboratory-identified CDI standardized infection ratio (SIR). The 2017 SIR of 0.83 (95% CI:0.77–0.87) corresponds to 17% fewer observed infections than was predicted based on national baseline data. However, Oregon’s CDI SIR has yet to meet the 2020 U.S. Department of Health and Human Services (HHS) reduction target of an SIR of 0.70, which corresponds to a 30% reduction in CDI.

FIGURE 1

Healthcare-onset *C. difficile* infections, Oregon Acute Care Hospitals, 2012–2017



Source: National Healthcare Safety Network (NHSN)

Methods to control the spread of *C. difficile* in hospitals include improving hand hygiene compliance, reducing unnecessary antibiotic prescribing, training healthcare employees to minimize risk of spread from patient-to-patient when caring for patients with active infections, and training hospital environmental services employees how to remove *C. difficile* from hospital environments. To help fight the spread of *C. difficile*, Oregon's inter-facility transfer communication law took effect in 2014, which requires health care facilities to notify receiving facilities of patient *C. difficile* status upon transfer. In 2017, the Healthcare-Associated Infections Program initiated implementation of CDC's Targeted Assessment for Prevention (TAP) Strategy. This strategy provides healthcare facilities with information to support CDI prevention, including new CDI data metrics, surveys of healthcare worker awareness and perceptions of CDI prevention strategies, and options for quality improvement projects.

Additional Resources: [Oregon NHSN HAI Report](#)

About the Data: Data source is the National Healthcare Safety Network (NHSN). Data is derived from mandatory case reporting by healthcare facilities. Data includes all cases of hospital-onset, laboratory identified *C. difficile* infections. The standardized infection ratio (SIR) is the Centers for Disease Control and Prevention's (CDC) recommended measure of performance relative to national baselines. The metric is the ratio of reported infections in a calendar year to the predicted infections for that same year based on risk-adjusted national baseline rates. CDC recently updated HAI baselines using data entered in NHSN in 2015. This "rebaseline" updates both the source of aggregate data and the risk adjustment methodology used to create the baselines. The changes in SIR from 2015 to 2016 may be, in part, a reflection of these updates. As of 2016, SIRs do not include critical access hospitals. CDI data for critical access hospitals can be found at: <https://go.usa.gov/xUMGy>.

For More Information Contact: Rebecca Pierce, Rebecca.A.Pierce@state.or.us

Date Updated: September 13, 2018

[Oregon State Health Profile](#)

OHA 9153-D (Rev) 09/13: This document can be provided upon request in an alternate format for individuals with disabilities or in a language other than English for people with limited English skills. To request this publication in another format or language, contact the Publications and Design Section at 503-378-3486, 711 for TTY, or email dhs-oha.publicationrequest@state.or.us.