

Oregon 2002 -2005 BRFSS Data - Behavior Risk Survey Topics by County

Data are adjusted to the 2000 U.S. Census Population.	
County	Diabetes
OREGON	4.2%
Baker	7.7%
Benton	3.7%
Clackamas	4.0%
Clatsop	5.2%
Columbia	3.1%
Coos	4.5%
Crook	3.7%
Curry	6.8%
Deschutes	4.0%
Douglas	5.2%
Grant	2.4%
Harney	3.5%
Hood River	1.4%**
Jackson	3.9%
Jefferson	5.9%
Josephine	4.9%
Klamath	4.7%
Lake	2.5%
Lane	3.9%
Lincoln	5.6%
Linn	5.4%
Malheur	6.3%
Marion	4.2%
Morrow	7.0%
Multnomah	4.1%
Polk	4.4%
Tillamook	2.9%
Umatilla	3.6%
Union	4.5%
Wallowa	4.3%
Washington	4.5%
Yamhill	3.5%
Gilliam/Wheeler	0%†
Sherman/Wasco	2.1%

Age-adjusted Prevalence of Selected Chronic Diseases and Related Risk Factors Among Adults by County, Oregon 2002-2005

Tables were created using combined BRFSS survey results from 2002 through 2005. Two numbers are provided for each condition or disease: an unadjusted rate and an age-adjusted rate.

Unadjusted rates, provided in separate tables, give a description of the absolute burden of a disease or risk factor for an individual county.

Age-adjusted rates supplied in this table allow you to compare rates for a county to the State's rate. They adjust for differences that could result from a county's population being, on average, older or younger than the State's. When comparing counties with Oregon or with each other, use these age-adjusted rates.

† Rate based on less than 50 respondents overall or fewer than 12 persons in one age group; may not accurately reflect behavior of entire county and results should be interpreted with caution. Estimates based on 10 or fewer respondents are not reported.

** Statistically significant difference compared to Oregon.

Three age groups were used for age adjusting: 18-34yrs., 35-54yrs, and 55+yrs.

Source: Oregon Behavioral Risk Factor Surveillance System - healthoregon.org/brfss