

TABLE 6-11. Years of Potential Life Lost before Age 65 from the Leading Causes of Death, by Year, Oregon Residents, 1985-1999

Year	Total	Unintentional Injury	Cancer	Heart Disease ¹	Suicide	Perinatal Conditions	Congenital Anomalies	Homicide ²	Alcohol-induced Deaths ³
1985	118,035	30,221	18,735	12,480	8,792	8,763	8,019	3,393	2,334
1986	119,185	29,922	17,928	12,309	9,447	7,710	6,897	5,369	2,559
1987	120,949	30,110	18,723	12,736	8,203	10,387	7,257	4,827	2,412
1988	119,533	29,720	18,838	12,256	9,745	8,309	6,402	4,038	2,433
1989	116,878	26,633	17,924	11,489	9,714	10,989	6,149	4,047	2,985
1990	117,310	26,397	19,097	10,260	9,609	7,586	6,602	3,505	2,647
1991	113,112	23,842	19,215	11,005	9,801	6,291	6,710	4,152	2,582
1992	114,350	21,758	18,655	10,670	10,492	7,069	6,220	4,973	2,845
1993	123,280	25,797	19,747	12,169	9,772	5,391	7,125	4,475	3,334
1994	126,313	25,604	21,242	11,189	11,467	6,809	5,848	5,568	3,491
1995	128,177	28,912	20,505	12,226	12,029	4,932	5,394	5,139	3,856
1996	126,458	28,627	21,610	12,764	11,304	6,155	5,238	4,884	4,086
1997	120,508	27,322	21,233	12,748	10,937	6,596	5,867	4,081	3,783
1998	122,992	27,500	22,356	12,404	11,771	5,128	6,310	4,224	4,011
1999	117,350	21,710	21,254	13,390	9,807	7,276	6,523	3,724	3,142

Year	Diabetes	Cerebrovascular Disease	Chronic Lower Respiratory Disease	Acquired Immune Deficiency Syndrome	Sudden Infant Death Syndrome	Undetermined External Cause	Pneumonia and Influenza	Aortic Aneurysm	Epilepsy
1985	1,287	1,960	1,342	271	6,907	1,672	768	283	390
1986	1,038	1,540	1,438	825	7,805	1,925	955	406	214
1987	1,224	1,794	1,126	2,135	7,420	1,783	784	298	258
1988	1,631	1,783	1,197	3,076	6,387	1,176	1,220	240	334
1989	1,171	1,533	1,467	3,304	5,999	1,606	1,070	529	240
1990	1,181	1,770	1,341	4,778	7,098	1,427	1,494	404	280
1991	1,388	1,801	1,309	5,796	5,484	1,112	900	428	248
1992	1,916	2,087	1,213	6,479	5,423	1,706	1,224	575	261
1993	1,594	2,399	1,424	7,884	5,873	1,746	1,469	373	427
1994	1,890	2,799	1,309	8,419	4,064	1,747	1,434	515	406
1995	1,811	2,052	1,509	8,214	4,906	2,021	901	595	281
1996	2,019	2,277	1,625	5,559	3,033	2,265	1,115	368	258
1997	2,036	2,432	1,660	2,286	2,323	1,413	1,313	483	399
1998	2,447	2,520	1,392	1,668	2,903	1,342	1,177	435	320
1999	2,441	2,226	1,720	1,700	1,679	1,596	768	291	282

¹ Includes alcoholic cardiomyopathy.

² Excludes legal intervention.

³ Includes the alcohol-linked disorders represented by ICD-10 codes F10, G31.2, I42.6, K29.2, K70, O35.4, P04.3, R78.0, X45, X65, and Y15.