

**TABLE 6-3. Selected Leading Causes of Death with Rates,  
Oregon Residents, 1983-2002**

Year	Total	Major Cardiovascular Disease			Malignant Neoplasms	Chronic Lower Respiratory Disease	Pneumonia and Influenza	Diabetes Mellitus
		Diseases of the Heart <sup>1</sup>	Cerebrovascular Diseases	Arteriosclerosis				
Number of Deaths								
1983	22,361	7,910	2,021	470	4,943	971	634	329
1984	23,101	8,010	1,919	431	5,387	957	725	343
1985	23,824	8,192	2,000	432	5,410	1,097	838	317
1986	23,328	7,788	1,926	417	5,272	1,090	742	328
1987	24,181	7,936	1,958	440	5,594	1,233	743	395
1988	24,557	7,662	2,010	378	5,801	1,203	900	439
1989	24,679	7,482	2,006	355	5,819	1,272	924	450
1990	25,073	7,482	1,912	332	6,056	1,304	966	483
1991	24,935	7,139	2,004	307	6,268	1,353	791	540
1992	25,714	7,255	2,138	314	6,362	1,273	841	575
1993	27,596	7,652	2,202	341	6,622	1,595	1,014	642
1994	27,361	7,417	2,394	300	6,599	1,469	885	662
1995	28,190	7,529	2,483	298	6,824	1,460	899	705
1996	28,900	7,676	2,632	256	6,784	1,676	946	739
1997	28,750	7,500	2,582	237	6,790	1,648	909	816
1998	29,346	7,276	2,636	228	7,007	1,638	1,010	870
1999*	29,356	7,252	2,817	198	6,904	1,762	684	855
2000*	29,541	7,104	2,567	230	6,989	1,696	637	847
2001*	30,128	7,086	2,604	195	7,091	1,743	576	1,033
2002*	31,082	7,245	2,639	210	7,232	1,842	661	1,034
Rates <sup>2</sup>								
1983	848.6	300.2	76.7	17.8	187.6	36.9	24.1	12.5
1984	868.5	301.1	72.1	16.2	202.5	36.0	27.3	12.9
1985	890.4	306.2	74.7	16.1	202.2	41.0	31.3	11.8
1986	877.2	292.8	72.4	15.7	198.2	41.0	27.9	12.3
1987	898.9	295.0	72.8	16.4	208.0	45.8	27.6	14.7
1988	895.9	279.5	73.3	13.8	211.6	43.9	32.8	16.0
1989	884.2	268.1	71.9	12.7	208.5	45.6	33.1	16.1
1990	880.7	262.8	67.2	11.7	212.7	45.8	33.9	17.0
1991	851.0	243.7	68.4	10.5	213.9	46.2	27.0	18.4
1992	863.2	243.8	71.8	10.5	213.6	42.7	28.2	19.3
1993	908.4	251.9	72.5	11.2	218.0	52.5	33.4	21.1
1994	887.8	240.7	77.7	9.7	214.1	47.7	28.7	21.5
1995	900.1	240.4	79.3	9.5	217.9	46.6	28.7	22.5
1996	908.5	241.3	82.7	8.0	213.3	52.7	29.7	23.2
1997	893.7	233.1	80.3	7.4	211.1	51.2	28.3	25.4
1998	898.1	222.7	80.7	7.0	214.4	50.1	30.9	26.6
1999*	889.4	219.7	85.3	6.0	209.1	53.4	20.7	25.9
2000*	859.6	206.7	74.7	6.7	203.4	49.3	18.5	24.6
2001*	867.8	204.1	75.0	5.6	204.3	50.2	16.6	29.8
2002*	886.9	206.7	75.3	6.0	206.4	52.6	18.9	29.5

<sup>1</sup> Excludes alcoholic cardiomyopathy prior to 1999. <sup>2</sup> All rates per 100,000 population.

\* Beginning in 1999, causes of death were classified using the rubrics and methodology of the tenth revision of the International Classification of Disease (which supplanted the ninth revision). Therefore, data for deaths classified prior to this date should not be compared to 1999 and more recent data without applying ICD-9/ICD-10 comparability ratios. See Appendix B.

**TABLE 6-3. Selected Leading Causes of Death with Rates,  
Oregon Residents, 1983-2002 (Continued)**

Year	Alcohol-induced Deaths <sup>3</sup>	Alzheimer's Disease	Parkinson's Disease	Acquired Immune Deficiency Syndrome	External Causes			
					Unintentional Injuries	Suicide	Homicide <sup>4</sup>	Firearms (Any Manner)
Number of Deaths								
1983	337	114	75	2	1,156	417	110	352
1984	343	154	99	6	1,185	423	127	360
1985	308	200	104	24	1,207	417	118	325
1986	325	245	102	70	1,184	450	181	383
1987	311	309	111	78	1,185	400	157	348
1988	330	344	131	114	1,190	461	143	375
1989	334	355	130	135	1,151	459	142	391
1990	334	386	147	190	1,115	456	106	382
1991	306	462	144	224	1,013	460	126	363
1992	320	488	139	249	1,032	492	154	420
1993	363	550	169	305	1,185	472	142	392
1994	352	599	193	330	1,187	525	180	447
1995	358	688	232	333	1,293	526	154	439
1996	419	740	236	223	1,295	533	143	430
1997	382	718	214	93	1,281	538	125	428
1998	380	806	275	71	1,337	569	134	441
1999*	304	868	256	73	1,144	499	109	391
2000*	383	905	278	62	1,211	502	93	378
2001*	431	1,038	293	64	1,257	524	107	360
2002*	442	1,125	306	87	1,382	517	106	376
Rates <sup>2</sup>								
1983	12.8	4.3	2.8	0.1	43.9	15.8	4.2	13.4
1984	12.9	5.8	3.7	0.2	44.5	15.9	4.8	13.5
1985	11.5	7.5	3.9	0.9	45.1	15.6	4.4	12.1
1986	12.2	9.2	3.8	2.6	44.5	16.9	6.8	14.4
1987	11.6	11.5	4.1	2.9	44.1	14.9	5.8	12.9
1988	12.0	12.6	4.8	4.2	43.4	16.8	5.2	13.7
1989	12.0	12.7	4.7	4.8	41.2	16.4	5.1	14.0
1990	11.7	13.6	5.2	6.7	39.2	16.0	3.7	13.4
1991	10.4	15.8	4.9	7.6	34.6	15.7	4.3	12.4
1992	10.7	16.4	4.7	8.4	34.6	16.5	5.2	14.1
1993	11.9	18.1	5.6	10.0	39.0	15.5	4.7	12.9
1994	11.4	19.4	6.3	10.7	38.5	17.0	5.8	14.5
1995	11.4	22.0	7.4	10.6	41.3	16.8	4.9	14.0
1996	13.2	23.3	7.4	7.0	40.7	16.8	4.5	13.5
1997	11.9	22.3	6.6	2.9	39.8	16.7	3.9	13.3
1998	12.1	24.7	8.4	2.2	40.9	17.4	4.1	13.5
1999*	9.2	26.3	7.8	2.2	34.7	15.1	3.3	11.8
2000*	11.1	26.3	8.1	1.8	35.2	14.6	2.7	11.0
2001*	12.4	29.9	8.4	1.8	36.2	15.1	3.1	10.4
2002*	12.6	32.1	8.7	2.5	39.4	14.8	3.0	10.7

3 Includes the alcohol-linked disorders represented by ICD-10 codes F10, G31.2, G62.1, I42.6, K29.2, K70, O35.4, P04.3, R78.0, X45, X65 and Y15. 4 Included legal intervention prior to 1999. Data shown now exclude legal intervention.

\* Beginning in 1999, causes of death were classified using the rubrics and methodology of the tenth revision of the International Classification of Disease (which supplanted the ninth revision). Therefore, data for deaths classified prior to this date should not be compared to 1999 and more recent data without applying ICD-9/ICD-10 comparability ratios. See Appendix B.