

TABLE 6-50. Crude Death Rates for Selected Leading Causes of Mortality, United States, 1992-2006

Year	Total	Heart Disease	Cancer	Cerebrovascular Disease	CLRD ¹	Unintended Injury	Pneumonia and Influenza	Suicide	Diabetes
1992	852.9	277.2	206.0	59.2	37.5	34.9	20.7	12.0	20.0
1993	880.0	284.1	207.5	61.1	40.8	36.0	22.4	12.1	21.3
1994	875.4	277.1	207.1	61.9	40.6	36.0	21.8	12.0	22.2
1995	880.0	276.5	206.8	63.1	40.8	36.4	22.3	11.9	23.0
1996	872.5	272.4	205.3	63.3	41.6	36.7	22.3	11.6	23.7
1997	864.7	267.6	203.5	62.7	42.4	36.6	22.5	11.4	23.9
1998	864.2	263.7	202.1	58.9	43.1	37.1	23.7	11.3	24.4
1999	877.0	265.9	201.6	61.4	45.5	35.9	23.4	10.7	25.1
2000	873.6	257.9	200.5	60.3	44.9	34.0	24.3	10.3	24.9
2001	848.5	245.8	196.0	57.9	43.7	35.7	22.0	10.8	25.1
2002	847.3	241.7	193.2	56.4	43.3	37.0	22.8	11.0	25.4
2003	841.9	235.6	191.5	54.2	43.5	37.6	22.4	10.8	25.5
2004	816.5	222.2	188.6	51.1	41.5	38.1	20.3	11.0	24.9
2005	825.9	220.0	188.7	48.4	44.2	39.7	21.3	11.0	25.3
2006	810.4	211.0	187.0	45.8	41.6	40.6	18.8	11.1	24.2

Year	Arteriosclerosis ²	Alzheimer's Disease	Alcohol-induced ³	Homicide (excluding legal intervention)	Hypertension	Acquired Immune Deficiency Syndrome	Parkinson's Disease	Congenital Anomalies	Amyotrophic Lateral Sclerosis
1992	6.4	10.2	7.5	9.9	4.5	14.3	3.0	4.4	1.5
1993	6.5	11.4	7.5	9.9	4.9	15.7	3.5	4.3	1.4
1994	6.4	12.3	7.6	9.4	5.0	17.5	3.8	4.1	1.5
1995	6.2	13.3	7.6	8.6	5.2	17.7	4.1	4.1	1.5
1996	6.1	13.4	7.3	7.8	5.5	12.7	4.5	4.0	1.6
1997	5.8	13.8	7.2	7.3	5.7	6.7	4.6	3.9	1.6
1998	5.5	13.8	7.1	6.6	5.9	5.4	4.9	3.9	1.6
1999	5.5	16.3	7.1	6.2	6.2	5.4	5.4	3.8	1.9
2000	5.2	17.8	7.0	5.9	6.5	5.2	5.7	3.8	1.9
2001	4.9	18.9	7.0	7.1	6.8	5.0	5.8	3.7	1.9
2002	4.8	20.4	6.9	6.1	7.0	4.9	5.9	3.7	2.0
2003	4.5	21.8	7.0	6.1	7.5	4.7	6.2	3.6	2.0
2004	4.0	22.5	7.2	5.9	7.9	4.4	6.1	3.6	1.9
2005	4.0	24.2	7.3	6.1	8.4	4.2	6.6	3.5	1.9
2006	2.9	24.2	7.4	6.2	8.0	4.0	6.5	3.5	-

All rates per 100,000 population. A "-" indicates that the data are not available. Rates for 2006 are preliminary.

1. CLRD consists principally of bronchitis, emphysema, asthma, and chronic airways obstruction.
2. Beginning in 2006, the National Center for Health Statistics changed the ICD-10 codes to include only ICD-10 code I70.
3. Includes the alcohol-linked disorders represented by ICD-9 codes 291.0-291.9, 303, 305.0, 357.5, 425.5, 535.5 and 571.0-571.3, prior to 1999. For current ICD-10 components, see Table 6-6, footnotes 38-39.

NOTE: Beginning in 1999, causes of death were classified using the rubrics and methodology of the tenth revision of the International Classification of Disease (which supplanted the ninth revision). Final ICD-9/ICD-10 comparability ratios have been applied to all rates prior to 1999, except ALS and alcohol-induced deaths, where ratios were not calculated. See Appendix B.