

**TABLE 7-9. Late neonatal deaths¹ by weeks of gestation and weight
Oregon residents, birth cohort 2012**

Birthweight (in grams)	Total	Weeks of gestation									N.S.
		<20	20-23	24-27	28-31	32-35	36	37-39	40	41+	
Total ²	30	–	1	8	–	3	2	12	3	1	–
001-349	–	–	–	–	–	–	–	–	–	–	–
350-499	–	–	–	–	–	–	–	–	–	–	–
<500	–	–	–	–	–	–	–	–	–	–	–
500-749	5	–	1	4	–	–	–	–	–	–	–
750-999	4	–	–	4	–	–	–	–	–	–	–
1000-1249	–	–	–	–	–	–	–	–	–	–	–
1250-1499	–	–	–	–	–	–	–	–	–	–	–
1500-1999	2	–	–	–	–	1	–	1	–	–	–
2000-2499	2	–	–	–	–	2	–	–	–	–	–
<2500	13	–	1	8	–	3	–	1	–	–	–
2500+	17	–	–	–	–	–	2	11	3	1	–
2500-2999	4	–	–	–	–	–	1	3	–	–	–
3000-3499	8	–	–	–	–	–	1	5	2	–	–
3500-3999	3	–	–	–	–	–	–	2	–	1	–
4000-4499	1	–	–	–	–	–	–	–	1	–	–
4500+	1	–	–	–	–	–	–	1	–	–	–

¹ Late neonatal deaths occur from day 7 through 27 after birth.

² Includes unknown weight.

– Quantity is zero.

NOTE: Calculated gestation from reported date of last menses. If calculated gestation is unknown, the clinical estimate of gestation is used. Total includes reports with unknown birthweight and/or gestation.