

TABLE 6-20. Tobacco-linked deaths by cause of death, Oregon residents, 2014

Selected causes of death (and their ICD-10 codes)	Total	Linked ¹		Not linked		Unknown	
		Number	Percent	Number	Percent	Number	Percent
Total ²	34,160	7,270	21.3	19,019	55.7	7,871	23.0
Malignant neoplasms	3,633	1,876	51.6	1,083	29.8	674	18.6
Oral cavity, lip, pharynx (C00.0-C14.8)	138	72	52.2	40	29.0	26	18.8
Esophagus (C15)	239	80	33.5	79	33.1	80	33.5
Stomach (C16)	139	20	14.4	82	59.0	37	26.6
Pancreas (C25)	569	53	9.3	381	67.0	135	23.7
Larynx (C32)	37	23	62.2	1	2.7	13	35.1
Lung, bronchi, and trachea (C33-C34)	1,982	1,527	77.0	194	9.8	261	13.2
Cervix uteri (C53)	51	9	17.6	33	64.7	9	17.6
Kidney, other urinary tract (C64-C65)	182	29	15.9	110	60.4	43	23.6
Urinary bladder (C67)	178	59	33.1	67	37.6	52	29.2
Acute myeloid leukemia (C92.0)	118	4	3.4	96	81.4	18	15.3
Cardiovascular disease	8,373	1,828	21.8	4,196	50.1	2,349	28.1
Ischemic heart disease (I20-I25)	3,316	994	30.0	1,392	42.0	930	28.0
Other heart disease (I00-I09, I26-I51)	2,926	399	13.6	1,709	58.4	818	28.0
Cerebrovascular disease (I60-I69)	1,821	303	16.6	991	54.4	527	28.9
Atherosclerosis (I70)	41	16	39.0	17	41.5	8	19.5
Aortic aneurysm (I71)	143	65	45.5	45	31.5	33	23.1
Other arterial disease (I72-I78)	126	51	40.5	42	33.3	33	26.2
Respiratory diseases	2,312	1,553	67.2	406	17.6	353	15.3
Pneumonia and influenza (J09-J18)	449	62	13.8	260	57.9	127	28.3
Bronchitis and emphysema (J40-J43)	138	118	85.5	10	7.2	10	7.2
Other chronic airways obstruction (J44)	1,725	1,373	79.6	136	7.9	216	12.5
Perinatal conditions ³	91	—	—	83	91.2	8	8.8
Selected perinatal conditions ⁴	66	—	—	61	92.4	5	7.6
Sudden infant death syndrome (R95)	25	—	—	22	88.0	3	12.0
Other causes	19,751	2,013	10.2	13,251	67.1	4,487	22.7

¹ The Oregon death certificate asks 'Did tobacco use contribute to death?' This is followed by four checkboxes: 'Yes,' 'No,' 'Probably,' and 'Unknown.' The linked category includes deaths listed as 'Yes' or 'Probably' or if a contributing cause is F17.

² The causes of death shown in this table are those linked to tobacco use by the federal Centers for Disease Control and Prevention (CDC. Annual Smoking-Attributable Mortality, Years of Potential Life Lost, and Productivity Losses -- United States, 1997-2001. MMWR 2005; 54:625-628.).

³ The number of infant deaths resulting from exposure to tobacco combustion products is underreported.

⁴ The category includes the following conditions: other disorders related to short gestation and low birthweight (P07), respiratory distress of newborn (P22), congenital pneumonia (P23), neonatal aspiration syndromes (P24), and other respiratory conditions originating in the perinatal period (P25-P28).

— Quantity is zero.