

Comparative Life Cycle Assessment of Expanded Polystyrene Dispositions (Updated)

Materials Management

July 19, 2022

Material Lists Technical Workgroup Meeting #4

Agenda

- Goal and Scope
- Results
- Interpretation and Limitations
- Potential Next Steps



Goal and Scope

Project Goals/Objectives

- Using Comparative Life Cycle Assessment
 - Quantify the environmental impacts of different end of life management scenarios for Expanded Polystyrene (EPS) to identify trade-offs and key variables.



Scope – Functional (Declared) Unit

- **Function:** Disposition of EPS through different end of life pathways
- **Magnitude/unit:** 1 us ton (short ton)




Scope – Key Variables Evaluated

- Collection - Marginal vs Additional Drop-Off vs On Route
- Densification – Onsite vs Offsite vs Undensified
 - Transport – Densified vs Undensified
- End of Life Dispositions - Chemical Recycling (Oregon) vs Mechanical Recycling (Asia) vs Distant Landfilling vs Local Landfilling

Scenarios Evaluated

Scenario Number	Collection	Densification	Disposition
S1	Drop-Off (Marginal)	On-site	Pyrolysis (in-state)
S2	Drop-Off (Additional)	On-site	Pyrolysis (in-state)
S3	Drop-Off (Marginal)	On-site	Mechanical Recycling (Asia)
S4	Drop-Off (Additional)	On-site	Mechanical Recycling (Asia)
S5	Drop-Off (Marginal)	Off-Site	Pyrolysis (in-state)
S6	Drop-Off (Additional)	Off-Site	Pyrolysis (in-state)
S7	Drop-Off (Marginal)	Off-Site	Mechanical Recycling (Asia)
S8	Drop-Off (Additional)	Off-Site	Mechanical Recycling (Asia)
S9	Drop-Off (Marginal)	None/Undensified	Pyrolysis (in-state)
S10	Drop-Off (Additional)	None/Undensified	Pyrolysis (in-state)
S11	Drop-Off (Marginal)	None/Undensified	Distant Landfill
S12	Drop-Off (Additional)	None/Undensified	Distant Landfill
S13	On Route to Transfer Station	None/Undensified	Distant Landfill
S14	Drop-Off (Marginal)	None/Undensified	Nearby Landfill
S15	Drop-Off (Additional)	None/Undensified	Nearby Landfill
S16	On Route Direct to Landfill	None/Undensified	Nearby Landfill
S17	Drop-Off (Marginal)	On-site	Mechanical Recycling (California)
S18	Drop-Off (Additional)	On-site	Mechanical Recycling (California)
S19	Drop-Off (Marginal)	Off-Site	Mechanical Recycling (California)
S20	Drop-Off (Additional)	Off-Site	Mechanical Recycling (California)

Scope – System Boundary



Scope – System Boundary

- **Temporal Coverage** – 2016-2022
- **Geographical Coverage** – Oregon
- **Technological Coverage** – This study is intended to represent materials management options for expanded polystyrene the foreground system covers technology and processes related to transportation of EPS to central locations or collection depots, mechanical densification, transport to end markets, chemical recycling, mechanical recycling, or landfilling. The background system includes electricity, thermal energy, and energy carriers (e.g. fuels).

Scope – Data Sources

- Primary Data Sources
 - Mechanical Densification – from Tillamook County
 - Pyrolysis – from Agilyx/Regenyx Air Contaminant Discharge Permit - 2020 Annual Report
- Secondary Data Sources
 - Truck Emissions – diesel combustion from USLCI (US DOE)
 - Truck Fuel Efficiency – US EPA Smartway
 - Passenger Vehicle Emissions – GaBi Database
 - Ship Emissions – GaBi Database
 - Mechanical Recycling – GaBi Database
 - Landfilling – GaBi Database
 - Fuels (Diesel or Gasoline) – GaBi Database
 - Production Emissions for Displaced Materials (Styrene and Polystyrene) – GaBi Database

Scope – Selected Impact Categories and Indicators


TRACI 2.1 LCIA Categories

- Acidification Potential (AP)
- Eutrophication Potential (EP)
- Ecotoxicity (ETP)
- Global Warming Potential (GWP100)
- Particulate Matter (PM2.5) Potential
- Human Toxicity Potential (HTP) – Cancer
- Human Toxicity Potential (HTP) – NonCancer
- Ozone Depletion Potential (ODP)
- Smog Formation Potential (SFP)

Environmental Indicators


- Fossil Resource use
- Water Consumption
- Primary Energy Demand

Global Warming Potential




Source: thinkstep, used with permission

Acidification Potential



Source: thinkstep, used with permission

Eutrophication Potential



© thinkstep AG 2015, all rights reserved

Source: thinkstep, used with permission


Smog Formation Potential



Source: thinkstep, used with permission

Ozone Depletion Potential

© thinkstep AG 2015, all rights reserved



Source: thinkstep, used with permission

Primary Energy Demand




© thinkstep AG 2015, all rights reserved

+



Source: thinkstep, used with permission and iStockphoto.com/DrAfter123

Freshwater Consumption




Source: thinkstep, used with permission

Preliminary Results


Life Cycle Impact Assessment (LCIA) and Indicators

Scenario Number	Collection	Densification	Disposition
S1	Drop-Off (Marginal)	On-site	Pyrolysis (in-state)
S2	Drop-Off (Additional)	On-site	Pyrolysis (in-state)
S3	Drop-Off (Marginal)	On-site	Mechanical Recycling (Asia)
S4	Drop-Off (Additional)	On-site	Mechanical Recycling (Asia)
S5	Drop-Off (Marginal)	Off-Site	Pyrolysis (in-state)
S6	Drop-Off (Additional)	Off-Site	Pyrolysis (in-state)
S7	Drop-Off (Marginal)	Off-Site	Mechanical Recycling (Asia)
S8	Drop-Off (Additional)	Off-Site	Mechanical Recycling (Asia)
S9	Drop-Off (Marginal)	None/Undensified	Pyrolysis (in-state)
S10	Drop-Off (Additional)	None/Undensified	Pyrolysis (in-state)
S11	Drop-Off (Marginal)	None/Undensified	Distant Landfill
S12	Drop-Off (Additional)	None/Undensified	Distant Landfill
S13	On Route to Transfer Station	None/Undensified	Distant Landfill
S14	Drop-Off (Marginal)	None/Undensified	Nearby Landfill
S15	Drop-Off (Additional)	None/Undensified	Nearby Landfill
S16	On Route Direct to Landfill	None/Undensified	Nearby Landfill
S17	Drop-Off (Marginal)	On-site	Mechanical Recycling (California)
S18	Drop-Off (Additional)	On-site	Mechanical Recycling (California)
S19	Drop-Off (Marginal)	Off-Site	Mechanical Recycling (California)
S20	Drop-Off (Additional)	Off-Site	Mechanical Recycling (California)


LCIA Results – Global Warming Potential (GWP)




LCIA Results – Global Warming Potential (GWP)




LCIA Results – Global Warming Potential (GWP)




LCIA Results – Global Warming Potential (GWP)




LCIA Results – Global Warming Potential (GWP)




LCIA Results – Global Warming Potential (GWP)




LCIA Results – Global Warming Potential (GWP)




LCIA Results – Global Warming Potential (GWP)




LCIA Results – Acidification Potential (AP)




LCIA Results – Eutrophication Potential (EP)




LCIA Results – Ecotoxicity Potential (ETP)




LCIA Results – Particulate Matter (PM 2.5)




LCIA Results – Human Toxicity Potential (Cancer)




LCIA Results - Human Toxicity Potential (NonCancer)




LCIA Results – Ozone Depletion Potential (ODP)




LCIA Results – Smog Formation Potential (SFP)




Indicator Results – Fossil Resource Use



Indicator Results – Bluewater Consumption




Indicator Results – Primary Energy Demand (PED)



Damage Costs


Damage Costs by Life Cycle Stage

Social Cost Contribution Analysis - All Scenarios



Damage Costs by Impact Category

Social Cost Contribution Analysis - All Scenarios



Interpretation

Key Findings



- **Convenience of drop-off sites matters** - idea of marginal vs additional transport was the single biggest variable influencing results.
- **Densification is justified** when transport distances are large
- **Densification** can be on-site or off-site
- **Disposition results are mixed**
 - The best disposition varies by impact category
 - Also depends on whether your landfill is nearby or distant

Assumptions and Limitations

Assumptions

- Average distance traveled for drop-off (additional) is 4 miles (so 8 miles round trip) all of these emissions are allocated to EPS recovery and so do count towards the impacts of this system.
- Average distance traveled for drop-off (marginal) is 4 miles (so 8 miles round trip) however the emissions are allocated to the primary purpose for the trip (e.g. grocery store) and so do not count towards the impacts of this system.
- Densification of EPS is based on mechanical densifier technology only (however thermo-mechanical densifiers are also used in practice)
- Transport for drop-off is by passenger vehicle
- Transport to landfill is by truck
- Transport to chemical recycling is by truck
- Transport to mechanical recycling is a combination of transport by truck and ocean ship
- Have scaled-up the impacts of landfilling by a factor of approximately 5. This reflects an assumption that landfill operations for an undensified material (e.g. EPS) will increase because of the volume of this material compared to municipal solid waste generically (based on the ratio of the density of EPS compared to the average density of MSW).
- Model assumes 1:1 substitution for primary material production as a recycling credit. In other words, for each unit of EPS recovered (after losses are accounted for) an equivalent unit of primary production is avoided (e.g. Styrene or Polystyrene).

Assumptions and Limitations (cont.)

Limitations

- No information on co-products (char, syngas, wax, etc.) of pyrolysis are included, so all emissions associated with pyrolysis are allocated exclusively to styrene monomer produced. Depending on the amount and quality of co-products, when included, a reduction in the process emissions for pyrolysis is expected (though it is predicted to be small if allocation is based on the economic value of these co-products)
- No direct human health exposures are accounted for by processors of this material (e.g. those handling EPS at the recycling facility)
- The effects of mismanagement of these materials (e.g. litter) are not accounted for in the model or impact results.
- Domestic and international recycling processes are modeled using the same underlying data. As such, no regional variations in recycling technology, environmental laws, or energy systems are accounted for. It's possible that these differences, should they exist, could affect the recycling process emissions profile.

Feedback and/or Questions

Thank You!

Peter Canepa (peter.canepa@state.or.us)

