

# State Department of Geology and Mineral Industries

702 Woodlark Building  
Portland, Oregon

Roberts Manganese

Sixes River Area

Curry County

The grade of the "ore" is low, averages about 30% Mn. and is high in silica. Assays were stated by Mr. Roberts to have run 45%, but apparently on very carefully picked ore. Chert is very abundant in the manganese boulders.

The distance from the deposit to the end of the Floras Creek road is a distinct disadvantage, although several logging roads connect with that road from points near the deposit. Resurfacing and regrading of at least 3 miles of road would be necessary before the deposit could produce ore.

The tonnage of commercial ore is very low. Low grade manganese (30% Mn) is present in considerable quantities up to a probable 200 tons, in sight on the talus slope and exposed in the face.

Mr. Roberts has leased the property to Toy L. Young (Chinese), mining operator from Oakland, California, who is believed to have some connection with H.L. Wadell of Golden Surf Mining Interests. Mr. Roberts is to get \$2.00 per ton of ore mined.

Recommendations: It was recommended that before further work is done that the deposit be carefully sampled and ore carefully hand picked to determine what tonnage, if any, of commercial ore can be expected. It is not believed that the deposit can be worked profitably due to the distance from the end of the road, the distance to the highway, the low manganese content and the high silica content.

Informant: Lee J. Roberts

Randall E. Brown



# State Department of Geology and Mineral Industries

702 Woodlark Building  
Portland, Oregon

Report by: Randall E. Brown  
Date of visit: Feb. 12, 1942

## Roberts Manganese

## Sixes River Area

## Curry County

Manganese oxides are exposed 1.5 miles northeast of Edson Butte near the South Fork of Floras Creek. The manganese occurs in a series of cherts which crop out around a small hill, in place. The chert lens in which the manganese occurs is continuous for more than 100 feet and averages about 3 feet thick. The grade of ore is below 40% except locally, and is highly siliceous. The deposit is located near an old logging road which would require rebuilding for 3 miles to permit operation.

Owner: Lee J. Roberts                      Langlois, Oregon

Area: Deeded land

Location: Located two miles beyond the end of the Floras Creek road in the NW 1/4 of Sec. 13, T. 31 S., R. 14 W. a half mile southwest of the South Fork of Floras Creek.

Deposit is 12 miles from US Highway 101 at a point 1/2 mile south of Langlois, and 32 miles south of Coquille, the nearest railhead.

History: Property was recently leased to Toy L. Young, mining operator from Oakland, California, and at present is being opened up to expose the manganese zone with intent of stockpiling the ore.

Topography: Deposit is exposed on the south side of a small hill, about 50 feet above the base of the hill and 30 feet from the top. The talus slope below the exposure contains considerable manganese boulders. Slope is downhill to the end of the Floras Creek road. Deposit is in cutover land, in moderately rugged country, at an altitude of 1,000 feet.

Development: Deposit is exposed over a distance of 100 feet and from 3 to 10 feet vertically. The manganese zone is terminated on either end by the ends of the hill. No work has been done to determine the depth of the manganese zone.

Geology: The country rock, according to Diller, is basalt. If covering the area designated by Diller, the ridge containing the manganese is surrounded by the basalt. However, it is believed that the deposit lies on the northern edge of the basalt plug. The rock cropping out above and below the chert zone is Myrtle sandstone. The manganese occurs in a red chert lens which constitutes the basal member of a series of cherts resting on sandstone bedrock.



# State Department of Geology and Mineral Industries

702 Woodlark Building  
Portland, Oregon

White chert crops out to the crest of the hill, about 30 feet above the top of the red chert lens, with traces of sandstone at the hill summit.

The manganese occurs as pods, boulders and irregular masses of psilomelane and pyrolusite within the red chert. Only traces of stain occur in either the white chert or the underlying sandstone. The highest grade ore is found at the western end of the deposit where the manganese zone is less than 3 feet thick. Toward the east end the zone thickens to ten feet but the manganese decreases in grade. Apparently the effect of weathering has been to concentrate the manganese in a narrower zone on the west.

The chert composes the entire top of the hill and can be traced completely around the hill. The red chert crops out around the hill but is only stained on the north side. The attitude of the chert is: Strike N. 65 E, dip 15 to 40° N.

This deposit clearly shows the relationship between the red chert, the white chert and the manganese. It is possible that the iron in the red chert is being replaced by manganese, and that the lack of iron in the white chert may prevent precipitation within that rock.

Mining and metallurgy: The ore is the siliceous type, common to this type of manganese deposits and can probably not be concentrated. The deposit must be blasted, but the "ore" can be handled easily from the talus slope below.

An assay of typical "ore" from the deposit gave: Mn 29.8%