

Request for Amendment 5 for the Biglow Canyon Wind Farm

Exhibit G. Fish and Wildlife Habitat

**Submitted to
Oregon Energy Facility Siting Council**

Prepared for



Portland General Electric Company

Prepared by



October 2026

Table of Contents

1.0	Introduction.....	1
1.1	Analysis Area	2
1.2	Agency Consultation	2
2.0	Description of Biological and Botanical Surveys – OAR 345-022-0060(3)(a)	2
2.1	Information Review	3
2.2	Field Surveys.....	4
2.2.1	Habitat Categorization and Special Status Wildlife Surveys	4
2.2.2	Botanical Surveys	5
2.2.3	Raptor Nest Surveys.....	6
2.2.4	Wetlands and Waters Surveys.....	7
3.0	Identification of Habitat – OAR 345-022-0060(3)(b)(c).....	7
3.1	ODFW Habitat Categorization.....	7
3.2	Description of Fish and Wildlife Habitat in the Analysis Area	8
3.3	Quantity of Habitat Types by Habitat Category within the Analysis Area	9
4.0	Identification of State Sensitive Species and ODFW Site-Specific Issues of Concern – OAR 345-022-0060(3)(d).....	10
4.1	State Sensitive Species.....	10
4.2	ODFW Site-Specific Issues of Concern.....	17
5.0	Baseline Survey of Habitat Use by State Sensitive Species – OAR 345-022-0060(3)(e).....	17
5.1	Results of Field Surveys	17
5.1.1	2024 Habitat Categorization and Special Status Wildlife Surveys	17
5.1.2	2024 Botanical Surveys.....	18
5.1.3	Raptor Nest Surveys.....	19
5.1.4	Wetlands and Waters Surveys.....	19
6.0	Description of Potential Adverse Impacts – OAR 345-022-0060(3)(f)	19
6.1	Potential Impacts to Fish and Wildlife	20
6.2	Potential Impacts to State Sensitive Species.....	21
6.2.1	Mammals.....	21
6.2.2	Birds.....	22
6.2.3	Reptiles and Amphibians	29
7.0	Avoidance and Mitigation Measures – OAR 345-022-0060(3)(g).....	30

7.1	Project Design.....	30
7.2	Construction and Operation	31
7.3	Environmental Training and Sensitive Resource Awareness	32
7.4	Avian Protection.....	32
7.5	Mitigation	32
8.0	Monitoring Program – OAR 345-022-0060(3)(h).....	33
9.0	Conclusion	33
10.0	References	34
11.0	Approval Standards and Submittal Requirements	39

List of Tables

Table G-1. Summary of Field Surveys Conducted in 2024 and 2026.....	4
Table G-2. ODFW Habitat Categorization	8
Table G-3. Description of Habitat Types within the Analysis Area	9
Table G-4. Acres of Habitat Type by ODFW Habitat Categories within the Analysis Area and Site Boundary	10
Table G-5. State Sensitive Species with the Potential to Occur within the Analysis Area	12
Table G-6. Disturbance by Habitat Category and Land Cover Type	20
Table G-7. Approval Standards and Submittal Requirements Matrix	39

List of Figures

Figure G-1. Fish and Wildlife Analysis Area
Figure G-2. 2024-2026 Survey Area in the Facility Site Boundary
Figure G-3. Habitat Categories in the Analysis Area
Figure G-4. Habitat Subtypes in the Analysis Area
Figure G-5. 2024-2026 Special Status Species Observations in the Analysis Area and Its Vicinity

List of Attachments

Attachment G-1. Biological Survey Reports (Provided under Confidential Cover)
Attachment G-2. Draft Habitat Mitigation Plan

Acronyms and Abbreviations

Acronym/Abbreviation	Definition
AC	alternating current
APLIC	Avian Powerline Interaction Committee
BCWF or Facility	Biglow Canyon Wind Farm
BESS	battery energy storage system
BGEPA	Bald and Golden Eagle Protection Act
BIGL 2 or Project Developer	BIGL 2, LLC
Certificate Holder or PGE	Portland General Electric Company
Council or EFSC	Oregon Energy Facility Siting Council
gen-tie	generation tie
HMP	Habitat Mitigation Plan
kV	kilovolt
MW	megawatt
MWh	megawatt-hour
NHD	National Hydrography Dataset
NWI	National Wetlands Inventory
OAR	Oregon Administrative Rules
ODA	Oregon Department of Agriculture
ODFW	Oregon Department of Fish and Wildlife
ODOE	Oregon Department of Energy
ORBIC	Oregon Biodiversity Information Center
PWCA	Priority Wildlife Connectivity Area
PV	photovoltaic
RFA	Request for Amendment
SGCN	Species of Greatest Conservation Need
Site Certificate	Site Certificate for Amendment 4
Solar Facility	solar energy generation and battery storage components
SWAP	Oregon State Wildlife Action Plan
WMMP	Wildlife Monitoring and Mitigation Plan

1.0 Introduction

The Portland General Electric Company (PGE or Certificate Holder) submits this Request for Amendment (RFA) 5 to the Site Certificate on Amendment 4, issued March 27, 2026 (Site Certificate) for the Biglow Canyon Wind Farm (BCWF or Facility) to add a second phase of solar photovoltaic (PV) energy generation and battery storage capacity to the BCWF following the solar power generation and battery storage capacity approved under RFA 4.

BCWF, owned and operated by PGE, is located within an approved site boundary comprising approximately 25,000 acres, approximately 2.5 miles northeast of the town of Wasco in Sherman County, Oregon (Background Exhibit Figure 1). The BCWF operates under the Site Certificate from the Oregon Energy Facility Siting Council (Council or EFSC) as administered by the Oregon Department of Energy (ODOE). BCWF currently consists of 217 operating wind turbines, with a maximum blade tip height of 445 feet, and a peak generating capacity of 450 megawatts (MW) (Wind Facility). RFA 4, approved March 27, 2026, added 125 MW alternating current (AC) generating capacity from PV solar arrays and 500 megawatt-hours (MWh) in battery storage capacity. The solar energy generation and battery storage components (Solar Facility) described in RFA 4 are currently under construction.

In RFA 5, PGE proposes to add up to 200 MW AC generating capacity from PV solar arrays and 1,000 MWh in battery storage capacity (RFA 5 Solar Facility) by expanding the overall BCWF site boundary by approximately 839 acres. The RFA 5 Solar Facility is located across approximately 622 acres of the existing BCWF site boundary and 839 acres of expanded site boundary, for a total of approximately 1,461 acres (together, RFA 5 Site Boundary; see Background Exhibit Figure 4).

The RFA 5 Site Boundary provides a conservative estimate of the maximum area needed for development and micrositing through final design and construction for the RFA 5 Solar Facility, rather than the anticipated disturbance footprint. Primary infrastructure includes solar arrays, inverters, battery energy storage system (BESS) facilities and their subcomponents (i.e., inverters), one collector substation, and medium-voltage collector lines (herein collectively referred to as “Solar Components”). Additionally, the Solar Components include operations and maintenance (O&M) structures, site access roads, internal roads, perimeter fencing, facility entry gates, and temporary laydown areas. The maximum generating capacity from the Solar Components will be 200 MW AC. Power generated from the Solar Components will be transferred to the grid via a 230-kilovolt (kV) generation tie (gen-tie) transmission line, approximately 2.7 miles long. The RFA 5 Solar Facility includes the Solar Components and the gen-tie transmission line.

PGE will own and operate the RFA 5 Solar Facility as a part of the BCWF, which, to date, has been developed by BIGL 2, LLC (BIGL, or Project Developer). BIGL 2, in its capacity as the Project Developer, supports PGE in this RFA 5 and may construct and temporarily operate the RFA 5 Solar Facility on behalf of PGE under a Build-Transfer Agreement.

The information contained herein supports demonstration of compliance with the Fish and Wildlife Habitat approval standard for Oregon Administrative Rule (OAR) 345-022-0060.

1.1 Analysis Area

In accordance with OAR 345-001-0010(35)(c), the analysis area for fish and wildlife habitat and species consists of the RFA 5 Site Boundary and the area 0.5 miles from the RFA 5 Site Boundary (Figure G-1). Portions of the RFA 5 Site Boundary are designated as the Solar Components, which includes the anticipated disturbance footprint of the Solar Components (Solar Siting Area) and the gen-tie transmission line (Transmission Corridor). The RFA 5 Site Boundary was initially planned as part of RFA 4 and was removed from consideration prior to receiving the Site Certificate on Amendment 4; therefore, surveys referenced in this RFA 5 Exhibit G may also discuss species that occur within the vicinity of the Analysis Area (more than 0.5 miles but less than 5 miles from the RFA 5 Site Boundary).

1.2 Agency Consultation

For the BCWF RFA 4, which at the time included what is currently the RFA 5 Site Boundary, PGE had consulted with the Oregon Department of Fish and Wildlife (ODFW) regarding the appropriate protocols for documenting the presence of state sensitive species (inclusive of state threatened and endangered species) and the classification of fish and wildlife habitat. An email was sent to ODFW on May 22, 2024, by PGE with a biological study plan associated with the RFA 4 Site Boundary and additional lands that were subsequently omitted from RFA 4 (including the RFA 5 Site Boundary as well as lands omitted from consideration from development). ODFW responded on June 6, 2024, approving of the biological study plan and provided additional recommendations for siting of the Solar Components. The background review and field survey methods specific to botanical surveys were submitted to Oregon Department of Agriculture (ODA) for review on June 10, 2024. ODA provided concurrence with the methods on June 13, 2024, prior to Tetra Tech conducting field surveys.

On April 7, 2026, Tetra Tech, on behalf of PGE, sent an email to ODFW and ODOE with a pre-construction raptor nest survey plan for the solar components described in RFA 4. ODFW and ODOE approved of the survey methodology on April 20, 2026.

2.0 Description of Biological and Botanical Surveys – OAR 345-022-0060(3)(a)

(3) To assist the Council in determining whether the standard outlined in (1) through (2) has been met, the Applicant must submit information about the fish and wildlife habitat and the fish and wildlife species, other than the species addressed in OAR-022-0070(3) (the Threatened and Endangered Species Exhibit) that could be affected by the proposed facility, providing evidence to support a finding by the Council as required by this rule. The applicant must include:

(a) A description of biological and botanical surveys performed that support the information in this exhibit, including a discussion of the timing and scope of each survey;

Fish and wildlife species that might be affected by the Solar Components were identified through review of the original application materials for BCWF (Orion Sherman County Wind Farm, LLC 2005); the application materials for BCWF RFA 4, which included the RFA 5 Site Boundary (PGE 2025); and a desktop review of the RFA 5 Site Boundary in 2026. The 2026 desktop review was conducted to assess updates to reference databases and the recently published Oregon State Wildlife Action Plan (SWAP; ODFW 2026a), which replaced the Oregon Conservation Strategy and included new Species of Greatest Conservation Need (SGCN) and species that are pending sensitive status. A detailed summary of the desktop review and the results of the 2024 surveys are provided below.

2.1 Information Review

Tetra Tech conducted a desktop assessment including a review of existing information regarding habitat types and special status species with the potential to occur within 0.5 miles of the RFA 5 Site Boundary. The term “special status species” includes federal and state endangered, threatened, proposed, and candidate fish, wildlife, and plant species, as well as state sensitive wildlife species and SGCN as determined by the SWAP. Special status species also include eagles. While surveys were performed with an emphasis on special status species, this exhibit focuses on a subset of that group consisting of only state sensitive species and SGCN.¹

Tetra Tech reviewed ODFW’s sensitive species list and the SWAP to identify state sensitive species and SGCN with the potential to occur within the analysis area based on the ecoregion (ODFW 2021, ODFW 2026a) and then removed any that had no documented occurrences or modeled range in Sherman County (OSULP and INR 2021, ODFW 2026b, ORBIC 2023, ORBIC 2025). Species were further excluded for consideration if no potential habitat was present within the analysis area (ODFW 2026a, ODFW 2026b, NatureServe 2026, OregonFlora 2026, eBird 2026, StreamNet 2026, Xerces 2026, Xerces et al. 2021, Attachment G-1). Oregon Biodiversity Information Center (ORBIC) data (ORBIC 2024) were then reviewed to determine known species occurrence locations within the analysis area. The results of this desktop review and the results from field surveys that verified available habitat for wildlife species and provided further documentation of sensitive species within the analysis area are discussed in Section 4.1.

Other resources consulted during the desktop review included the National Land Cover Database, aerial photography, the ODFW Compass mapping tool (ODFW 2026b), as well as results from surveys conducted in support of RFA 4 (Attachment G-1) and pre- and post-construction surveys performed for the BCWF (Orion Sherman County Wind Farm 2005, WEST 2005, WEST 2009, WEST 2010, WEST 2012a, WEST 2012b, WEST 2012c, WEST 2013, PGE 2023). Exhibit D addresses the

¹ The Oregon SWAP was published in early 2026 and largely overlaps with the current state sensitive species list (ODFW 2021). The SWAP includes several species that are pending state sensitive status which will likely be implemented in the next update to the state sensitive species list, anticipated to occur in 2026.

Sherman County Comprehensive Plan (Goal 5), which includes the potential for natural resources to occur within the RFA 5 Site Boundary.

2.2 Field Surveys

Table G-1 provides a summary of biological field surveys conducted at the RFA 5 Site Boundary in 2024 and 2026 within portions of the Analysis Area (Figure G-2). The survey reports are included in Attachment G-1, except for the wetlands and waters survey report, which is included in Exhibit O, Attachment O1-1.

Table G-1. Summary of Field Surveys Conducted in 2024 and 2026

Survey	Survey Timing	Reference	Extent
Habitat Categorization and Special Status Wildlife Surveys	June 2024	Attachment G-1 ¹	Habitat Categorization Survey Area (combined RFA 4 and RFA 5 Site Boundaries and additional lands that have since been omitted from this RFA 5)
Botanical Surveys	July 2024		Botanical Survey Area (combined RFA 4 and RFA 5 Site Boundaries and additional lands that have since been omitted from this RFA 5)
Raptor Nest Surveys	June 2024, May 2026		0.5-mile buffer of combined RFA 4 and RFA 5 Site Boundaries and additional lands that have since been omitted from this RFA 5. Pre-construction survey in 2026 encompassed the RFA 4 Site Boundary plus a 0.5-mile buffer.
Wetlands and Waters Surveys	April and May 2024	Exhibit O, Attachment O1-1	Wetland Study Area (approximately 1,167 acres) encompassing RFA 4 Site Boundary and RFA 5 Site Boundary.
1. The pre-construction raptor nest survey report contains confidential material and will be provided under separate cover.			

2.2.1 Habitat Categorization and Special Status Wildlife Surveys

Habitat categorization and special status wildlife surveys were performed in June 2024 by Tetra Tech within the RFA 5 Site Boundary and additional lands that have since been omitted (Attachment G-1). The survey area covered 4,636 acres including the entire RFA 5 Site Boundary, the RFA 4 Solar Micrositing Area, and some additional lands that have since been omitted from consideration. The objective of the surveys was to categorize habitats per the habitat categories set forth in ODFW's Fish and Wildlife Mitigation Policy (OAR 635-415-0025), consistent with the habitat categorization developed during permitting of the BCWF, and to document the presence of special status wildlife species.

During the habitat categorization survey, habitat categories established by the ODFW Fish and Wildlife Habitat Mitigation Policy were used to rank the habitat subtypes present within the RFA 5 Site Boundary. Habitat mapping was performed using a minimum of 1-acre mapping unit (except for specialized habitat such as cliffs or rock outcrops). These surveys were conducted by a biologist

familiar with vegetation of the Columbia Plateau Ecoregion and were scheduled during the growing season to aid in identification of vegetative species. Meandering transects were walked by a biologist to categorize the composition and structure of habitat by noting the dominant vegetation, presence of large trees, existing disturbance, or other habitat features such as shrub and grass composition and density. Each polygon was then assigned a habitat type, subtype, and habitat quality category that was guided by the habitat categorization for the BCWF.

For the portions of the analysis area that were not covered by the survey, habitat mapping is based on a desktop analysis using aerial photography interpretation and the Certificate Holder's familiarity with the site acquired through operation of the BCWF.

Special status wildlife surveys were conducted concurrently with habitat categorization surveys. The objective of the special status wildlife survey was to document the presence of any special status species within the survey area, as well as general wildlife species. During surveys, the biologist walked meandering transects through nonagricultural areas, scanning and listening for wildlife species and recognizable signs of wildlife (e.g., scat, tracks, burrows and nests). Agricultural areas were surveyed by visually scanning and listening for wildlife from accessible roads or bordering non-agricultural lands. The biologist documented the location of special status species, or recognizable sign, and recorded the number of individuals and their behavior, if observed. Special habitats and unique features (i.e., cliffs, ponds, talus slopes, rock outcrops, raptor nests, and big game) were also documented if encountered. Following the survey, the digitized data were downloaded and processed using geographic information system software and reviewed for quality control and assurance.

2.2.2 Botanical Surveys

Botanical surveys were conducted on July 9 and 10, 2024 (Attachment G-1). The purpose of the botanical surveys was to document the presence of federally or state listed endangered, threatened, and candidate vascular plant species and state and county-designated noxious weeds within the RFA 5 Site Boundary. The surveys were designed in accordance with Council standards set forth in OAR 345-022-0060 and OAR 345-022-0070², and in coordination with the ODA.

Prior to field surveys, Tetra Tech conducted a review of existing information to identify federal and state endangered, threatened, proposed, and candidate plant species with the potential to occur. No federally listed, proposed, or candidate plant species have the potential to occur within or near the RFA 5 Site Boundary; however, one state endangered species, one state threatened species, and four state candidate species have the potential to occur in Sherman County. No suitable habitat exists within the RFA 5 Site Boundary for these species.

² The surveys were designed in accordance with the Council standards that were previously set forth in OAR 345-021-0010(1)(p) and (q) but have since moved to OAR 345-022-0060 and OAR 345-022-0070 on April 2, 2025.

Tetra Tech reviewed lists of species designated as noxious weeds in Oregon and Sherman County (ODA 2022³; SCNWD 2024). Existing literature and other sources were also reviewed to familiarize surveyors with identification of designated noxious weeds that would potentially be encountered within the survey area.

Tetra Tech conducted botanical field surveys using the Intuitive Controlled survey method, a standard and commonly accepted survey protocol (USFS and BLM 1998). This method incorporates meandering transects that traverse the survey area and targets the full array of major vegetation types, aspects, topographical features, habitats, and substrate types. While en route, the surveyors searched for target species, and when the surveyors arrived at an area of high potential habitat, they conducted a complete survey for the target species. The botanist maintained a running list of vascular plant species encountered and made informal collections of unknown species for later identification. Data collected for any rare plant population, if encountered, would include species phenology, number of plants observed, habitat information and associated species, and any visible threats such as development, grazing, erosion, or noxious weeds.

2.2.3 Raptor Nest Surveys

PGE has performed raptor nest surveys at regular intervals during operation of the BCWF, with the most recent survey being from 2022 (PGE 2023). Prior to conducting surveys, Tetra Tech reviewed available raptor nest data from the most recent 2022 raptor nest survey, which extended out 2 miles from the BCWF and encompassed the entire RFA 5 Site Boundary and Exhibit G Analysis Area. The survey area of the 2024 raptor nest survey covered the RFA 4 Site Boundary, RFA 5 Site Boundary, and a 0.5-mile buffer, plus additional areas not addressed in this RFA 5 (Attachment G-1).

The 2024 survey was performed from the ground in mid-June. The biologist checked the status of all known nests documented during the 2022 PGE survey that occurred within the survey area and searched for new nests. Data recorded for each raptor nest included the date; time; unique nest identification; Universal Transverse Mercator coordinates; species occupying the nest; nest status; and nest condition. Nest status (active or inactive) was determined using a combination of cues such as adult behavior (e.g., diving, calling, nest building/repair, incubation, brooding, or feeding of young), presence of eggs or young, or whitewash.

A pre-construction survey was conducted in 2026 to support the construction of the solar components approved under Amendment 4 of the Site Certificate for BCWF. The survey area consisted of the RFA 4 Site Boundary plus a 0.5-mile buffer, which overlaps with the northern portion of the RFA 5 Site Boundary and Analysis Area described in this RFA 5 (see Attachment G-1, 2026 raptor nest survey report provided under separate cover).

³ The ODA Noxious Weed Policy and Classification System was revised in 2024, after initial desktop review was completed. Survey data were reviewed against the 2024 ODA Noxious Weed Policy and Classification System to support this RFA 5 Exhibit G, as described in Section 5.1.2.

2.2.4 Wetlands and Waters Surveys

Tetra Tech conducted pedestrian surveys to delineate wetlands and other waters on July 9 to 12, 2024 (Attachment O1-1). The survey area covered the entirety of the RFA 4 Site Boundary, RFA 5 Site Boundary, and additional lands that have since been omitted. Prior to the surveys, Tetra Tech reviewed the National Wetlands Inventory (NWI), National Hydrography Dataset (NHD), hydric soils data, and historical aerial photographs in Google Earth to identify potential wetlands and other waters prior to performing field work. Wetlands and surface water data were obtained from the U.S. Fish and Wildlife Service NWI (USFWS 2024), which includes NWI and miscellaneous wetland mapping by state and federal agencies, non-governmental organizations, academia and consultants, and from the U.S. Geological Survey NHD (USGS 2024). Soils data were also obtained from the National Resources Conservation Service Web Soil Survey (NRCS 2024). Tetra Tech used aerial imagery from Google Earth because a wide variety of imagery was available.

The desktop wetland data were used to focus the wetland delineation's field effort while the desktop surface water data were used to focus the non-wetlands water evaluation as necessary.

Data collected during wetlands and waters surveys informed habitat categorization and helped with the determination of state sensitive species presence described in this exhibit. Results from the 2024 wetlands and waters surveys are summarized in Exhibit O and presented in the corresponding survey report (Attachment O1-1).

3.0 Identification of Habitat – OAR 345-022-0060(3)(b)(c)

(b) Identification of all fish and wildlife habitat in the analysis area, classified by the general fish and wildlife habitat categories as set forth in OAR 635-415-0025 and the sage-grouse specific habitats described in the Greater Sage-Grouse Conservation Strategy for Oregon at OAR 635-140-0000 through 635-140-0025 (core, low density, and general habitats), and a description of the characteristics and condition of that habitat in the analysis area, including a table of the areas of permanent disturbance and temporary disturbance (in acres) in each habitat category and subtype;

(c) A map showing the locations of the habitat identified in (b);

3.1 ODFW Habitat Categorization

The ODFW Fish and Wildlife Habitat Mitigation Policy (OAR 635-415-0015) provides a framework for assigning one of six category types to habitats based on the relative importance of these habitats to fish and wildlife species. The definition of each category type, as well as an example of each category type within the analysis area, is shown in Table G-2. The Solar Components will not be located in sage-grouse habitat, and therefore the Greater Sage-Grouse Conservation Strategy for Oregon does not apply.

Table G-2. ODFW Habitat Categorization

ODFW Habitat Category	Definition ¹	Examples of ODFW Habitat Categories within Analysis Area
1	Irreplaceable, essential habitat for a fish or wildlife species, population, or a unique assemblage of species and is limited on either a physiographic province or site-specific basis, depending on the individual species, population or unique assemblage.	None identified in the Analysis Area.
2	Essential habitat for a fish or wildlife species, population, or unique assemblage of species and is limited either on a physiographic province or site-specific basis depending on the individual species, population or unique assemblage.	None identified in the Analysis Area.
3	Essential habitat for fish and wildlife, or important habitat for fish and wildlife that is limited either on a physiographic province or site-specific basis, depending on the individual species or population.	<ul style="list-style-type: none"> • Upland Trees • Intermittent Streams • Grasslands • Freshwater Emergent Wetlands
4	Important habitat for fish and wildlife species.	<ul style="list-style-type: none"> • Grasslands
5	Habitat for fish and wildlife having high potential to become either essential or important habitat.	None identified in the Analysis Area.
6	Habitat that has low potential to become essential or important habitat for fish and wildlife.	<ul style="list-style-type: none"> • Agricultural Areas – These areas include irrigated and non-irrigated cropland. • Developed Areas – These areas include existing infrastructure and developed areas.
1. Source: OAR 635-415-0025.		

3.2 Description of Fish and Wildlife Habitat in the Analysis Area

The analysis area covers approximately 5,824 acres in northern Sherman County, Oregon, in the Columbia Plateau. The analysis area comprises primarily agricultural lands with small areas of remnant grassland and woodland habitat with elevations ranging from approximately 420 to 1,600 feet above sea level. Vegetation within much of the analysis area has been modified due to historic and current agricultural activity, construction and management of the Facility, and historic and current grazing activity. Non-native, invasive grasses and forbs are common throughout the analysis area due to these land uses. Only limited areas of native habitat types occur, and even in those areas, non-native, invasive grasses and forbs, including cheatgrass (*Bromus tectorum*), bulbous bluegrass (*Poa bulbosa*), cereal rye (*Secale cereale*), and prickly lettuce (*Lactuca serriola*), are abundant.

The analysis area does not overlap with Priority Wildlife Connectivity Areas (PWCAs) or any big game winter range habitat (ODFW 2013, ODFW 2023); therefore, these special habitat areas will not be discussed further in this RFA 5 Exhibit G.

Habitat types and subtypes within the analysis area are described in Table G-3 and displayed in Figure G-2.

Table G-3. Description of Habitat Types within the Analysis Area

Habitat Type	Habitat Sub-Type	Description
Agriculture	Irrigated Cropland	Cultivated crops and associated disturbed areas. The analysis area overlaps with one small portion of irrigated barley in the northeast.
	Non-Irrigated Cropland	Cultivated crops and associated disturbed areas. Non-irrigated cropland is mostly dryland wheat and barley.
Developed	Developed	Paved and gravel roads, gravel quarries, residences, equipment staging, permanent footprint of existing wind facility.
Grassland-steppe	Grasslands	Weedy grasslands with minimal shrub component and trace forbs. Most grasslands within the analysis area are dominated by cheatgrass (<i>Bromus tectorum</i>).
Surface Water	Intermittent Streams	NHD data.
Upland	Upland Trees	Consists mostly of live non-native black locust (<i>Robinia pseudoacacia</i>) trees and snags containing an understory of weedy annual grasses and native shrubs.
Wetland	Freshwater Emergent Wetland	Two delineated wetlands and associated ponds at the bottom of the gravel quarries.

3.3 Quantity of Habitat Types by Habitat Category within the Analysis Area

Table G-4 shows the acreages within the analysis area and the RFA 5 Site Boundary of each habitat type with the assigned habitat category. The locations of each habitat type by category within the analysis area are shown on Figures G-3 and G-4.

The most prominent habitat category in the Analysis Area is Category 6 (99 percent). Category 6 includes Agriculture and Developed habitat types, which are dominated by Non-irrigated Cropland subtypes but also include Irrigated Cropland and Developed subtypes. The remaining 1 percent of the Analysis Area consists of Category 3 Upland, Surface Water, Grassland-steppe, and Wetlands, and Category 4 Grassland-steppe habitat types. Representative photos of the analysis area are included in the Habitat Categorization and Special Status Wildlife Survey Report (Attachment G-1).

Presence of a particular habitat category within the analysis area does not indicate that this habitat will necessarily be impacted by the Solar Components. Section 6.1 presents the areas of permanent disturbance and temporary disturbance (in acres) in each habitat category and habitat type with discussion on potential impacts to fish and wildlife habitat.

Table G-4. Acres of Habitat Type by ODFW Habitat Categories within the Analysis Area and Site Boundary

ODFW Habitat Category	Habitat Type	Habitat Sub-Type	Acres in Analysis Area ¹	Acres in Site Boundary ¹
3	Upland	Upland Trees	26.4	19.1
	Surface Water	Intermittent Streams	25.4	-
	Grassland-steppe	Grasslands	1.1	<1
	Wetland	Freshwater Emergent Wetland	0.4	-
Category 3 Total			53.2	19.1
4	Grassland-steppe	Grasslands	17.1	3.1
Category 4 Total			17.1	3.1
6	Agriculture	Non-Irrigated Cropland	5,645.5	1,434.9
	Developed	Developed	92.2	3.6
	Agriculture	Irrigated Cropland	15.6	-
Category 6 Total			5,753.3	1,438.5
Grand Total			5,823.6	1,460.8
Note: Numbers may not sum correctly due to rounding. "-" means no acres while <1 means greater than zero but less than 0.5 acres present within the analysis area and the RFA 5 Site Boundary. 1. Acres represent area within the analysis area and the RFA 5 Site Boundary, not impact areas. Impacts are discussed in Section 6.0.				

4.0 Identification of State Sensitive Species and ODFW Site-Specific Issues of Concern – OAR 345-022-0060(3)(d)

(d) Based on consultation with the Oregon Department of Fish and Wildlife (ODFW) and appropriate field study and literature review, identification of all State Sensitive Species that might be present in the analysis area and a discussion of any site-specific issues of concern to ODFW;

The Certificate Holder identifies all state sensitive species, in addition to SGCN, below in compliance with OAR 345-022-0060(3)(d).

4.1 State Sensitive Species

Based on the desktop analysis and field surveys, 35 state sensitive and/or SGCN species and 1 protected eagle species have potential to occur in the analysis area (Table G-5). Bald eagles (*Haliaeetus leucocephalus*) are addressed in Exhibit G as species of concern protected under the Bald and Golden Eagle Protection Act (BGEPA). Golden eagles (*Aquila chrysaetos*), also protected under the BGEPA, were included in the 2026 SWAP as an SGCN and are pending sensitive status in Oregon (ODFW 2026a). Of the 35 state sensitive species that have potential to occur in the analysis area, 7 are sensitive-critical and 11 are sensitive in the Columbia Plateau Ecoregion (ODFW 2021). Another 10 species are pending sensitive status and may or may not be considered sensitive or sensitive-critical in

the Columbia Plateau Ecoregion when their status is updated (ODFW 2026a). The remaining 7 species are, at this time, only considered SGCN in the Columbia Plateau Ecoregion, though they are considered sensitive or sensitive-critical in other ecoregions (ODFW 2021, ODFW 2026a). The 35 state sensitive and/or SGCN species include 14 mammals, 15 birds, 2 reptiles, 1 amphibian, and 3 invertebrate species. A query of the StreamNet database (StreamNet 2026) and the Certificate Holder's request for information from ORBIC (ORBIC 2024) did not identify any state sensitive fish species with potential to occur in the analysis area. Therefore, no fish are included in Table G-5 below or addressed in this Exhibit G. State threatened and endangered species are addressed in Exhibit H.

Table G-5. State Sensitive Species with the Potential to Occur within the Analysis Area

Common Name	Scientific Name	State Status ¹ (Columbia Plateau Ecoregion)	Expected Habitats	Observed or Expected Occurrence within Analysis Area ²	Potential Use of Habitat Within the Analysis Area
Mammals					
California myotis	<i>Myotis californicus</i>	SGCN	Inhabits various habitats, including sea coasts, desert scrub, oak-juniper woodlands, montane and humid coastal forests, mountain meadows, canyons, riparian woodlands, grasslands, rural residential areas, and towns. Roosts in a wide range of sheltered sites, including trees, shrubs, mines, caves, bridges, and buildings.	Moderate. Recorded in Sherman County and predicted habitat exists within the analysis area. Has not been detected during fatality monitoring at the BCWF.	Forage: Upland Trees, aquatic habitats, existing structures, and their vicinities. Roosting: Upland Trees, existing structures.
Canyon bat	<i>Parastrellus hesperus</i>	PS, SGCN	Inhabits deserts, woodlands, and shrublands. Roosts among boulders or in cracks and crevices of rock faces. Highly associated with canyon and cliff habitat, and also found in caves and mines. Foraging occurs in canyons, along cliffs, in riparian zones, and over lava beds. Foraging habitat must be in close proximity to roost sites.	Low. Predicted to occur in Sherman County but this species prefers canyon and cliff habitats, which do not occur within the analysis area. This species may only pass through the analysis area. Not detected during fatality monitoring at the BCWF.	Forage: Upland Trees, aquatic habitats, existing structures, and their vicinities. Roosting: Upland Trees, existing structures.
Hoary bat	<i>Lasiurus cinereus</i>	S, SGCN	Generally associated with forested habitat. Often found in coniferous forests east of the Cascade Range. Species prefers late-successional conifer forests for roosting. Often forages along riparian corridors and forest openings or forest borders, where it can easily fly. Sometimes found in open cover types like grasslands and meadows, especially when migrating or foraging.	High, during migration. Recorded in Sherman County and predicted habitat within analysis area. Frequently found during fatality monitoring at the BCWF.	Forage: Upland Trees, aquatic habitats, existing structures, and their vicinities. Roosting: Upland Trees, existing structures.
Little brown myotis	<i>Myotis lucifugus</i>	PS, SGCN	Occupies a wide range of habitat types and roosting structures, including trees with exfoliating bark, snags, rock crevices/cliffs, caves/mines, wood piles, and buildings in urban areas. Limited by available roosting habitat and/or closure/disturbance at hibernacula and maternity colonies.	Moderate. Has been recorded during acoustic surveys in similar habitat just east of the John Day River. This species has not been detected during fatality monitoring at the BCWF.	Forage: Upland Trees, aquatic habitats, existing structures, and their vicinities. Roosting: Upland Trees, existing structures.
Long-legged myotis	<i>Myotis volans</i>	SGCN	Highly associated with montane forests, but can also be found in desert and riparian habitats. Often associated with late successional conifer forests or other forested habitat with late successional components (especially snags). Forages in forest riparian areas and along forest edges. This species often uses tall, large-diameter snags as roosts, but may also use caves, mines, bridges, and rock crevices as day roosts, night roosts, and hibernacula.	Low. Recorded in Sherman County but this species’ preferred habitat does not occur within the analysis area. Not detected during fatality monitoring at the BCWF.	Forage: Upland Trees, aquatic habitats, existing structures, and their vicinities. Roosting: Upland Trees, existing structures.
North American porcupine	<i>Erethizon dorsatum</i>	PS, SGCN	Coniferous and mixed forests, but also may be found in riparian zones, grasslands, shrublands, and deserts. Winter dens may be in rock outcrops, live hollow trees, hollow logs, or outbuildings. Known to forage in agricultural fields.	High. North American porcupine was observed during wildlife surveys for RFA 4 and during pre- and post-construction surveys at BCWF. Habitat within the analysis area is not high quality for this species and individuals may only pass through the analysis area to more suitable habitat; however, porcupine may rely on winter wheat and barley when other food sources are limited.	Forage: All habitat types. Den: Upland Trees.
Pallid bat	<i>Antrozous pallidus</i>	S, SGCN	Lives in dry, open habitat such as grasslands, shrub-steppe, and dry forest ecotones. Uses crevices in cliffs, caves, mines, or bridges (and sometimes, buildings) for day, night, or maternity roosts, or hibernacula.	Moderate. Recorded in Sherman County and predicted habitat within analysis area. Not detected during fatality monitoring at the BCWF.	Forage: Upland Trees, aquatic habitats, existing structures, and their vicinities. Roosting: Upland Trees, existing structures.
Silver-haired bat	<i>Lasionycteris noctivagans</i>	S, SGCN	Relies on late-successional conifer forests but may be found in other habitat types during migration.	High, during migration. Recorded in Sherman County and predicted habitat within analysis area. Frequently found during fatality monitoring at the BCWF.	Forage: Upland Trees, aquatic habitats, existing structures, and their vicinities. Roosting: Upland Trees, existing structures.

Common Name	Scientific Name	State Status ¹ (Columbia Plateau Ecoregion)	Expected Habitats	Observed or Expected Occurrence within Analysis Area ²	Potential Use of Habitat Within the Analysis Area
Spotted bat	<i>Euderma maculatum</i>	S, SGCN	Roosts in caves, cliffs, canyon walls, and trees near meadows. Forages in meadows, shrub-steppe, and along water sources.	Low. Recorded in Sherman County but no predicted habitat within analysis area. Not detected during fatality monitoring at the BCWF.	Forage: Upland Trees, aquatic habitats, existing structures, and their vicinities. Roosting: Upland Trees, existing structures.
Townsend’s big-eared bat	<i>Corynorhinus townsendii</i>	SC, SGCN	Occurs in a variety of habitats but most likely to be found near caves, mines, or other roosting areas such as hollow trees, bridges and isolated buildings. Forages in a variety of habitats, including arid pine forests, sage-steppe, and stream corridors.	Moderate. Recorded in Sherman County and predicted habitat within analysis area. Not detected during fatality monitoring at the BCWF.	Forage: Upland Trees, aquatic habitats, existing structures, and their vicinities. Roosting: Upland Trees, existing structures.
Western long-eared myotis	<i>Myotis evotis</i>	PS, SGCN	Occupies a wide range of habitat types and roosting structures, including large trees and snags, caves and mines, and cliffs. Prefers to forage near water. Sensitive to disturbance at roost sites.	Low. Recorded in Sherman County but preferred habitat does not exist within the analysis area. Not detected during fatality monitoring at the BCWF.	Forage: Upland Trees, aquatic habitats, existing structures, and their vicinities. Roosting: Upland Trees, existing structures.
Western small-footed myotis	<i>Myotis ciliolabrum</i>	PS, SGCN	Strongly associated with habitat occurring near cliffs or rocky structures, or caves/mines. Prefers roosting in crevices. Can forage in complex habitats. Sensitive to loss of habitat and disturbance at roosts.	Low. Recorded in Sherman County but preferred habitat does not exist within the analysis area. Not detected during fatality monitoring at the BCWF.	Forage: Upland Trees, existing structures, and their vicinities. Roosting: Upland Trees, existing structures.
White-tailed jackrabbit	<i>Lepus townsendii</i>	SGCN	Requires open areas in bunch grass and rabbitbrush dominated habitats. Avoids forested areas.	Moderate. Habitat within the analysis area is low quality, but this species has been observed during pre- and post-construction surveys at BCWF.	Forage: All habitat types. Den: Grasslands, Upland Trees where canopy is thin, infrequently disturbed areas within and near non-irrigated cropland.
Yuma myotis	<i>Myotis yumanensis</i>	PS, SGCN	Occupies a wide range of habitats and roosting structures. Roosts can include caves, bridges, or other anthropogenic structures. Often found near bodies of water, such as streams, rivers, and ponds. Especially vulnerable to loss or disturbance of roosting habitat, hibernacula, and maternity sites.	Low. Recorded in Sherman County but preferred habitat does not exist within the analysis area. Not detected during fatality monitoring at the BCWF.	Forage: Upland Trees, aquatic habitats, existing structures, and their vicinities. Roosting: Upland Trees, existing structures.
Birds					
American avocet	<i>Recurvirostra americana</i>	PS, SGCN	Alkali and freshwater ponds, lakes, and wetlands with extensive shallow areas that provide critical foraging habitat. Prefers sites with minimal disturbance and rely on broad, open shorelines or islands with little to no vegetation for nesting.	Low. Recorded in Sherman County but no predicted habitat exists within the analysis area. May only pass through during migration.	Forage: Freshwater Emergent Wetlands. Nesting: None.
Bald eagle	<i>Haliaeetus leucocephalus</i>	None	Typically builds its nest in tall, sturdy trees near water sources, such as rivers, lakes, and coastlines where it can find fish, its primary food source. Known to scavenge opportunistically on carcasses in otherwise unsuitable habitat particularly during winter and migration.	High. Bald eagles are known to nest along the John Day River, but no suitable nesting habitat for bald eagles exists within the analysis area. No bald eagles or bald eagle nests detected during the surveys conducted for RFA 4. No bald eagle nests found during raptor nests surveys for the BCWF but bald eagles are regularly detected during avian use surveys for the BCWF. Use of the Solar Components is expected to be mostly limited to migration and winter, when the species sometimes feeds on carrion.	Forage: All habitat types. Nesting: None
Black tern	<i>Chlidonias niger</i>	PS, SGCN	Shallow freshwater marshes, flooded meadows, and wetlands with emergent vegetation. Often associated with bulrush, sedge rush, or tufted hairgrass/rush habitats with some standing water.	Low. The analysis area exists within this species’ range, but no suitable habitat exists within the analysis area. May only pass through during migration.	Forage: Freshwater emergent wetlands Nesting: None

Common Name	Scientific Name	State Status ¹ (Columbia Plateau Ecoregion)	Expected Habitats	Observed or Expected Occurrence within Analysis Area ²	Potential Use of Habitat Within the Analysis Area
Brewer’s sparrow	<i>Spizella breweri</i>	S, SGCN	Migrant and summer resident that uses sagebrush shrublands, generally with a canopy height greater than 5 feet. Typically nests in brush or in clumps of grass.	Moderate. Not detected during the 2024 surveys. Also not detected during avian use surveys for the BCWF, but one individual found during fatality monitoring for the BCWF. Limited suitable habitat within the analysis area.	Forage: Grasslands and agricultural areas. Nesting: None.
Burrowing owl (Western)	<i>Athene cunicularia hypugaea</i>	SC, SGCN	Occurs in open grassland and shrub-steppe habitats as well as highly disturbed habitats such as agricultural areas. Typically nests in burrows excavated by burrowing mammals. Spring and summer resident that travels south for warmer climates during the winter.	Moderate. Suitable habitat within the analysis area, and eBird sightings documented in the vicinity of the analysis area. This species was detected during pre-construction surveys for the BCWF.	Forage: Grasslands and agricultural areas. Nesting: All terrestrial habitats where burrows exist.
Common nighthawk	<i>Chordeiles minor</i>	S, SGCN	Long-distance migrant present in Oregon during its breeding season, from May to August. Inhabits any kind of open or semi-open terrain. Nesting habitat is characterized by open landscapes with little ground cover and is most abundant in sagebrush and rocky scablands and rimrock habitats of eastern Oregon. Tends to roost and nest on bare ground, including gravel roads.	Moderate. Suitable habitat within the analysis area, and eBird sightings documented in the vicinity of the Analysis Area. Not detected during the 2024 surveys or during avian use surveys, but one individual found during fatality monitoring for the BCWF.	Forage: All terrestrial habitat types. Nesting: All terrestrial habitat types except active agricultural fields.
Ferruginous hawk	<i>Buteo regalis</i>	SC, SGCN	Found in open, arid landscapes. Species often uses grasslands and shrub-steppe with scattered shrubs or trees for perching and nesting. Also nests on cliffs, rimrock, and hillsides.	High. Suitable nesting and foraging habitat within the analysis area. Not detected during the 2024 surveys. Low numbers have been detected during avian use surveys but no nests detected during raptor nest surveys for the BCWF.	Forage: All terrestrial habitat types. Nesting: Upland Trees, utility structures.
Golden eagle	<i>Aquila chrysaetos</i>	PS, SGCN	Inhabits a variety of habitats including forests, canyons, shrub lands, grasslands, and oak woodlands. Nests on cliffs, utility towers, or in large trees. Forages in open habitats.	High. Known historical nest territories along the nearby cliffs of the John Day River. Analysis area lacks nesting habitat but provides foraging habitat. Not detected during the 2024 surveys. Species is regularly detected during BCWF avian use surveys but no nests detected within survey area during raptor nest surveys for the BCWF.	Forage: All habitat types. Nesting: None
Grasshopper sparrow (Western)	<i>Ammodramus savannarum perpallidus</i>	S, SGCN	Migrant and summer resident east of the Cascades. Occurs is dry grassland habitat, generally with low to moderate grass height and low percent shrub cover.	Moderate. Suitable habitat within the analysis area and eBird sightings documented in the vicinity of the analysis area. Not detected during the 2024 surveys, but regularly detected during avian use surveys for the BCWF.	Forage: Grasslands, agricultural areas. Nesting: Grasslands.
Lewis’s woodpecker	<i>Melanerpes lewis</i>	SC, SGCN	Ponderosa pine forests, oak woodlands, oak-pine woodlands, cottonwood riparian forests, and areas burned by wildfires.	Low. Recorded in Sherman County but limited suitable habitat within the analysis area. Not detected during the 2024 surveys, or during pre- or post-construction surveys at the BCWF. Compass observations occur nearby in Wasco, Oregon; therefore, this species may pass through the analysis area during migration.	Forage: Upland Trees, residential areas (yards). Nesting: Within the vicinity of surface waters.
Loggerhead shrike	<i>Lanius ludovicianus</i>	S, SGCN	Found in a variety of open, brushy habitats, including agricultural areas. In Oregon, often uses tall sagebrush for nesting and roosting and open areas with grasses and significant bare ground for foraging.	High. Detected within the RFA 5 Site Boundary during 2024 special status wildlife surveys. Also detected during avian use surveys for the BCWF. Suitable nesting and foraging habitat within the analysis area.	Forage: All terrestrial habitats. Nesting: Upland Trees, residential areas (yards).
Long-billed curlew	<i>Numenius americanus</i>	SC, SGCN	Prefers open habitat with relatively short grass and little woody vegetation but can occur in dryland wheat.	High. Suitable nesting and foraging habitat within the analysis area. Not detected during the 2024 surveys but detected during avian use surveys for the BCWF.	Forage: Grasslands, agricultural areas. Nesting: Grasslands, agricultural areas.

Common Name	Scientific Name	State Status ¹ (Columbia Plateau Ecoregion)	Expected Habitats	Observed or Expected Occurrence within Analysis Area ²	Potential Use of Habitat Within the Analysis Area
Sagebrush sparrow	<i>Artemisiospiza nevadensis</i>	SC, SGCN	Sagebrush obligates. Found in shrub-steppe habitat, particularly in big sagebrush communities. They require high shrub cover and low grass and litter cover in relatively large patches.	Low. No suitable habitat for this species exists within the analysis area, although eBird sightings have been documented within the vicinity of the analysis area. May pass through the analysis area during migration.	Forage: All terrestrial habitats. Nesting: None.
Short-eared owl	<i>Asio flammeus flammeus</i>	SGCN	Large expanses of marshes and wet prairies for foraging and nesting including wetlands, grasslands, shrub-steppe, and agricultural lands. During winter, they utilize open areas, including pastures, hay meadows, marshes, and tidal areas, and may sometimes be found in trees or dense shrubs.	Low. Suitable winter habitat exists within the analysis area, but eBird relative abundance data indicates this species does not often occur within the vicinity of the analysis area. Observed incidentally during avian use surveys at BCWF.	Forage: All habitat types. Nesting: All habitat types.
Swainson's hawk	<i>Buteo swainsoni</i>	S, SGCN	Open-country specialists that hunt and forage in grassland, shrubsteppe, and agricultural areas. Nests are frequently in lone trees or isolated shrubs in open country.	High. Suitable nesting and foraging habitat within the analysis area. Not detected during the 2024 surveys but detected during avian use surveys for the BCWF and active Swainson's hawk nests detected during raptor nest surveys for the BCWF.	Forage: All habitat types. Nesting: All habitat types.
Wilson's phalarope	<i>Phalaropus tricolor</i>	PS, SGCN	Breeding habitat occurs in shallow freshwater and saline ponds, marshes, and wet meadows. Non-breeding habitat occurs on lake shores, mudflats, salt marshes, freshwater marshes, alkaline ponds, and sewage ponds.	Low. No suitable breeding or non-breeding habitat occurs within the analysis area. This species is a migrant and may only pass through the analysis area. Not detected at BCWF during any survey.	Forage: Freshwater Emergent Wetlands. Nesting: None.
Reptiles					
Northern sagebrush lizard	<i>Sceloporus graciosus graciosus</i>	S	Closely associated with steppe habitat with sandy soils and sparse vegetation in the grass/forb layer. Also occurs in chaparral, juniper woodlands, and coniferous forests.	Low. Recorded in Sherman County but limited suitable habitat exists within the analysis area. Not detected during the 2024 surveys, nor during pre- or post-construction surveys at the BCWF.	Forage: Terrestrial habitats where soils are sandy and cover is available. Breeding: None.
Western painted turtle	<i>Chrysemys picta bellii</i>	SC, SGCN	Marshy ponds, small lakes, slow-moving streams, and quiet, off-channel portions of rivers. Prefer waters with muddy bottoms and aquatic vegetation.	Low. Recorded in Sherman County but limited suitable habitat within the analysis area. Not detected during the 2024 surveys, nor during pre- or post-construction surveys at the BCWF.	Forage: Freshwater Emergent Wetlands. Nesting: None.
Amphibians					
Western toad	<i>Anaxyrus boreas</i>	S, SGCN	Breeds in wetlands, lakes, ponds, and streams and side channels with slow moving water. Adults occur in habitat with open forest canopies or landscapes with few to no trees, south-facing slopes, close proximity to water, and high density of cover (e.g., logs, rocks, burrows).	Moderate. Recorded in Sherman County but limited suitable habitat exists within the analysis area. Observed incidentally during pre- and post-construction surveys at BCWF.	Forage: All habitat types where cover is available. Breeding: Aquatic habitats with water present.
Invertebrates					
Monarch butterfly	<i>Danaus plexippus</i>	SGCN	Primary breeding habitat occurs where milkweed (<i>Asclepias</i> spp.) grows, which includes grasslands, scrublands, croplands, roadsides, urban and suburban parks and gardens, and sand dune areas.	Moderate, during migration. Narrowleaf milkweed (<i>Asclepias fascicularis</i>) was recorded during surveys conducted for RFA 4 and in the BCWF Wind HMA. Though these observations were not mapped, this plant prefers moist soils and is primarily found along the Columbia River Gorge and its associated drainages outside of the Analysis Area for this RFA 5. Monarch butterfly would likely only pass through the analysis area during migration.	Forage: All habitats except for agricultural areas. Breeding: Where milkweed may occur, such as drainages, streams, and residential areas.

Common Name	Scientific Name	State Status ¹ (Columbia Plateau Ecoregion)	Expected Habitats	Observed or Expected Occurrence within Analysis Area ²	Potential Use of Habitat Within the Analysis Area
Morrison bumblebee	<i>Bombus morrisoni</i>	SGCN	Open dry scrub, shrubland, grassland, and dry coniferous forests. Assumed to nest underground (e.g., in abandoned rodent burrows).	Low. This species was not detected within the vicinity of the analysis area during PNW Bumble Bee Atlas surveys. Suitable habitat for this species does not occur within the analysis area or its vicinity.	Forage: All terrestrial habitat types. Nesting: All terrestrial habitat types.
Western bumblebee	<i>Bombus occidentalis</i>	SGCN	Uses a wide variety of natural, agricultural, urban, and rural habitat types.	Low. This species was not detected within the vicinity of the analysis area during PNW Bumble Bee Atlas surveys. Habitat within the vicinity of the analysis area is low quality for this species.	Forage: All terrestrial habitat types. Nesting: All terrestrial habitat types.
<p>1. Oregon Status: S = State Sensitive; SC = State Sensitive—Critical; SGCN = Species of Greatest Conservation Need; PS = Pending Sensitive Status.</p> <p>2. The occurrence locations provided by ORBIC (2024) are from a variety of sources that vary in accuracy and precision. Because some ORBIC records, particularly older records, have a high level of uncertainty, some occurrence locations are represented by large polygons in order to reflect this lack of precision. These polygons are sometimes more than one mile in diameter around a single occurrence location, and do not necessarily reflect the actual location of the sensitive species. As a result, the ORBIC occurrence information was used as a rough approximation of occurrence.</p>					

4.2 ODFW Site-Specific Issues of Concern

The only site-specific issue for the Solar Components identified by ODFW was to avoid residual grassland habitats where possible, to leave cover for the sensitive birds in the area (J.L. Thompson, personal communication, June 6, 2024). Impacts to grassland habitats have been minimized through micrositeing. Approximately 1.1 acres of Category 3 Grassland-steppe habitat and 17.1 acres of Category 4 grassland-steppe habitat were identified within the analysis area (Table G-4). Construction of the Solar Components is expected to result in 0.9 acre of permanent disturbance to Category 4 grassland-steppe habitat (Table G-6). No disturbance is anticipated for Category 3 grassland-steppe habitat.

5.0 Baseline Survey of Habitat Use by State Sensitive Species – OAR 345-022-0060(3)(e)

(e) A baseline survey of the use of habitat in the analysis area by species identified in (d) performed according to a protocol approved by the Department and ODFW;

The desktop analysis described in Section 2.1 was performed to determine which sensitive species may be using the habitats available within the analysis area for breeding, foraging, or stop-over habitat during migration. Baseline field surveys, including habitat categorization and special status wildlife surveys, botanical surveys, raptor nest surveys, and wetland and waters surveys, were completed in 2024 within portions of the analysis area. In addition to the baseline surveys, a pre-construction raptor nest survey in support of the Solar Components approved in Amendment 4 of the Site Certificate was conducted in 2026, overlapping with the northern portion of the analysis area. Biologists observed two sensitive wildlife species, the loggerhead shrike and North American porcupine, during the 2024 surveys, though porcupine was not considered sensitive or pending sensitive status at the time the surveys were completed. The results of the surveys are described below. Additional details on the field survey methods and results are provided in Attachment G-1. The use of habitats within the analysis area by state sensitive species is summarized in Table G-5.

5.1 Results of Field Surveys

5.1.1 2024 Habitat Categorization and Special Status Wildlife Surveys

Habitat types were identified within the Habitat Categorization Survey Area, which covered 4,636 acres including the entire RFA 4 Site Boundary, RFA 5 Site Boundary, and some additional lands that have since been omitted. All habitat types were then categorized according to ODFW's Fish and Wildlife Habitat Mitigation Policy Habitat Categorization flowchart (ODFW 2008). The most prominent habitat category in the Habitat Categorization Survey Area was Category 6, Agriculture habitat type, which included both Irrigated and Non-irrigated Croplands, and Developed habitat subtypes. Additional habitat types included Category 4 Grasslands habitat subtype, Category 3 Sagebrush Shrub-steppe and Upland Trees habitat subtypes, as well as

Category 2 Intermittent and Riparian Tree habitat subtype (Attachment G-1). The RFA 5 Analysis Area does not contain Shrub-steppe or Riparian habitat. The survey results did not consider overlap with PWCAs and big game winter range layers, though no winter range or PWCAs exist within the RFA 5 Site Boundary or the analysis area.

Eighteen bird species, five mammal species, and two invertebrate species were observed during the surveys. Two special status species, loggerhead shrike and North American porcupine, were documented during surveys, though the porcupine was not observed within the analysis area (Figure G-5; Attachment G-1). North American porcupine was not considered sensitive or pending sensitive status at the time the surveys were completed, and thus was not discussed in detail in the report.

5.1.2 2024 Botanical Surveys

Botanical surveys were conducted to document the presence of federally or state listed endangered, threatened, and candidate vascular plant species and state and county-designated noxious weeds within the Botanical Survey Area, which covered 3,980 acres including the entire RFA 4 Site Boundary, RFA 5 Site Boundary, and some additional lands that have since been omitted from this RFA 5 (Attachment G-1). The botanist observed that vegetation within much of the Botanical Survey Area has been modified due to historic and current agricultural activity, construction and management of the Facility, and historic and current grazing activity. No threatened, endangered, proposed, or candidate vascular plant species were observed within the Botanical Survey Area. In addition, no suitable habitat for the candidate species Henderson's ricegrass (*Eriocoma [Achnatherum] hendersonii*), hepatic monkeyflower (*Erythranthe jungermannioides*), and sessile mouseltail (*Myosurus sessilis*) was observed within the Botanical Survey Area. Limited suitable habitat for the state threatened Lawrence's milkvetch (*Astragalus collinus* var. *laurentii*) was observed within the Botanical Survey Area, but historic and current anthropogenic activities have resulted in degradation of the limited suitable habitat for Lawrence's milkvetch within the Botanical Survey Area. Limited suitable habitat for Lawrence's milkvetch occurs within the RFA 5 Site Boundary. Twelve noxious weed species were identified within the Survey Area. Of the noxious weed species identified, 10 were documented within the analysis area (Attachment G-1).

The results of the botanical surveys were reviewed against the 2024 ODA noxious weed list (ODA 2024) and the OregonFlora collection database (OregonFlora 2026) in support of this RFA 5 Exhibit G and Exhibit H. There are no noxious weeds in the 2024 ODA noxious weed list that were not already documented during the botanical survey. In addition, according to the OregonFlora database, there are now recorded occurrences of spreading stickseed (*Hackelia diffusa* var. *diffusa*; state candidate) within Sherman County (OregonFlora 2026). This species was not a target species in the 2024 botanical survey report; however, based on the results of the report, there is no suitable habitat for this species within the analysis area. Spreading stickseed is discussed further in Exhibit H.

5.1.3 Raptor Nest Surveys

No state sensitive species were observed nesting within the Raptor Nest Survey Area during the 2024 raptor nest surveys or the 2026 pre-construction raptor nest surveys conducted for the approved solar generation facility described in RFA 4 (Attachment G-1). A total of five nests were observed within the 2024 Raptor Nest Survey Area, including one in-use great horned owl (*Bubo virginianus*) nest, one inactive great horned owl nest, and three inactive nests with unknown species determinations. Two active red-tailed hawk nests (*Buteo jamaicensis*) and two inactive nests with unknown species determinations were observed in the 2026 pre-construction Raptor Nest Survey Area, all of which were observed outside of the analysis area. The nearest nest to the RFA 5 Site Boundary was an active red-tailed hawk nest approximately 0.75 miles northwest of the substation (Attachment G-1, 2026 raptor nest survey report provided under separate cover).

5.1.4 Wetlands and Waters Surveys

The 2024 wetland and waters surveys covered the entirety of the RFA 4 Site Boundary, RFA 5 Site Boundary, and some additional lands that have since been omitted (Exhibit O, Attachment O1-1). A total of two palustrine emergent wetlands, one riverine wetland, three ephemeral waterways, and one desktop delineated intermittent waterway were found within the Wetland and Waters Survey Area. Of these results, only the two palustrine emergent wetlands are within the analysis area, though outside of the RFA 5 Site Boundary (Figure G-4).

6.0 Description of Potential Adverse Impacts – OAR 345-022-0060(3)(f)

(f) A description of the nature, extent and duration of potential adverse impacts on the habitat identified in (b) and species identified in (d) that could result from construction, operation and retirement of the proposed facility;

This section describes potential impacts to habitat and state sensitive species that are known to occur or have the potential to occur within the analysis area. Permanent impacts to fish and wildlife habitat are discussed as well as disturbance impacts to mammals, birds, reptiles, amphibians, and invertebrates. As described in detail in Section 3.1.5 of the RFA 5 Background document, the Certificate Holder proposes to construct the Solar Components over a period of 18 months.

Construction and operation of the Solar Components will result in both permanent and temporary impacts to wildlife and their habitats (see Table G-6 in Section 6.1). Due to the multi-year construction schedule of the Solar Components, both permanent and temporary impacts to fish and wildlife habitat will occur in phases over this time period. Permanent impact areas are those that will be converted from the existing condition to a different condition for the life of the Solar Components.

Direct impacts to habitat include permanent loss of some specific habitat types; indirect impacts may include increased potential for the invasion of noxious weeds, particularly along fence lines and roads. These habitats are identified and described in Section 3.0, and Table G-6 in Section 6.1 provides the number of acres that will be permanently and temporarily impacted by the Solar Components, organized by habitat type and sub-type. The entire area within the fenceline is considered permanently disturbed; all temporary disturbance areas are outside the fenceline. Permanent disturbance will also result from support poles associated with generation tie connections from the proposed collector substation to the existing Biglow Canyon Substation, BESS, O&M structures, the substation, and a portion of the substation access road. These components are described in detail in Section 3.0 of the RFA 5 Background document.

Construction of the Solar Components is expected to involve up to 44.1 acres of temporary impacts (Table G-6). The specific extent of each component's temporary impact is detailed in Section 3.4.3 of the RFA 5 Background document and described in terms of a total, worst-case scenario impact for the full duration of construction. The RFA 5 Background document presents temporary and permanent impacts of the Solar Components. Restoration of temporary impact areas will occur following construction per the Comprehensive Solar Revegetation and Soil Management Plan (Exhibit C, Attachment C-2).

6.1 Potential Impacts to Fish and Wildlife

Potential impacts on fish and wildlife habitat include the temporary and permanent disturbance to habitat associated with the construction and operation of the Solar Components. The acres of temporary and permanent disturbances are summarized in Table G-6. Some impacts will be avoided or minimized as described in Section 7.0. Impacts that cannot be avoided will be mitigated as described in the Biglow Solar Habitat Mitigation Plan (HMP; Attachment G-2). Impacts to state sensitive species are described in Section 6.2. No PWCAs or big game winter range are present within the analysis area; therefore, the construction and operation of the Solar Components will not adversely affect habitat within PWCAs or big game winter range.

Table G-6. Disturbance by Habitat Category and Land Cover Type

ODFW Habitat Category	Land Cover Type	Temporary Disturbance (acres)	Permanent Disturbance (acres)
3	Upland – Upland Trees	-	5.0
Category 3 Total		-	5.0
4	Grassland-steppe – Grasslands	-	0.9
Category 4 Total		-	0.9
6	Agriculture – Non-irrigated Cropland	43.9	1,237.1
	Developed	0.2	1.1
Category 6 Total		44.1	1,238.2
Grand Total		44.1	1,244.1
Note: Totals in this table may not sum correctly due to rounding. "-" = no impact; <0.1 = greater than zero but less than 0.05 acre.			

6.2 Potential Impacts to State Sensitive Species

This section addresses potential adverse impacts to state sensitive species identified in Section 4.0. Habitat loss and modification resulting from construction activities will occur in permanent impact areas and temporary construction areas, and the associated impacts on wildlife will vary by species. In addition to these habitat-related impacts, potential adverse impacts to sensitive species due to construction and operation activities may include potential nesting and breeding disturbance, collision with solar arrays, security perimeter fencing, overhead collector lines, vehicles and heavy equipment, and disturbance related to artificial lighting.

6.2.1 Mammals

6.2.1.1 Bats

Twelve state sensitive and/or SGCN bat species have the potential to occur within the analysis area: California myotis (SGCN), canyon bat (pending sensitive status, SGCN), hoary bat (sensitive, SGCN), little brown myotis (pending sensitive status, SGCN), long-legged myotis (SGCN), pallid bat (sensitive, SGCN), silver-haired bat (sensitive, SGCN), spotted bat (sensitive, SGCN), Townsend's big-eared bat (sensitive-critical, SGCN), western long-eared myotis (pending sensitive status, SGCN), western small-footed myotis (pending sensitive status, SGCN), and Yuma myotis (pending sensitive status, SGCN). Two state sensitive species (hoary bat and silver-haired bat) were detected during fatality monitoring at the BCWF (Table G-5). The timing and frequency of detections suggest that these species are relatively common and fly through much of the area during the late summer and fall months, concurrent with their migration period. The 10 additional state-sensitive and/or SGCN bat species with the potential to occur in the analysis area were not detected during fatality surveys at BCWF. The Facility is not sited near typical breeding or roosting habitat for these species, and there is limited aquatic foraging habitat such as wetland, ponds, and slow-moving streams within the site boundary (Table G-4, Figure G-4). Additionally, construction activities will typically occur during daylight hours when bats are generally absent, and thus construction activities are not anticipated to disturb foraging bats.

Collision with solar panels and security fencing is a potential threat to bats. Post-construction bat mortality data at utility-scale PV solar energy sites are limited. Smallwood (2022) reviewed 18 California fatality monitoring reports from 1982 to 2018 and calculated an average fatality estimate of 0.06 bats/MW/year at PV solar arrays, 2.56 bats/kilometer/year along perimeter fences, and zero fatalities associated with gen-tie power lines. Any impacts to bats would most likely occur in late summer and fall, during the migratory period for tree-roosting bats. Insects may be attracted to lighting around structures, which may in turn attract bats to forage near Solar Components infrastructure. Thus, artificial lighting at night may increase the risk of collision fatalities. However, the potential for collision risk due to artificial night lights will be avoided and minimized, as described in Section 7.0. As a result, construction and operation of the Solar Components are anticipated to have minimal impact on these bat species, including during their migratory period.

6.2.1.2 Other Mammals

The two remaining state sensitive and/or SGCN mammal species that may be impacted by the Solar Components are North American porcupine (pending sensitive status, SGCN) and white-tailed jackrabbit (SGCN). North American porcupine was observed during the 2024 surveys (Attachment G-1). White-tailed jackrabbit was observed during multiple surveys that were conducted in support of the application for the Existing Facility (Orion Sherman County Wind Farm 2005). Both of these species have been observed incidentally during post-construction surveys and monitoring for BCWF (WEST 2009, 2010, 2012a, 2012b, 2012c, 2013). Potential impacts to these species and other terrestrial mammals include habitat loss, barriers to dispersal, and direct mortality. Both of these species may occur in all of the habitats mapped within the RFA 5 Site Boundary and studies indicate that both North American porcupine and white-tailed jackrabbit may rely on cultivated wheat and barley when native food sources are scarce (Johnson 1977; Berteaux et al. 2005; Bear and Hansen 1966; Flinders and Hansen 1971, 1972; Brunton 1981; USFWS 2017). However, an abundance of similar agricultural lands in the vicinity of the RFA 5 Site Boundary remain available to provide forage habitat for North American porcupine and white-tailed jackrabbit individuals that may be displaced. In addition, the implementation of speed limits within the RFA 5 Site Boundary are expected to reduce direct mortality impacts to terrestrial mammals. As a result, impacts from the construction and operation of the Solar Components are expected to be minimal for these species and other terrestrial mammals.

6.2.2 Birds

Fourteen state sensitive and/or SGCN bird species and one protected eagle species have the potential to occur within the analysis area and could be potentially impacted by the Solar Components (Table G-5). Other, non-sensitive bird species potentially occurring within the RFA 5 Site Boundary could also be impacted by the Solar Components. Direct habitat loss will occur from the development of the RFA 5 Site Boundary, and habitat fragmentation may reduce the functionality of this area for birds. However, most of the Solar Components will be located in disturbed habitat and an abundance of similar agricultural lands in the vicinity of the RFA 5 Site Boundary remain available to provide habitat for avian individuals potentially displaced.

Temporary impacts from noise and visual disturbance associated with the presence of personnel, vehicles, and equipment during construction could disrupt normal feeding or breeding activities or cause birds to avoid the area. Operations will have reduced levels of human activity compared to construction but could still cause birds to alter their typical behavior and/or avoid the area.

Unlike avian fatality data available for wind turbines, there have been limited standard fatality data associated with PV solar energy infrastructure collected and available across the country. The available data on avian mortality at utility-scale solar energy sites suggests mortality at PV facilities is low, compared to collisions occurring with other anthropogenetic infrastructure (e.g., wind turbines, communication towers, tall buildings). In a study by Kosciuch et al. (2020) that analyzed fatality monitoring data from 10 PV solar facilities in California and Nevada, a fatality estimate of

2.5 birds/MW/year was calculated, but this was reduced to an average annual fatality rate of 1.8 birds per MW per year when an outlier project was removed.

It is theorized that waterbirds confuse the reflective array of solar panels located near ground level to be a body of water and attempt to land on them, and other bird species may fly into the reflective panels like they would windows, unable to perceive the difference between the sky and the glass. The Kosciuch et al. (2020) study showed that water obligate bird fatalities were present in most of the studies (80 percent). Water obligates are species that depend on water for mobility and cannot take flight from land, while water associates are species that rely on water for foraging, reproduction, and/or roosting. Kosciuch et al. (2020) stated that the causal mechanism responsible for water obligate mortalities at PV facilities could not be concluded with their study, and their results were not predictive outside of the vicinity of the sites included in their study.

In a recent solar study that included six small solar arrays located next to Lake Michigan, no water obligate mortalities were recorded and only two water associate mortalities showed evidence of death caused by colliding with solar panels (Rodriguez et al. 2023). In addition, an analysis that summarized bird fatalities to date at PV solar facilities in Alberta, Canada, was conducted and found no correlation with water associate and water obligate mortalities being present at higher numbers than other avian species (Kosciuch et al. 2022).

The results from three publicly available studies from large-scale California PV facilities reported the avian taxonomic groups most highly represented as fatalities were passerines (49 percent), followed by doves/pigeons (22 percent), although loons and grebes ranked second at one facility (23 percent; WEST 2014). Passerines are “perching birds” that make up almost half of all bird species, so that high percentage could easily correlate with the high number of birds in that taxonomic group. When analyzing the fatality species composition, often the species with the largest populations spend most of their time close to the ground and inhabit landscapes with relatively low-growing vegetation or are habituated to human structures. These species often represent the highest number of fatalities at PV facilities (e.g., horned larks [*Eremophila alpestris*], western meadowlarks [*Sturnella neglecta*], mourning doves [*Zenaida macroura*], house sparrows [*Passer domesticus*], house finches [*Haemorhous mexicanus*], gray partridge [*Perdix perdix*]; Kosciuch et al. 2020, Kosciuch et al. 2022, WEST 2014).

In Oregon, results of a fatality study at a 56-MW PV facility near Prineville detected only three bird fatalities during one year of standardized searches, including only two native birds: horned lark and a dark-eyed junco (*Junco hyemalis*) (ODOE 2020). These findings, which are the first for the region in Oregon, imply that significant fatality events are unlikely at PV solar facilities in the area. However, low numbers of common ground-dwelling bird species fatalities are possible (ODOE 2020).

Although the fatality numbers from solar studies suggest the probability for waterbird fatality events associated with hitting solar panels is low, waterbirds may attempt to land on the solar panels at the Solar Components, which is located in the vicinity of the Columbia River, a known waterbird stop-over area, especially in times of poor weather and low visibility, and during nocturnal migration.

Artificial lighting has been shown to attract nocturnally migrating birds, resulting in collisions with structures such as communication towers, guy wires, and wind turbines. Research has shown that lighting at solar energy facilities can be designed to minimize that attraction, including using lights that are directed downward, have the lowest allowable intensity, and are on timers or motion sensors.

Interactions with overhead electrical systems have the potential to negatively affect birds, especially raptors, primarily through electrocutions and collisions. Electrocution on overhead electric systems is a primary cause of anthropogenic mortality for golden eagles in North America (Dwyer et al. 2020). Powerline collisions appear to be a less frequent source of mortality for raptors compared to electrocutions (Loss et al. 2014). Collisions with powerlines are possible; however, transmission lines will be constructed following the Avian Powerline Interaction Committee (APLIC) recommendations for collision avoidance (APLIC 2012). Powerline electrocution is not expected at the Solar Components because all transmission lines will be constructed following the latest APLIC design standards (APLIC 2012), as specified in Site Certificate Condition GEN-FW-01.

The security perimeter fencing proposed for the Solar Components, estimated to total up to approximately 8.5 miles of fence, is an additional collision hazard for birds. The security perimeter fencing will be chain-link and up to 8 feet in height with additional strands of barbed or razor wire on top.

During the 2024 survey effort and the 2026 pre-construction survey, no nests of sensitive raptor species were observed in the analysis area. The only sensitive raptor species observed nesting during long-term raptor nesting surveys at the BCWF was the Swainson's hawk (state sensitive). Two great horned owl nests and three inactive nests were recorded during the 2024 raptor nest surveys, and two red-tailed hawk nests and two inactive nests were recorded during the 2026 pre-construction survey (Attachment G-1). Most of the nests were associated with Emigrant Canyon, east of the analysis area. Construction activities occurring near nests during the sensitive breeding season could cause the nests to fail. The protective buffers and survey requirements identified in Site Certificate Condition CON-FW-01 will be implemented for raptor nests that are determined to be active during construction.

Construction occurring during the breeding season (as early as February and lasting through August) could cause active nests to fail due to direct destruction of nests or nearby disturbance causing nest abandonment. Active nests of all native birds are protected under the Migratory Bird Treaty Act. As described above, 14 state sensitive bird species and 1 eagle species may occur within the analysis area. Impacts to sensitive bird species with the potential to occur within the RFA 5 Site Boundary are addressed below. Measures described in Section 7.0 will be used to minimize or avoid these potential impacts.

- **American avocet (pending sensitive status, SGCN):** No American avocet were recorded during 2024 surveys (Attachment G-1), nor were any detected during avian use surveys for the BCWF. This species uses alkali and freshwater ponds, lakes, and wetlands with extensive shallow areas and prefer sites with minimal disturbance (ODFW 2026a). No suitable habitat for American avocet exists within the RFA 5 Site Boundary, but this species

may stop at the Freshwater Emergent Wetland within the analysis area during migration. Potential adverse impacts from construction and operation of the Solar Components include potential collision with the solar arrays, perimeter fencing, overhead collector line, and vehicles during the spring and summer months. American avocet fatalities have been documented at PV solar facilities, though uncommon, and this species is a nocturnal migrant (Kagan et al. 2014); therefore, this species may be attracted to artificial lights produced by the Solar Components during migration, creating additional collision risks.

- **Bald eagle (no state status, BGEPA-protected):** No bald eagle nests or bald eagle individuals were recorded during 2024 and 2026 surveys (Attachment G-1). Bald eagles have been observed during avian use surveys for the BCWF. There is no suitable nesting habitat for the bald eagle within the analysis area. Bald eagle use of the Solar Components is expected to be mostly limited to migration and winter, when the species sometimes feeds on carrion (Buehler 2022). Although bald eagles may occasionally pass through the area, construction and operation of the Solar Components are not expected to adversely impact bald eagles.
- **Black tern (pending sensitive status, SGCN):** No black terns were recorded during 2024 surveys (Attachment G-1) nor were any detected during avian use surveys for the BCWF. This species relies on shallow, freshwater marshes, flooded meadows, and wetlands with emergent vegetation (ODFW 2026a). No suitable habitat for this species occurs within the RFA 5 Site Boundary and the Freshwater Emergent Wetland habitat within the analysis area is poor quality habitat for this species; however, black tern may pass through the Analysis Area during migration. However, while this species has not been recorded as a fatality at PV solar facilities, other tern species have been recorded, although rarely (Kosciuch et al. 2020). Potential adverse impacts from construction and operation of the Solar Components include potential collision with the solar arrays, perimeter fencing, overhead collector line, and vehicles during the spring and summer months. This species is known to be both a diurnal and a nocturnal migrant (Heath et al. 2020); therefore, this species may be attracted to artificial lights produced by the Solar Components during migration, creating additional collision risks.
- **Brewer's sparrow (sensitive, SGCN):** No Brewer's sparrows were recorded during 2024 surveys (Attachment G-1). Additionally, no Brewer's sparrows were detected during avian use surveys for the BCWF; a single individual was found during fatality monitoring for the BCWF (Table G-5). This species uses shrublands, generally with a canopy height of more than 5 feet. Despite the lack of detections, Brewer's sparrows could potentially occur in the limited shrub-steppe habitat within the RFA 4 Site Boundary, just northeast of the analysis area, and could therefore move through the RFA 5 Site Boundary during migration. Construction and operation of the Solar Components may result in potential nesting disturbance in the northeastern portion of the RFA 5 Site Boundary and potential collision risks with vehicles, solar arrays, perimeter fencing, and overhead electrical lines. Sparrow fatalities have been documented at PV solar facilities (WEST 2014), although no Brewer's sparrows have been reported specifically. Brewer's sparrows are a nocturnal migrant (ADW

2024); therefore, this species may be attracted to artificial lights produced by the Solar Components during migration, increasing collision risks.

- **Burrowing owl (sensitive-critical, SGCN):** No burrowing owls were recorded during 2024 and 2026 surveys (Attachment G-1), but the species was detected during pre-construction surveys at the BCWF (Table G-5). The species inhabits wide-open, sparsely vegetated areas and could occur in grassland, shrubland, agricultural, and developed habitats within the RFA 5 Site Boundary, especially if burrowing mammals are present. Potential adverse impacts from the construction and operation of the Solar Components include the loss of foraging and breeding habitat, potential collapse of burrows and nesting disturbance, and potential collision risks with vehicles, solar arrays, perimeter fencing, and overhead collector lines during the breeding season. Burrowing owl fatalities have been documented at PV solar facilities (WEST 2014). This species is a nocturnal migrant and, therefore, may be attracted to artificial lights produced by the Solar Components during migration, increasing collision risks.
- **Common nighthawk (sensitive, SGCN):** No common nighthawks were recorded during the 2024 surveys (Attachment G-1), but one individual was found during fatality monitoring for the BCWF (Table G-5). Common nighthawk is most commonly found in open or semi-open terrain. The species tends to roost and nest on gravel or sparsely vegetated grasslands (Brigham et al. 2020). It is an aerial insectivore that feeds in low-light conditions at dusk and dawn, often near water, but also on insects attracted to artificial lights (Brigham et al. 2020). Potential impacts to common nighthawk during construction and operation of the Solar Components include roosting and nesting disturbance and collision with vehicles, solar arrays, perimeter fencing, and overhead collector lines during the breeding season and collision with vehicles and turbine blades. Nighthawks have been documented at PV solar facilities (WEST 2014). Common nighthawk is a crepuscular insectivore. Though there is still uncertainty regarding the relationship between this species and artificial lighting, studies indicate that extra-territorial individuals are more likely to hunt insect prey under artificial lights in non-urban areas, while individuals performing territorial activities (e.g., nesting) tend to avoid artificial light (Adams et al. 2024, Adams et al. 2025). Therefore, this species may be attracted to the artificial lights produced by the Solar Components, increasing collision risks.
- **Ferruginous hawk (sensitive-critical, SGCN):** No ferruginous hawks were recorded during 2024 and 2026 surveys (Attachment G-1). Low numbers of ferruginous hawks have been detected during avian use surveys for the BCWF but no nests have been found during raptor nest surveys for the BCWF (Table G-5). The potential adverse impact to this species includes loss of foraging habitat, as no nests were observed within the analysis area. Limited hawk fatalities have been documented at PV solar facilities, and no ferruginous hawk fatalities specifically; therefore, the solar arrays should not create a collision risk to this species.

- **Golden eagle (pending sensitive status, SGCN, BGEPA):** No golden eagles were recorded during 2024 and 2026 surveys (Attachment G-1). Golden eagles were regularly detected during avian use surveys for the BCWF but no nests were detected within the survey area during raptor nest surveys for the BCWF (Table G-5). The potential adverse impact to this species includes loss of foraging habitat and vehicular collisions. No golden eagle fatalities have been recorded at solar fatalities, and the species is assumed not to be at risk of colliding with solar arrays. The implementation of speed limits will likely reduce the risk of foraging golden eagles from being struck by vehicles and equipment.
- **Grasshopper sparrow (sensitive, SGCN):** No grasshopper sparrows were recorded during the 2024 surveys (Attachment G-1), but grasshopper sparrows were regularly recorded during avian use surveys for the BCWF (Table G-5). This species uses dry grassland habitat, generally with low to moderate grass height and low percent shrub cover. It could potentially occur in the limited grassland habitat within the RFA 5 Site Boundary. Impacts to appropriate habitat for this species during the construction and operation of the Solar Components include permanent impact to 0.9 acre of Grassland habitat and no temporary impacts (Table G-6). Other potential adverse impacts from collision risks exist with vehicles, heavy equipment, solar arrays, perimeter fencing, and overhead collector line. Sparrow fatalities have been documented at PV solar facilities (WEST 2014). Grasshopper sparrow is also a nocturnal migrant (Hill and Renfrew 2018); therefore, this species may be attracted to artificial lights produced by the Solar Components during migration, causing additional collision risks.
- **Lewis's woodpecker (sensitive-critical, SGCN):** No Lewis's woodpeckers were recorded during 2024 surveys (Attachment G-1) or during pre- or post-construction surveys at the BCWF (Table G-5). Lewis's woodpecker is associated with open forests, often at lower elevations, and nest in woodlands of the river valleys of eastern Oregon (Csuti et al. 2001). The RFA 5 Site Boundary does not provide breeding habitat, but the species has the potential to pass through the Solar Components during migration. As a diurnal migrant (Abele et al. 2004), this species will not be adversely impacted by artificial lighting. This species has also not been documented as a fatality at solar fatalities. Impacts to Lewis's woodpeckers from construction and operation of the Solar Components are anticipated to be minimal.
- **Loggerhead shrike (sensitive, SGCN):** One loggerhead shrike was recorded during 2024 surveys (Attachment G-1). The shrike was observed in Upland Trees habitat in the southern portion of the RFA 5 Site Boundary. Additionally, low numbers (one to two individuals) of loggerhead shrikes have been occasionally detected during avian use surveys for the BCWF. This species uses tall sagebrush for nesting and roosting, and forages in open areas with grasses and bare ground (ODFW 2026a). Potential adverse impacts to loggerhead shrike include habitat loss, nesting disturbance, and collisions associated with vehicles, solar arrays, perimeter fencing, and overhead collector lines. Loggerhead shrike fatalities have been documented at PV solar facilities, and this species is a nocturnal migrant

(WEST 2014); therefore, this species may be attracted to artificial lights produced by the Solar Components during migration, increasing collision risks.

- **Long-billed curlew (sensitive-critical, SGCN):** No long-billed curlews were recorded during the 2024 surveys (Attachment G-1), but the species has been detected during avian use surveys for the BCWF (Table G-5). Long-billed curlew is typically found in open grasslands and could occur in grassland or agricultural habitats within the RFA 5 Site Boundary. Potential adverse impacts from construction and operation of the Solar Components include loss of potential nesting and foraging habitat as well as potential collision with the solar arrays, perimeter fencing, overhead collector line, and vehicles during the spring and summer months. Long-billed curlew fatalities have been documented at PV solar facilities and this species is a nocturnal migrant (WEST 2014); therefore, this species may be attracted to artificial lights produced by the Solar Components during migration, creating additional collision risks. Additionally, long-billed curlew is susceptible to human disturbance during the breeding season, which can result in nest abandonment or disruption of brood-rearing (Dugger and Dugger 2002); construction of the Solar Components may adversely impact active breeding attempts if construction occurs in proximity to long-billed curlew during the breeding season.
- **Short-eared owl (SGCN):** No short-eared owl individuals or nests were recorded within the analysis area during the 2024 and 2026 surveys. However, short-eared owl was observed incidentally during avian use surveys for the BCWF (Table G-5). Short-eared owl requires large expanses of open country habitat, such as wetlands, grasslands, shrub-steppe, and agricultural lands (ODFW 2026a). During the winter, this species will also use areas with tree roosts (Wiggins et al. 2020). There are few documented fatalities of short-eared owl at PV solar facilities, though the causes of death are often undetermined (Walston et al. 2015; Conkling et al. 2023). Limited suitable nesting habitat exists within the RFA 5 Site Boundary, but this species is known to occur within the vicinity of the analysis area during the winter and spring (Fink et al. 2023). Construction of the Solar Components may result in loss of foraging habitat during the winter and migratory period, as well as increased potential of collision with the solar arrays, perimeter fencing, overhead collector line, and vehicles. Implementation of speed limits within the RFA 5 Site Boundary should minimize collision risk with vehicles.
- **Swainson's hawk (sensitive, SGCN):** No Swainson's hawk individuals or nests were recorded within the Analysis Area during the 2024 and 2026 surveys. However, Swainson's hawk has been detected during avian use surveys for the BCWF and has been a regularly occurring nesting raptor species at the BCWF (Table G-5). Swainson's hawk is an open-country specialist that forages extensively while in flight. Swainson's hawk is more likely to hunt in agricultural areas, such as hay and alfalfa fields, pastures, and grain crops than is most other raptor species (Bechard et al. 2020). Construction of the Solar Components may result in loss of foraging habitat and reduce nesting habitat if trees are removed. To prevent disturbance to nesting Swainson's hawk, the Certificate Holder will not conduct ground-

disturbing activities within 0.25 miles of active Swainson's hawk nests during the nesting season, as described in Section 7.0. Limited hawk fatalities have been documented at PV solar facilities, and no Swainson's hawk fatalities specifically (WEST 2014); therefore, the solar arrays should not create a collision risk to this species.

- **Wilson's phalarope (pending sensitive status, SGCN):** No Wilson's phalarope individuals were recorded within the analysis area during the 2024 surveys or during surveys at the BCWF (Table G-5). Wilson's phalarope breeds in wetlands and is strongly dependent on saline lakes for food (ODFW 2026a). This species is rare in Sherman County and suitable foraging or breeding habitat does not exist within the RFA 5 Site Boundary; however, Wilson's phalarope may occur as a migrant within the vicinity of the analysis area. Wilson's phalarope is not commonly documented as a fatality at PV solar generation facilities, but other shorebirds are (Conkling et al. 2023). Construction of the Solar Components may result in collision with solar arrays, perimeter fencing, overhead collector line, and vehicles. In addition, Wilson's phalarope is a nocturnal migrant; therefore, this species may be attracted to artificial lights produced by the Solar Components during migration, creating additional collision risks.

6.2.3 Reptiles and Amphibians

No state sensitive reptiles or amphibians were documented during the 2024 surveys (Attachment G-1); however, western toad (sensitive, SGCN) was observed during pre- and post-construction surveys at the BCWF. No state sensitive reptiles were observed during the pre- and post-construction surveys at the BCWF. In addition to western toad, two state sensitive reptile species have the potential to occur within the analysis area: northern sagebrush lizard (sensitive) and western painted turtle (sensitive-critical, SGCN; Table G-5). Potential impacts to these species and other reptiles and amphibians include habitat loss and direct mortality. There is no suitable habitat for the northern sagebrush lizard (shrub-steppe), western painted turtle (riparian, surface water, and wetland habitats), or western toad (surface water, wetland habitats) within the RFA 5 Site Boundary, though habitat exists for these species within the analysis area and its vicinity. Best management practices employed on-site in accordance with the Erosion and Sediment Control Plan (Exhibit C, Attachment C-1) are expected to result in minimal impacts to aquatic reptiles and amphibians such as the western painted turtle and western toad. Additional measures outlined in Section 7.0 will be used to further minimize impacts to aquatic reptiles and amphibians.

Use of machinery for clearing, grading, and trenching the Solar Components may cause some mortality of non-aquatic reptiles, such as northern sagebrush lizard. Impacts during operation are anticipated to be limited to direct mortality caused by vehicles and are expected to be low.

7.0 Avoidance and Mitigation Measures – OAR 345-022-0060(3)(g)

(g) A description of any measures proposed by the applicant to avoid, reduce, or mitigate the potential adverse impacts described in (f) in accordance with the general fish and wildlife habitat mitigation goals and standards described in OAR 635-415-0025 and a description of any measures proposed by the applicant to avoid, minimize, and provide compensatory mitigation for the potential adverse impacts described in (f) in accordance with the sage-grouse specific habitat mitigation requirements described in the Greater Sage-Grouse Conservation Strategy for Oregon at OAR 635-140-0000 through 635-140-0025, and a discussion of how the proposed measures would achieve those goals and requirements; and

This section describes measures to avoid, minimize, and mitigate for impacts to state sensitive and other wildlife species and their habitats, and describes how those measures are expected to achieve the habitat mitigation goals of OAR 635-415-0025. The RFA 5 Site Boundary does not occur in sage-grouse habitat; therefore, the mitigation requirements associated with the Greater Sage-Grouse Conservation Strategy for Oregon do not apply.

7.1 Project Design

Measures employed during design of the Solar Components to avoid and minimize impacts to fish and wildlife habitat, as well as state sensitive species, included the following:

- To the extent feasible, Solar Components have been sited on previously disturbed habitat, including cropland and developed areas, thereby minimizing impacts to other habitats more suitable for sensitive and other wildlife species, grasslands and wetlands. This siting effort is expected to minimize impacts to wildlife generally and to sensitive species in particular.
- Per Site Certificate Condition GEN-FW-01, the Certificate Holder will construct all overhead transmission lines following PGE's avian-safe design standards which are consistent with the suggested practices manual published by APLIC (APLIC 2012). This is expected to minimize the risk of electrocution and collision to all bird species, and to eagles, Swainson's hawks, and ferruginous hawks in particular. Per Site Certificate Condition GEN-LU-12, the Certificate Holder will implement permanent down-shield lighting at the O&M buildings, substation, and other Solar Components. Outdoor lighting will be sited, limited in intensity, shielded, and hooded in a manner that prevents the lighting from projecting onto adjacent properties and roadways. This is expected to minimize the risk of avian collision with Solar Components infrastructure for all birds and bats in general, but to nocturnal migrant species (including Brewer's sparrows and grasshopper sparrows) and to the crepuscular, insectivorous common nighthawk in particular. Down-shield lighting will be in place year-round, mitigating impacts to birds and bats both during migration and while foraging for insects at any time of the year.

7.2 Construction and Operation

During construction and operation of the Solar Components, measures for avoiding and reducing impacts to wildlife and plants, including state sensitive species, will be implemented as follows:

- Per Site Certificate Condition GEN-LU-12, the Certificate Holder will use the minimum lighting necessary for nighttime construction, repairs, or emergencies. The lighting will be directed downward to limit the illumination to the work area.
- Per Site Certificate Condition CON-SP-02, the Certificate Holder shall control the introduction and spread of noxious weeds in accordance with the methods, monitoring procedures and success criteria set forth in the Comprehensive Solar Revegetation and Soil Management Plan (Exhibit C, Attachment C-2).
- Per Site Certificate Condition CON-SP-02, the Certificate Holder shall restore areas that are temporarily disturbed during construction in accordance with the methods, monitoring procedures and success criteria set forth in the Comprehensive Solar Revegetation and Soil Management Plan (Exhibit C, Attachment C-2).
- Per Site Certificate Conditions PRE-FW-01 and OPR-FW-05, the Certificate Holder shall submit a final Habitat Mitigation Plan and implement it once operations of the Solar Components begin. The draft Biglow Solar HMP is included as Attachment G-2.
- Per Site Certificate Condition OPR-FW-01, the Certificate Holder shall conduct wildlife monitoring and mitigation in accordance with the Wildlife Monitoring and Mitigation Plan (WMMP) that was incorporated in the Final Order on Amendment 2. Components of the WMMP applicable to the Solar Components include ongoing environmental training for Solar Components personnel and reporting requirements governing incidental wildlife injuries and deaths observed at the Solar Components and raptor nest monitoring. Raptor nest monitoring performed for BCWF covers the entirety of the RFA 5 Site Boundary, and this RFA does not propose any modifications to the methods or timing of future raptor nest monitoring efforts associated with BCWF and the Solar Components.
- Per Site Certificate Condition CON-FW-01, the Certificate Holder will perform pre-construction raptor nest surveys to determine the location of active raptor nests. Survey methods shall be approved by ODFW. Condition CON-FW-01 includes nest buffers and seasonal restriction dates within which the Certificate Holder would not engage in high-impact construction activities (defined in Condition CON-FW-01).
- Per Site Certificate Conditions CON-SP-02, CON-SP-03, and CON-WM-01, implementation of the Comprehensive Solar Revegetation and Soil Management Plan, a Spill Prevention, Control, and Countermeasure (SPCC) Plan, and a Construction Waste Management Plan will ensure measures are in place to avoid and minimize environmental effects of spills and hazardous materials during construction activities. Per Site Certificate Conditions OPR-WM-02, OPR-SP-06, and OPR-SP-07, the Comprehensive Solar Revegetation and Soil

Management Plan, and Operations SPCC Plan, and an Operations Waste Management Plan will be implemented during operations.

- Per Site Certificate Condition PRE-WL-01, jurisdictional waters and their buffers will be flagged to prevent work activities from occurring within sensitive aquatic habitats within the RFA 5 Site Boundary.
- Per Site Certificate Conditions OPR-WM-01, OPR-SP-01, OPR-SP-02, and GEN-SP-02, operations of the Solar Components will avoid spills and discharge of sediment, sanitary wastewater, and hazardous materials into wildlife habitat.

During construction of the underground collector system, the Certificate Holder shall open the smallest necessary sections of trench during each day of construction and backfill the trenches as soon as is practical after power lines have been set in the trenches. If trenches are not filled by the end of the day, they shall be covered with a sturdy material (e.g., plywood or similar material) overnight. Doing so is expected to safeguard animals from falling into the trenches and becoming trapped.

7.3 Environmental Training and Sensitive Resource Awareness

Per Site Certificate Condition OPR-FW-01, the Certificate Holder will follow the WMMP, which includes providing ongoing environmental training to construction and operation staff. In addition, as prescribed in the Comprehensive Solar Revegetation and Soil Management Plan (Exhibit C, Attachment C-2), a Worker Environmental Awareness Program will be implemented, outlining to construction and operation staff how to avoid impacts to sensitive species and their habitat, review exclusion areas/seasonal avoidance buffers, and other environmental issues. Training includes reporting injured or dead wildlife discovered incidental to construction and maintenance operations in accordance with the procedures described in the WMMP (BCWF 2007).

7.4 Avian Protection

The Certificate Holder will implement the following avian protection measures:

- Condition CON-FW-01 – nest buffers and seasonal restrictions.
- Condition GEN-FW-01 – design and construct all aboveground transmission line structures following the practices suggested by APLIC.
- Company-wide Avian Protection Plan (PGE 2020).
 - Avian-safe design, consistent with Condition GEN-FW-01.
 - Consideration of risk factors for avian collisions and use of flight diverters, if warranted.
 - Internal reporting system for tracking avian mortalities.

7.5 Mitigation

After avoidance and minimization measures have been implemented, some impacts to wildlife habitat and sensitive species will remain. Permanent habitat loss will be mitigated for according to

ODFW Habitat Mitigation Policy goals and standards, as described in the RFA 5 Site Boundary HMP (Attachment G-2). Included in this plan are measures for protecting and enhancing sufficient acreages of wildlife habitat to meet the mitigation goals and standards for those habitat subtypes and categories temporarily and permanently impacted by the Solar Components. The mitigation will be in place for the duration of the Solar Components.

8.0 Monitoring Program – OAR 345-022-0060(3)(h)

(h) A description of the applicant's proposed monitoring plans to evaluate the success of the measures described in (g).

The Certificate Holder will conduct revegetation monitoring as described in the Comprehensive Solar Revegetation and Soil Management Plan (Exhibit C, Attachment C-2). The Certificate Holder will conduct habitat mitigation monitoring as described in the HMP (Attachment G-2). Monitoring for noxious weeds will occur per the Comprehensive Solar Revegetation and Soil Management Plan (Exhibit C, Attachment C-2).

No formal monitoring program is applicable to some of the measures discussed in Section 7.0. For instance, other than through professional review of the information included in this RFA, there is no monitoring program for design of the Solar Components. Similarly, adherence to speed limits will be expected but not formally monitored.

Environmental monitors will be on-site during construction to support implementation of the environmental training and sensitive resource awareness.

9.0 Conclusion

As part of the RFA and Solar Components siting process, fish and wildlife habitats within the Analysis Area were identified and categorized pursuant to OAR 635-415-0025. Based on survey results, the Solar Components were adjusted to minimize all impacts to Category 3 and 4 habitat. Unavoidable habitat impacts will be mitigated consistent with OAR 635-415-0025.

Therefore, based on the information provided in this exhibit, there is sufficient evidence upon which the Council may find that the design, construction, and operation of the Solar Components, taking into account the proposed mitigation measures, are consistent with the fish and wildlife mitigation goals and standards of OAR 635-415-0025. Accordingly, the Certificate Holder demonstrates compliance with OAR 345-022-0060.

10.0 References

- Abele, S.C., V.A. Saab, and E.O. Garton. 2004. Lewis's Woodpecker (*Melanerpes lewis*): a technical conservation assessment. [Online]. USDA Forest Service, Rocky Mountain Region.
https://www.fs.usda.gov/Internet/FSE_DOCUMENTS/stelprdb5182072.pdf
- Adams, C.A., C.C.S. Clair, E.C. Knight, and E.M. Bayne. 2024. Behaviour and landscape contexts determine the effects of artificial light on two crepuscular bird species. *Landscape Ecology* 39(4): 83. <https://doi.org/10.1007/s10980-024-01875-3>
- Adams, C.A., C.C.S. Clair, E.C. Knight, and E.M. Bayne. 2025. Artificial light alters spatial and temporal habitat use by a crepuscular aerial insectivore. *Ibis* 168(1): 308–328.
<https://doi.org/10.1111/ibi.13425>
- ADW (Animal Diversity Web). 2024. Brewer's sparrow (*Spizella breweri*) species account.
https://animaldiversity.org/accounts/Spizella_breweri/. Accessed August 2026.
- APLIC (Avian Powerline Interaction Committee). 2012. Reducing Avian Collisions with Power Lines: The State of the Art in 2012. Edison Electric Institute and APLIC. Washington, D.C.
- Bear, G.D., and R.M. Hansen. 1966. Food habits, growth and reproduction of white-tailed jackrabbits in southern Colorado. Tech. Bull. Agric. Exp. Sta., Colorado State University 90:1–59.
- Bechard, M. J., C. S. Houston, J. H. Saransola, and A. S. England. 2020. Swainson's Hawk (*Buteo swainsoni*), version 1.0. In *Birds of the World* (A. F. Poole, Editor). Cornell Lab of Ornithology, Ithaca, NY, USA. <https://doi.org/10.2173/bow.swahaw.01>
- Berteaux, D., I. Klvana, and C. Trudeau. 2005. Spring-to-Fall Mass Gain in a Northern Population of North American Porcupines. *Journal of Mammalogy* 86(3):514–519.
- Brigham, R. M., J. W. Ng, R. G. Poulin, and S. D. Grindal. 2020. Common Nighthawk (*Chordeiles minor*), version 1.0. In *Birds of the World* (A. F. Poole, Editor). Cornell Lab of Ornithology, Ithaca, NY, USA. <https://doi.org/10.2173/bow.comnig.01>
- Brunton, D.F. 1981. Nocturnal aggregations of white-tailed jack-rabbits at Rimbey, Alberta. *Blue Jay* 39:120–121.
- Buehler, D. A. 2022. Bald Eagle (*Haliaeetus leucocephalus*), version 2.0. In *Birds of the World* (P. G. Rodewald and S. G. Mlodinow, Editors). Cornell Lab of Ornithology, Ithaca, NY, USA.
<https://doi.org/10.2173/bow.baleag.02>
- Conkling, T.J., A.L. Fesnock, and T.E. Katzner. 2023. Numbers of wildlife fatalities at renewable energy facilities in a targeted development region. *PLoS ONE* 18(12): e0295552.
<https://doi.org/10.1371/journal.pone.0295552>
- Csuti, B., T.A. O'Neil, M.M. Shaughnessy, E.P. Gaines, and J.C. Hak. 2001. *Atlas of Oregon Wildlife*, 2nd edition. Oregon State University Press, Corvallis. 525 pp.

- Dugger, B. D., and K. M. Dugger. 2002. Long-billed Curlew (*Numenius americanus*), version 2.0. In *The Birds of North America* (A. F. Poole and F. B. Gill, Editors). Cornell Lab of Ornithology, Ithaca, NY, USA. <https://doi.org/10.2173/bow.lobcur.01>
- Dwyer, J. F., Murphy, R. K., Stahlecker, D. W., Dwyer, A. M. and Boal, C. W. 2020. Golden Eagle perch-site use in the U.S. Southern Plains: Understanding electrocution risk. *Journal of Raptor Research* 54(2): 126-135.
- eBird. 2026. eBird: An online database of bird distribution and abundance [web application]. eBird, Cornell Lab of Ornithology, Ithaca, New York. <https://ebird.org/map>. Accessed August 2026.
- Fink, D., T. Auer, A. Johnston, M. Strimas-Mackey, S. Ligocki, O. Robinson, W. Hochachka, L. Jaromczyk, C. Crowley, K. Dunham, A. Stillman, C. Davis, M. Stokowski, P. Sharma, V. Pantoja, D. Burgin, P. Crowe, M. Bell, S. Ray, I. Davies, V. Ruiz-Gutierrez, C. Wood, A. Rodewald. 2024. eBird Status and Trends, Data Version: 2023; Released: 2025. Cornell Lab of Ornithology, Ithaca, New York. <https://doi.org/10.2173/WZTW8903>
- Flinders, J.T., and R.M. Hansen. 1971. Diets and habitats of jackrabbits within a shortgrass ecosystem. IBP Grassland Biome Technical Report No. 98.
- Flinders, J.T. and R.M. Hansen. 1972. Diets and habitats of jackrabbits in northeastern Colorado. Range Sci. Dept. Sci. Ser., Colorado State University 12:1–29.
- Heath, S. R., E. H. Dunn, and D. J. Agro. 2020. Black Tern (*Chlidonias niger*), version 1.0. In *Birds of the World* (S. M. Billerman, Editor). Cornell Lab of Ornithology, Ithaca, NY, USA. <https://doi.org/10.2173/bow.blkter.01>
- Hill, J.M., and R.B. Renfrew. 2018. Migratory patterns and connectivity of two North American grassland bird species. *Ecology and Evolution* 9(1):680–692. <https://doi.org/10.1002/ece3.4795>. Accessed August 2026.
- Johnson, Judith. 1977. Biology of the Porcupine (*Erethizon dorsatum*) in northwestern South Dakota. *Electronic Theses and Dissertations*. 144. <https://openprairie.sdstate.edu/etd/144>
- Kagan, R.A., T.C. Viner, P.W. Trail, and E.O. Espinoza. 2014. Avian Mortality at Solar Energy Facilities in Southern California: A Preliminary Analysis. National Fish and Wildlife Forensics Laboratory.
- Kosciuch, K., D. Riser-Espinoza, M. Gerringer, and W. Erickson. 2020. A summary of bird mortality at photovoltaic utility scale solar facilities in the Southwestern U.S. *PLoS ONE* 15(4): e0232034. <https://doi.org/10.1371/journal.pone.0232034>. Accessed August 2026.
- Kosciuch, K., D. Riser-Espinoza, K. Russell, J. Sullivan, and N. Bartok. 2022. Alberta Regional Solar Fatality Analysis. Summarized the incidence of water-associated and water obligate bird fatalities observed to date at PV solar facilities in Alberta, Canada.

- Loss, S.R., T. Will, S.S. Loss, and P.P. Marra. 2014. Bird-building collisions in the United States: estimates of annual mortality and species vulnerability. *Condor* 116: 8-23.
<https://doi.org/10.1650/CONDOR-12-090>
- NatureServe. 2026. NatureServe Explorer [web application]. NatureServe, Arlington, Virginia. Available online at: <https://explorer.natureserve.org/>. Accessed August 2026.
- NRCS (Natural Resources Conservation Service). 2024. Web Soil Survey. Available online at: <http://websoilsurvey.sc.egov.usda.gov/App/WebSoilSurvey.aspx>.
- ODA (Oregon Department of Agriculture). 2022. Noxious Weed Policy and Classification System. Noxious Weed Control Program. Salem, OR.
- ODA. 2024. Noxious Weed Policy and Classification System 2024. Noxious Weed Control Program. Salem, OR. Revised 2024. Available online at: <https://www.oregon.gov/oda/weeds/oregon-noxious-weeds/Documents/NoxiousWeedPolicyClassification.pdf> Accessed August 2026.
- ODFW (Oregon Department of Fish and Wildlife). 2008. The Mitigation Category Flow Chart. Available online at: https://www.dfw.state.or.us/habitat/mitigation/mitigation_category_flow.pdf
- ODFW. 2013. Deer and Elk Winter Range for Eastern Oregon. GIS Data Files. Publish Date: 1/9/2013.
<https://nrimp.dfw.state.or.us/DataClearinghouse/default.aspx?p=202&XMLname=885.xml>.
- ODFW. 2021. Sensitive Species List. Revised October 2021.
https://www.dfw.state.or.us/wildlife/diversity/species/docs/Sensitive_Species_List.pdf. Accessed August 2026.
- ODFW. 2023. Priority Wildlife Connectivity Areas Web Map. Oregon Department of Fish and Wildlife. Salem, Oregon.
<https://experience.arcgis.com/experience/6979b6598f904951bd0af1821e1595f1/>.
- ODFW. 2026a. Oregon's State Wildlife Action Plan. 2026. Salem, Oregon. Available online at: <https://swap.oregon.gov/> Accessed August 2026.
- ODFW. 2026b. Compass Online Fish and Wildlife Mapping Tool.
<http://dfw.state.or.us/maps/compass/>. Accessed August 2026.
- ODOE (Oregon Department of Energy). 2020. Montague Wind Power Facility - Final Order on Request for Amendment 5. September 25, 2020.
- ORBIC (Oregon Biodiversity Information Center). 2023. Rare, Threatened and Endangered Vascular Plant Species of Oregon (2023). Institute for Natural Resources Publications. 158. Available online at: <https://archives.pdx.edu/ds/psu/43182>
- ORBIC. 2024. Element Occurrence Record Digital Data Set for rare, threatened, or endangered species for the state of Oregon, May 2, 2024. ORBIC, Institute for Natural Resources. Portland State University, Portland, Oregon.

- ORBIC. 2025. Rare, Threatened and Endangered Zoology Species of Oregon (2025). Institute for Natural Resources Publications. 174. Available online at: <https://archives.pdx.edu/ds/psu/44183>
- OregonFlora. 2026. OregonFlora Portal. Available online at: <https://oregonflora.org/> Accessed July 2026.
- Orion Sherman County Wind Farm, LLC. 2005. Site Certificate Application for the Biglow Canyon Wind Farm. Submitted to the Oregon Energy Facility Siting Council. October 2005.
- OSULP and INR (Oregon State University Libraries & Press and Institute for Natural Resources). 2021. Oregon Explorer. <https://oe.oregonexplorer.info/wildlife/wildlifeviewer/>. Accessed August 2026.
- PGE (Portland General Electric Company). 2020. Avian Protection Plan. Revision 2. Prepared by PGE Environmental Services.
- PGE. 2023. Biglow Canyon Wind Farm Wildlife Monitoring, Habitat Mitigation and Revegetation 2022 Monitoring Report. Prepared by PGE Environmental Services. April 2023.
- PGE. 2025. Request for Amendment 4 to the Site Certificate for the Biglow Canyon Wind Farm. Submitted to Oregon Department of Energy. December 2025.
- Rodriguez, M., D. Riser-Espinoza, W. Erickson, and M. Tuma. 2023. Post-Construction Monitoring at the Two Creeks Solar Project. Manitowoc County, Wisconsin.
- SCNWD (Sherman County Noxious Weed District). 2024. Sherman County Noxious Weeds. <https://www.co.sherman.or.us/noxious-weeds-sherman-county/>. Accessed August 2026.
- Smallwood, S. 2022. Utility-Scale Solar Impacts to Volant Wildlife. Journal of Wildlife Management. Vol.86, Issue 4. <https://doi.org/10.1002/jwmg.22216>
- StreamNet. 2026. Fish distribution data for All Fish Species. GIS Data. Portland (OR). <https://www.streamnet.org/home/data-maps/sn-mapper/>. Accessed August 2026.
- USFS and BLM (U.S. Forest Service and U.S. Bureau of Land Management). 1998. Survey and Manage Survey Protocol – Vascular Plants.
- USFWS (U.S. Fish and Wildlife Service). 2017. Biology of jackrabbits (*Lepus* spp.) as prey of golden eagles (*Aquila chrysaetos*) in the western United States. Unpublished report prepared by the Western Golden Eagle Team, U.S. Fish and Wildlife Service. Available online at: <https://ecos.fws.gov/ServCat/Reference/Profile/87136>
- USFWS. 2024. National Wetlands Inventory. Wetlands Data by State, Oregon. <https://fwsprimary.wim.usgs.gov/wetlands/apps/wetlands-mapper/>.
- USGS (U.S. Geological Survey). 2024. National Hydrography Dataset. Available online at: <http://datagateway.nrcs.usda.gov/>.
- Walston, L.J. Jr., K.E. Rollins, K.P. Smith, K.E. LaGory, K. Sinclair, C. Turchi, T. Wendelin, and H. Souder. 2015. A Review of Avian Monitoring and Mitigation Information at Existing Utility-

- Scale Solar Facilities. Prepared for U.S. Department of Energy, SunShot Initiative and Office of Energy Efficiency & Renewable Energy. April 2015. Available online at:
https://blmsolar.anl.gov/related/avian-solar/docs/Avian_Monitoring_Mitigation_Solar.pdf
- WEST (Western EcoSystems Technology, Inc.). 2005. Wildlife and Habitat Baseline Study for the Proposed Biglow Canyon Wind Power Project, Sherman County, Oregon. March 2004 - August 2005. Prepared for Orion Energy LLC., Oakland, California. Prepared October 2005.
- WEST. 2009. Biglow Canyon Wind Farm Phase I Post-Construction Avian and Bat Monitoring First Annual Report Sherman County, Oregon. January 2008 – December, 2008. Prepared for Portland General Electric Company. Prepared April 2009.
- WEST. 2010. Biglow Canyon Wind Farm Phase I Post-Construction Avian and Bat Monitoring Second Annual Report, Sherman County, Oregon. January 26, 2009 – December 11, 2009. Prepared for Portland General Electric Company. Prepared April 2010.
- WEST. 2012a. Biglow Canyon Wind Farm – Phase III, Year 1 Avian and Bat Monitoring Report, Sherman County, Oregon. September 13, 2010 – September 9, 2011. Prepared for Portland General Electric Company. Prepared April 2012.
- WEST. 2012b. Biglow Canyon Wind Farm Phase II, Amended Year 1 Post-construction Avian and Bat Monitoring Report, Sherman County, Oregon. September 10, 2009 – September 12, 2010. Prepared for Portland General Electric Company. Original report prepared January 2011. Amended June 2012.
- WEST. 2012c. Biglow Canyon Wind Farm Phase II, Year 1 Post-construction Avian and Bat Monitoring Report, Sherman County, Oregon. September 13, 2010 – September 15, 2011. Prepared for Portland General Electric Company. Prepared April 2012.
- WEST. 2013. Biglow Canyon Wind Farm – Phase III, Year 2 Avian and Bat Monitoring Report, Sherman County, Oregon. September 19, 2011 – September 18, 2012. Prepared for Portland General Electric Company. Prepared April 2013.
- WEST. 2014. Sources of Avian Mortality and Risk Factors Based on Empirical Data from Three Photovoltaic Solar Facilities. June 2014.
- Wiggins, D. A., D. W. Holt, and S. M. Leasure. 2020. Short-eared Owl (*Asio flammeus*), version 1.0. In Birds of the World (S. M. Billerman, Editor). Cornell Lab of Ornithology, Ithaca, NY, USA. <https://doi.org/10.2173/bow.sheowl.01>
- Xerces, IDFG, and WDFW (Xerces Society for Invertebrate Conservation, Idaho Department of Fish and Game, and Washington Department of Fish and Wildlife). 2021. The Pacific Northwest Bumble Bee Atlas: Summary and Species Accounts. Appendices to the Final Report Submitted to the U.S. Fish and Wildlife Service. November 2021. Available online at: https://www.xerces.org/sites/default/files/publications/21-026_01_2.pdf Accessed July 2026.

Xerces. 2026. Bumble Bee Atlas Data Highlights. Available online at:
<https://www.bumblebeeatlas.org/pages/data-highlights> Accessed August 2026.

11.0 Approval Standards and Submittal Requirements

Table G-7. Approval Standards and Submittal Requirements Matrix

Requirements	Location
OAR 345-022-0060 Fish and Wildlife Habitat	–
Approval Standards	
To issue a site certificate, the Council must find that the design, construction and operation of the facility, taking into account mitigation, are consistent with:	–
(1) The general fish and wildlife habitat mitigation goals and standards of OAR 635-415-0025(1) through (6) in effect as of February 24, 2017; and	Section 7.0
(2) For energy facilities that impact sage-grouse habitat, the sage-grouse specific habitat mitigation requirements of the Greater Sage-Grouse Conservation Strategy for Oregon at OAR 635-415-0025(7) and OAR 635-140-0000 through -0025 in effect as of February 24, 2017.	–
Submittal Requirements	
(3) To assist the Council in determining whether the standard outlined in (1) through (2) has been met, the Applicant must submit information about the fish and wildlife habitat and the fish and wildlife species, other than the species addressed in OAR-022-0070(3) (the Threatened and Endangered Species Exhibit) that could be affected by the proposed facility, providing evidence to support a finding by the Council as required by this rule. The applicant must include:	–
(a) A description of biological and botanical surveys performed that support the information in this exhibit, including a discussion of the timing and scope of each survey;	Section 2.0
(b) Identification of all fish and wildlife habitat in the analysis area, classified by the general fish and wildlife habitat categories as set forth in OAR 635-415-0025 and the sage-grouse specific habitats described in the Greater Sage-Grouse Conservation Strategy for Oregon at OAR 635-140-0000 through 635-140-0025 (core, low density, and general habitats), and a description of the characteristics and condition of that habitat in the analysis area, including a table of the areas of permanent disturbance and temporary disturbance (in acres) in each habitat category and subtype;	Section 3.0
(c) A map showing the locations of the habitat identified in (b);	Figure G-2
(d) Based on consultation with the Oregon Department of Fish and Wildlife (ODFW) and appropriate field study and literature review, identification of all State Sensitive Species that might be present in the analysis area and a discussion of any site-specific issues of concern to ODFW;	Section 4.0
(e) A baseline survey of the use of habitat in the analysis area by species identified in (d) performed according to a protocol approved by the Department and ODFW;	Section 5.0
(f) A description of the nature, extent and duration of potential adverse impacts on the habitat identified in (b) and species identified in (d) that could result from construction, operation and retirement of the proposed facility;	Section 6.0

Requirements	Location
(g) A description of any measures proposed by the applicant to avoid, reduce, or mitigate the potential adverse impacts described in (f) in accordance with the general fish and wildlife habitat mitigation goals and standards described in OAR 635-415-0025 and a description of any measures proposed by the applicant to avoid, minimize, and provide compensatory mitigation for the potential adverse impacts described in (f) in accordance with the sage-grouse specific habitat mitigation requirements described in the Greater Sage-Grouse Conservation Strategy for Oregon at OAR 635-140-0000 through 635-140-0025, and a discussion of how the proposed measures would achieve those goals and requirements; and	Section 7.0
(h) A description of the applicant's proposed monitoring plans to evaluate the success of the measures described in (g).	Section 8.0

Figures

**Biglow Canyon
Wind Farm Request
for Amendment #5**

**Figure G-1
Fish and Wildlife
Analysis Area**

SHERMAN COUNTY, OR

-  RFA5 Site Boundary
-  Analysis Area (0.5-mile Buffer)
-  State Boundary
-  County Boundary
-  City/Town



Reference Map




\\tt-local\GIS\US\CES\Projects\PDX\BrightNight\BiglowCanyon\Maps_RFA5\Exhibit_GIPGE_BiglowCanyon_RFA5_ExhibitG_20260810.aprx




1:35,000

WGS 1984 UTM Zone 10N



NOT FOR CONSTRUCTION

\\tlocal\GIS\US\CES\Projects\PDX\BrightNight\BiglowCanyon\Maps_RFA5\Exhibit_GIPGE_BiglowCanyon_RFA5_ExhibitG_20260810.aprx




**Biglow Canyon
Wind Farm Request
for Amendment #5**

**Figure G-2
2024-2026 Survey Area
in the
Facility Site Boundary**


SHERMAN COUNTY, OR

- RFA5 Site Boundary
- State Boundary
- County Boundary
- City/Town
- 2026 Raptor Nest Survey Area
- 2024 Wetlands and Waters Survey Area
- 2024 Botanical Survey Area
- 2024 Habitat, Sensitive Species, Raptor Nest Survey Area




1:50,000

WGS 1984 UTM Zone 10N



NOT FOR CONSTRUCTION

\\tt.local\GIS\US\CES\Projects\PDX\BrightNight\BiglowCanyon\Maps_RFA5\Exhibit_GIPGE_BiglowCanyon_RFA5_ExhibitG_20260810.aprx



**Biglow Canyon
Wind Farm Request
for Amendment #5**


**Figure G-3
Habitat Categories
in the Analysis Area**

SHERMAN COUNTY, OR

- RFA5 Site Boundary
 - Analysis Area (0.5-mile Buffer)
 - State Boundary
 - County Boundary
 - City/Town
- Final Habitat Category
- 3
 - 4
 - 6




Reference Map




1:30,000

WGS 1984 UTM Zone 10N



NOT FOR CONSTRUCTION

\\tt.local\GIS\US\CES\Projects\PDX\BrightNight\BiglowCanyon\Maps_RFA5\Exhibit_GIPGE_BiglowCanyon_RFA5_ExhibitG_20260730.aprx



**Biglow Canyon
Wind Farm Request
for Amendment #5**

**Figure G-4
Habitat Subtypes
in the Analysis Area**

SHERMAN COUNTY, OR

- RFA5 Site Boundary
- Analysis Area (0.5-mile Buffer)
- State Boundary
- County Boundary
- City/Town
- Habitat Subtype**
 - Irrigated Cropland
 - Non-Irrigated Cropland
 - Developed
 - Grasslands
 - Intermittent Streams
 - Freshwater Emergent Wetlands
 - Upland Trees




Reference Map




1:30,000

WGS 1984 UTM Zone 10N



NOT FOR CONSTRUCTION

\\t.local\GIS\US\CES\Projects\BrightNight\BiglowCanyon\Maps_RFA5\Exhibit_GIPGE_BiglowCanyon_RFA5_20260810.aprx



**Biglow Canyon
Wind Farm Request
for Amendment #5**

**Figure G-5
2024-2026 Special Status
Species Observations
in the Analysis Area
and its Vicinity
SHERMAN COUNTY, OR**

- RFA5 Site Boundary
- Analysis Area (0.5-mile Buffer)
- State Boundary
- County Boundary
- City/Town
- Loggerhead Shrike
- Porcupine




Reference Map



1:33,000

WGS 1984 UTM Zone 10N



NOT FOR CONSTRUCTION

Attachment G-1. Biological Survey Reports

2024 Botanical Survey Report

2024 Botanical Survey Report

**Biglow Canyon Wind Farm
Request for Amendment 4
November 2024**

Prepared for



BIGL bn, LLC

Prepared by



TETRA TECH

This page intentionally left blank

Table of Contents

1.0	Introduction	1
2.0	Methods	2
2.1	Survey Area.....	2
2.2	Background Review.....	2
2.2.1	Threatened, Endangered, and Candidate Plants	2
2.2.2	Noxious Weeds.....	3
2.3	Field Survey Methods	3
2.3.1	Threatened, Endangered, and Candidate Plants	3
2.3.2	Noxious Weeds.....	4
3.0	Results	5
3.1	Threatened, Endangered, and Candidate Plants	5
3.2	Noxious Weeds	5
4.0	Conclusions	7
5.0	References	7

List of Tables

Table 1. Threatened, Endangered, and Candidate Vascular Plant Species with Potential to Occur at the Solar Components.....	3
Table 2. Noxious Weeds Observed Within the Survey Area	5

List of Figures

- Figure 1. Project Location
- Figure 2. Noxious Weed Observations

List of Attachments

- Attachment 1. Vascular Plants Observed During 2024 Field Surveys

Acronyms and Abbreviations

AC	alternating current
Solar Micrositing Area	Site Boundary Subject to RFA 4
BCWF or Existing Facility	Biglow Canyon Wind Farm
BIGL	BIGL bn, LLC
Certificate Holder / PGE	Portland General Electric Company
Council	Oregon Energy Facility Siting Council
ESA	Endangered Species Act
GPS	Global Positioning System
IPaC	Information for Planning and Consultation
MW	megawatt
OAR	Oregon Administrative Rule
ODA	Oregon Department of Agriculture
ORBIC	Oregon Biodiversity Information Center
RFA	Request for Amendment
Tetra Tech	Tetra Tech, Inc.
USFWS	U.S. Fish and Wildlife Service

1.0 Introduction

The Portland General Electric Company (PGE or Certificate Holder) is submitting a Request for Amendment (RFA) 4 to the Site Certificate on Amendment 3, issued October 31, 2008 (Site Certificate) for the Biglow Canyon Wind Farm (BCWF or Existing Facility) to add photovoltaic solar energy generation and battery storage (Solar Components) to the operating BCWF in Sherman County, OR.

BCWF, owned and operated by PGE, is located within an approved site boundary comprising approximately 25,000 acres, approximately 2.5 miles northeast of the town of Wasco in Sherman County, Oregon (Figure 1). The BCWF operates under the Site Certificate from the Oregon Energy Facility Siting Council (Council or EFSC) as administered by the Oregon Department of Energy. BCWF currently consists of 217 wind turbines, with a maximum blade tip height of 445 feet, and a peak generating capacity of 450 megawatts (MW).

In RFA 4, PGE proposes to add up to 385 MW alternating current (AC) generating capacity from photovoltaic solar arrays and 375 MW in battery storage capacity. RFA 4 seeks to expand the BCWF site boundary to include the Solar Components in portions of the existing site boundary and in the proposed expanded site boundary (together, Solar Micrositing Area or RFA 4 Site Boundary).

The Solar Micrositing Area is approximately 3,980 acres and provides a conservative estimate of the maximum area needed for development, micrositing, and temporary disturbances from the Solar Components during construction, rather than the anticipated disturbance footprint. Solar Components will include solar arrays, inverters, battery energy storage system facilities and their subcomponents (i.e., inverters), two collector substations, a total of approximately 3 miles of 230-kilovolt generation tie transmission line, medium voltage collector lines, operations and maintenance structures, site access roads, internal roads, perimeter fencing, facility entry gates, and temporary laydown areas. The maximum generating capacity from the Solar Components will be 385 MW AC and construction may take place in phases.

PGE will own and operate the Solar Components as a part of the BCWF, which, to date, have been developed by BIGL bn,LLC (BIGL). BIGL, in its capacity as the project developer, supports PGE in this RFA 4 and may construct and temporarily operate the Solar Components on behalf of PGE under a Build-Transfer Agreement. Tetra Tech Inc. (Tetra Tech) is providing support to PGE and BIGL through preparation of RFA 4.

This survey report presents the methods and results of the 2024 botanical surveys conducted by Tetra Tech for the Solar Micrositing Area. The purpose of the botanical surveys was to document the presence of federally or state listed endangered, threatened, and candidate vascular plant species and state and county-designated noxious weeds within the Solar Micrositing Area. The surveys were designed in accordance with Council standards set forth in Oregon Administrative Rules (OAR) 345-021-0010 (1)(p) and (a), OAR 345-022-0070, and in coordination with the Oregon Department of Agriculture (ODA).

2.0 Methods

The background review and field survey methods described in this section were submitted to ODA for review on June 10, 2024. ODA provided concurrence with these methods on June 13, 2024, prior to Tetra Tech conducting field surveys (J. Brown, personal communication, June 13, 2024).

2.1 Survey Area

The Survey Area for the 2024 botanical surveys consisted of the 3,980-acre Solar Micrositing Area as shown on Figure 1. All areas of suitable or marginally suitable habitat within the Solar Micrositing Area (i.e., Survey Area) were surveyed during botanical surveys conducted in 2024. Only active agricultural lands and developed lands (e.g., existing roads, existing buildings) were excluded from surveys.

2.2 Background Review

2.2.1 *Threatened, Endangered, and Candidate Plants*

Tetra Tech conducted a review of existing information to identify federal and state endangered, threatened, proposed, and candidate plant species with the potential to occur within the vicinity of the Survey Area. Sources of information included the following:

- U.S. Fish and Wildlife Service’s (USFWS) Information for Planning and Consultation (IPaC) Resource List for the Survey Area (USFWS 2024a);
- USFWS threatened, endangered, and candidate species list for Oregon (USFWS 2024b);
- Oregon Listed Plants by County (ODA 2024a);
- Oregon Biodiversity Information Center (ORBIC) 2019 Rare, Threatened and Endangered Species of Oregon (ORBIC 2019);
- ORBIC Rare, Threatened and Endangered Vascular Plant Species of Oregon (ORBIC 2023);
- ORBIC Element Occurrence Records for the vicinity of the Project (ORBIC 2024); and
- OregonFlora’s Online Guide to the Vascular Plants of Oregon (OregonFlora 2024).

The initial list of potential primary target species included all vascular plant species listed as endangered, threatened, proposed, or candidate for listing by the USFWS under the federal Endangered Species Act (ESA), or by the ODA under the Oregon ESA, that occur or have potential to occur within or near the Solar Components. Based on this review, Tetra Tech determined that no federally listed, proposed, or candidate plant species have the potential to occur within or near the Proposed Facility; however, one state-listed threatened and three state candidate vascular plant species have potential to occur within or near the Solar Components (Table 1).

Table 1. Threatened, Endangered, and Candidate Vascular Plant Species with Potential to Occur at the Solar Components

Scientific Name	Common Name	State Status ¹	Habitat	Survey Period
<i>Astragalus collinus</i> var. <i>laurentii</i>	Lawrence's milkvetch	T	Bunchgrass prairies, roadsides; sandy or rocky soils overlying basalt on dry slopes.	Surveys should be conducted when species is fruiting; typically late May to August
<i>Eriocoma</i> [<i>Achnatherum</i>] <i>hendersonii</i>	Henderson's ricegrass	C	Dry, shallow rocky soils described from basalt in sagebrush or ponderosa pine. Soils are often subject to frost heave.	May to June
<i>Erythranthe jungermannioides</i>	Hepatic monkeyflower	C	Moist basalt crevices and seeps in vertical cliff faces and canyon walls.	May (June) to August (depending on hydrology)
<i>Myosurus sessilis</i>	Sessile mousetail	C	Vernal pools and alkali flats.	April to June (depending on hydrology)
Sources: ODA 2024a, 2024b; OregonFlora 2024. 1. T = Threatened, C = Candidate for listing.				

2.2.2 Noxious Weeds

Prior to field surveys, Tetra Tech reviewed lists of species designated as noxious weeds in Oregon state and Sherman County (ODA 2022; SCNWD 2024). Existing literature and other sources were also reviewed to familiarize surveyors with identification of designated noxious weeds that would potentially be encountered within the Survey Area. Sources of information reviewed included:

- OregonFlora online guide to vascular plants of Oregon (OregonFlora 2024);
- Oregon Noxious Weed Profiles (ODA 2024c); and
- Oregon WeedMapper (ODA 2024d).

2.3 Field Survey Methods

2.3.1 Threatened, Endangered, and Candidate Plants

Field surveys were conducted in early July. The survey schedule was chosen based on the recommended survey period for Lawrence's milkvetch (*Astragalus collinus* var. *laurentii*), the primary target plant species with potential to occur in the Solar Components vicinity. Surveys were conducted by a Tetra Tech senior botanist familiar with suitable habitat for and identification of Lawrence's milkvetch as well as the other target plant species listed in Table 1. Prior to conducting

field surveys, fact sheets for the target plant species were compiled. These fact sheets were used by the surveyor in the field and included the following:

- Photos of each target species and its habitat;
- Information detailing habitat associations;
- Range and flowering period;
- Identifying features; and
- Characteristics distinguishing target species from similar species within its range.

Tetra Tech conducted botanical field surveys using the Intuitive Controlled survey method, a standard and commonly accepted survey protocol (USFS and BLM 1998). This method incorporates meandering transects that traverse the survey area, and that target the full array of major vegetation types, aspects, topographical features, habitats, and substrate types. While en route, the surveyors search for target species, and when the surveyors arrive at an area of high potential habitat (that was defined in the pre-field review or encountered during the field visit), they conduct a complete survey for the target species. Complete surveys include an examination of 100 percent of the habitat.

Standard Tetra Tech survey protocol includes recording Global Positioning System (GPS) locations of any target species encountered with a tablet using ArcGIS FieldMaps software. Survey methods also include completing ORBIC siting forms for any rare plant populations observed and taking photos to serve as digital specimen vouchers to illustrate identifying characteristics, plant habits, and habitat.

Data collected for each rare plant population, if encountered, would include:

- Species phenology;
- Number of plants observed;
- Habitat information and associated species; and
- Visible threats.

During surveys, Tetra Tech maintained a running list of vascular plant species encountered and made informal collections of unknown species for later identification. Identification was verified by the use of appropriate plant keys; in particular, *Flora of the Pacific Northwest* (Hitchcock and Cronquist 2018). Nomenclature follows that used by OregonFlora (2024). The final vascular plant species list for the surveys is included as Attachment 1 of this report.

2.3.2 Noxious Weeds

Noxious weed surveys were conducted concurrently with surveys for threatened, endangered, and candidate plant species. During the surveys, surveyors documented observations of state and county-listed noxious weeds within the survey area. When a noxious weed was encountered, the

location was recorded with a GPS point and the species, estimated size of infestation (i.e., small – less than 0.1 acre, medium – 0.1 to 1 acre, or large – 1 to 5 acres), and relative abundance (i.e., sparse [only a few individuals noted or low cover of species in area], common [many individuals of the species noted in area], or very high cover [dense population of the species]) was recorded.

3.0 Results

Field surveys were conducted on July 9 and 10, 2024. During the surveys, Tetra Tech observed that vegetation within much of the Survey Area has been modified due to historic and current agricultural activity, construction and management of the existing wind energy facility, and historic and current grazing activity. Non-native, invasive grasses and forbs are common throughout the Survey Area due to these land uses. Only limited areas of native habitat types (e.g., shrub-steppe) occur within the Survey Area and even in these areas, non-native, invasive grasses and forbs, including cheatgrass (*Bromus tectorum*), bulbous bluegrass (*Poa bulbosa*), cereal rye (*Secale cereale*), and prickly lettuce (*Lactuca serriola*) were often abundant.

3.1 Threatened, Endangered, and Candidate Plants

No threatened, endangered, proposed, or candidate vascular plant species were observed within the Survey Area. In addition, no suitable habitat for Henderson’s ricegrass (*Eriocoma* [*Achnatherum*] *hendersonii*), hepatic monkeyflower (*Erythranthe jungermannioides*), and sessile mousetail (*Myosurus sessilis*), and limited suitable habitat for Lawrence’s milkvetch, was observed within the Survey Area. As noted above, historic and current anthropogenic activities have resulted in degradation of the limited suitable habitat for Lawrence’s milkvetch within the Survey Area.

3.2 Noxious Weeds

Tetra Tech recorded 12 listed noxious weed species within the Survey Area. Table 2 lists the noxious weed species observed, their noxious weed designation, and the frequency of observations. Figure 2 displays the locations of noxious weeds observed during the field surveys.

Table 2. Noxious Weeds Observed Within the Survey Area

Scientific Name	Common Name	Status (State ¹ / Sherman County ²)	Frequency ³
<i>Aegilops cylindrica</i>	Jointed goatgrass	B / C	Infrequently observed. Observations primarily in southern portion of Survey Area.
<i>Centaurea diffusa</i>	Diffuse knapweed	B* / B (T)	Several small to medium-sized patches observed.
<i>Chondrilla juncea</i>	Rush skeletonweed	B (T)* / A (T)	Frequently observed in non-cultivated portions of Survey Area.
<i>Cirsium arvense</i>	Canada thistle	B* / B (T)	One observation in southern portion of Survey Area.

Scientific Name	Common Name	Status (State ¹ / Sherman County ²)	Frequency ³
<i>Cirsium vulgare</i>	Bull thistle	B* / C	Observed in two locations.
<i>Convolvulus arvensis</i>	Field bindweed	B* / B	Many small to large patches observed.
<i>Conyza canadensis</i>	Horseweed, marestail	Not listed / C	Commonly observed within Survey Area.
<i>Lactuca serriola</i>	Prickly lettuce	Not listed / C	Frequently observed within Survey Area.
<i>Onopordum acanthium</i>	Scotch thistle	B / B (T)	Several observations in northeastern portion of Survey Area.
<i>Rhaponticum</i> (<i>Acroptilon</i>) <i>repens</i>	Russian knapweed	B* / B	One observation in northeastern portion of Survey Area.
<i>Salsola tragus</i>	Russian thistle	Not listed / C	Frequently observed within Survey Area.
<i>Secale cereale</i>	Common rye, cereal rye	Not listed / C	Frequently observed within Survey Area.
<p>1. "B" Listed weeds: Weeds of economic importance, which are regionally abundant, but which may have limited distribution in some counties. "T" Designated weeds: designated group of weed species selected from either the A or B list as a focus for prevention and control by the Noxious Weed Control Program. Action against these weeds will receive priority. T-designated noxious weeds are determined by the Oregon State Weed Board and directs ODA to develop and implement a statewide management plan. Species marked with an (*) are targeted for biocontrol (ODA 2022).</p> <p>2. "A" Class: High Priority. Any noxious weed which greatly endangers the overall economic well – being of the County and has a small enough distribution where eradication is possible. "B" Class: Moderate Priority. A noxious weed which is well established in the County and has known negative impacts, but due to its distribution, eradication is not feasible. "C" Class – Low Priority. A noxious weed which is widespread throughout the County and has known economic impacts. "T" Class – Targeted List. A noxious weed from any Class that the Weed Advisory Board wishes to focus efforts and resources on. This T class list is reviewed annually. (SCNWD 2024).</p> <p>3. Patch/observation sizes are as follows : small = less than 0.1 acre, medium = 0.1 to 1.0 acre, large = greater than 1 acre.</p>			

Four noxious weed species were frequently observed in non-cultivated areas throughout the Survey Area: common rye (*Secale cereale*), prickly lettuce (*Lactuca serriola*), rush skeletonweed (*Chondrilla juncea*), and Russian thistle (*Salsola tragus*) (Figure 2). Observations of common rye typically consisted of large, dense infestations. Observations of prickly lettuce typically consisted of large, sparse to moderately dense infestations and observations of Russian thistle ranged from small, sparse to large, moderately dense infestations. Most observations of rush skeletonweed consisted of small or medium-sized sparse to moderately dense infestations, although one large infestation was also observed.

Diffuse knapweed (*Centaurea diffusa*), field bindweed (*Convolvulus arvensis*), marestail (*Conyza canadensis*) were all observed in several locations within the Survey Area. Diffuse knapweed (*Centaurea diffusa*) was primarily observed within the northeastern portion of the Survey Area, with observations ranging from small, sparse infestations to medium-sized, moderately dense infestations. Field bindweed was observed in the central and northern portion of Survey Area, with observations ranging from small, sparse infestations to moderately dense large infestations. Marestail was observed in the northern and southern portions of the Survey Area with observations ranging from sparse, medium to large, moderately dense infestations.

Jointed goatgrass (*Aegilops cylindrica*) and Scotch thistle (*Onopordum acanthium*) were each observed in four locations within the Survey Area. Three of the observations of jointed goatgrass were in the southern portion of the Survey Area and the fourth was in the central portion (Figure 2). The observation in the central portion of the Survey Area was a small, sparse infestation, whereas the observations in the southern portion consisted of medium-sized, moderately dense infestations and large, sparse infestations. All four observations of Scotch thistle were located in the northeastern portion of the Survey Area. While all four of these observations were small, they ranged in density from just a few individuals to dense cover of Scotch thistle.

The remaining three noxious weeds—bull thistle (*Cirsium vulgare*), Canada thistle (*Cirsium arvense*), and Russian knapweed (*Rhaponticum* [*Acroptilon*] *repens*)—were only observed in one to three locations within the Survey Area (Table 2). Bull thistle was observed in two locations within the Survey Area. The observation in the northeast portion of the Survey Area consisted of just a few individuals in a small area, whereas the observation in the southern portion of the Survey Area consisted of a medium-size, moderately dense infestation. Canada thistle and Russian knapweed were each observed in just one location. Both observations consisted of medium-sized, moderately dense infestation. The observation of Canada thistle was observed near a pond in the southern portion of the Survey Area and the observation of Russian knapweed was located along a road in the northeastern portion of the Survey Area.

4.0 Conclusions

Tetra Tech did not observe state listed or candidate vascular plant species during botanical surveys in 2024. Tetra Tech documented 12 state and/or county designated noxious weeds within the Survey Area, 4 of which were abundant throughout the Survey Area: common rye, prickly lettuce, rush skeletonweed, and Russian thistle.

5.0 References


- Hitchcock, C. L., and A. Cronquist. 2018. *Flora of the Pacific Northwest*. University of Washington Press; second edition. Seattle, WA.
- ODA (Oregon Department of Agriculture). 2022. Noxious Weed Policy and Classification System. Noxious Weed Control Program. Salem, OR. Available online at: <https://www.oregon.gov/oda/shared/Documents/Publications/Weeds/NoxiousWeedPolicyClassification.pdf>. Accessed June 2024.
- ODA.2024a. Oregon Listed Plants by County. Available online at: <https://www.oregon.gov/oda/programs/PlantConservation/Pages/ListedPlants.aspx>. Accessed April 2024.

- ODA. 2024b. Lawrence's milkvetch (*Astragalus collinus* var. *laurentii*) Fact Sheet. Available online at:
<http://www.oregon.gov/ODA/shared/Documents/Publications/PlantConservation/AstragalusCollinusLaurentiiProfile.pdf>. Accessed June 2024.
- ODA. 2024c. Oregon Noxious Weed Profiles. Available online at:
<https://www.oregon.gov/oda/programs/weeds/oregonnoxiousweeds/pages/aboutoregonweeds.aspx>. Accessed June 2024.
- ODA. 2024d. Oregon WeedMapper. Available online at:
<https://www.oregon.gov/oda/programs/Weeds/Pages/WeedMapper.aspx>. Accessed June 2024.
- ORBIC (Oregon Biodiversity Information Center). 2019. Rare, Threatened and Endangered Species of Oregon. Institute for Natural Resources, Portland State University, Portland, Oregon. 136 pp.
- ORBIC. 2023. Rare, Threatened and Endangered Vascular Plant Species of Oregon – An excerpt of the Rare, Threatened, and Endangered Species of Oregon publication. Available online at:
<https://inr.oregonstate.edu/orbic/rare-species/rare-species-oregon-publications>. Accessed April 2024.
- ORBIC. 2024. Element Occurrence Record Digital Data Set for Rare, Threatened and Endangered Species for the Biglow Solar Project in Sherman County. ORBIC, Institute for Natural Resources, Portland State University. Portland, OR. Received May 2, 2024.
- OregonFlora. 2024. Comprehensive Guide to the Vascular Plants of Oregon. Oregon State University. Corvallis, OR. Available online at: <https://oregonflora.org>. Accessed May 2024.
- SCNWD (Sherman County Noxious Weed District). 2024. Sherman County Noxious Weeds. Available online at: <https://www.co.sherman.or.us/noxious-weeds-sherman-county/>. Accessed June 2024.
- USFS (U.S. Forest Service) and BLM (U.S. Bureau of Land Management). 1998. Survey and Manage Survey Protocol – Vascular Plants.
- USFWS (U.S. Fish and Wildlife Service) 2024a. IPaC – Information for Planning and Consultation Resource List for the Project location in Wasco County, Oregon. Available online at:
<https://ipac.ecosphere.fws.gov/location/ECIIWBQU5JGEJN7MSYTNJNMQ4M/resources>. Accessed April 2024.
- USFWS. 2024b. Federally Listed, Proposed, Candidate, Delisted Species and Species of Concern under the Jurisdiction of the Fish and Wildlife Service which May Occur in Oregon. Available online at:
https://www.fws.gov/sites/default/files/documents/oregonspeciesstatelist_0.pdf. Accessed April 2024.

Figures

This page intentionally left blank

\\css706g\ists\ICES\Projects\PD\BrightNight\BiglowCanyon\Maps\NoxiousWeedsReport_20240828\BrightNight_Biglow_NoxiousWeeds_20241212.aprx



Biglow Canyon Wind Farm Request for Amendment 4

Figure 1
Project Location

SHERMAN COUNTY, OR

- Solar Micrositing Area/
Survey Area
- Proposed Transmission
Interconnect
- Existing Transmission
Lines (230-kV)
- Existing Biglow Canyon
Substation
- Existing Biglow Canyon
Wind Farm Turbine
- Oregon Energy Facility
Council Facilities
(Operating)**
 - Biglow Canyon Wind
Farm
 - Golden Hills Wind Project
 - Klondike III Wind Project
- Land Status**
 - Bureau of Land
Management
 - Private




Reference Map




1:55,000

WGS 1984 UTM Zone 10N



NOT FOR CONSTRUCTION

\\css706g\ists\ICES\Projects\PD\BrightNight\BiglowCanyon\Maps\NoxiousWeeds\Report_20240828\BrightNight_Biglow_NoxiousWeeds_20241212.aprx




Biglow Canyon Wind Farm Request for Amendment 4

Figure 2 Noxious Weed Observations

SHERMAN COUNTY, OR

- Map Grid
- Solar Micrositing Area/
Survey Area
- City/Town
- US Highway
- State Highway
- Local Roads
- County Boundary



\\cess706g\ists\1\CES\Projects\BRIGHTNight\BiglowCanyon\Maps\NoxiousWeedsReport_20240828\BrightNight_Biglow_NoxiousWeeds_20241212.aprx


- Noxious Weed
- Common Rye (*Secale cereale*)
 - Field Blindweed (*Convolvulus arvensis*)
 - Prickly Lettuce (*Lactuca serriola*)
 - Rush Skeletonweed (*Chondrilla juncea*)
 - Russian Thistle (*Salsola tragus*)
- Population Size
- <0.1 Acres
 - 0.1-1 Acres
 - 1-5 Acres

Biglow Canyon
Wind Farm Request
for Amendment 4


Figure 2.1
Noxious Weed
Observations

SHERMAN COUNTY, OR

- Solar Micrositing Area/
Survey Area
- Local Roads



Reference Map




1:6,000

WGS 1984 UTM Zone 10N

0 0.13 0.25 0.5 Miles

NOT FOR CONSTRUCTION

\\css706glst1\ICES\Projects\PD\BrightNight\BiglowCanyon\Maps\NoxiousWeedsReport_20240828\BrightNight_Biglow_NoxiousWeeds_20241212.aprx



Noxious Weed

- Common Rye (*Secale cereale*)
- Field Blindweed (*Convolvulus arvensis*)
- Prickly Lettuce (*Lactuca serriola*)
- Rush Skeletonweed (*Chondrilla juncea*)

Population Size

- <0.1 Acres
- 1-5 Acres

**Biglow Canyon
Wind Farm Request
for Amendment 4**


**Figure 2.2
Noxious Weed
Observations**

SHERMAN COUNTY, OR

□ Solar Micrositing Area/
Survey Area


— Local Roads

TETRA TECH **BRIGHTNIGHT**
Power when you need it



\\cess706glst1\ICES\Projects\1\CES\Projects\1\BrightNight\BiglowCanyon\Maps\NoxiousWeedsReport_20240828\BrightNight_Biglow_NoxiousWeeds_20241212.aprx

- Noxious Weed
- Bull Thistle (*Cirsium vulgare*)
- Common Rye (*Secale cereale*)
- Diffuse Knapweed (*Centaurea diffusa*)
- Field Blindweed (*Convolvulus arvensis*)
- Marestalk (*Conyza canadensis*)
- Prickly Lettuce (*Lactuca serriola*)
- Rush Skeletonweed (*Chondrilla juncea*)
- Russian Knapweed (*Rhaponticum repens*)
- Russian Thistle (*Salsola tragus*)
- Scotch Thistle (*Onopordum acanthium*)
- Population Size
- <0.1 Acres
- 0.1-1 Acres
- 1-5 Acres



Biglow Canyon
Wind Farm Request
for Amendment 4


Figure 2.3
Noxious Weed
Observations

SHERMAN COUNTY, OR

- Solar Micrositing Area/
Survey Area
- Local Roads




Reference Map



1:6,000

WGS 1984 UTM Zone 10N



NOT FOR CONSTRUCTION

\\cess706g\ists\1CES\Projects\1CES\BrightNight\BiglowCanyon\Maps\NoxiousWeedsReport_20240828\BrightNight_Biglow_NoxiousWeeds_20241212.aprx

Noxious Weed

●

 Common Rye (*Secale cereale*)

●

 Field Blindweed (*Convolvulus arvensis*)

●

 Jointed Goatgrass (*Aegilops cylindrica*)

Population Size

○

 <0.1 Acres

○

 0.1-1 Acres



Biglow Canyon
Wind Farm Request
for Amendment 4

Figure 2.4
Noxious Weed
Observations

SHERMAN COUNTY, OR

□

 Solar Micrositing Area/
Survey Area

—


 Local Roads

Tt

 TETRA TECH

BRIGHTNIGHT

Power when you need it



\\cess706g\ists\1\CES\Projects\PD\BrightNight\BiglowCanyon\Maps\NoxiousWeedsReport_20240828\BrightNight_Biglow_NoxiousWeeds_20241212.aprx

Noxious Weed

●

 Prickly Lettuce (*Lactuca serriola*)

Population Size

○

 1-5 Acres



Biglow Canyon
Wind Farm Request
for Amendment 4

Figure 2.5
Noxious Weed
Observations

SHERMAN COUNTY, OR

Solar Micrositing Area/
Survey Area


Local Roads

Tt

 TETRA TECH

BRIGHTNIGHT

Power when you need it



\\ces706glst1\ICES\Projects\PD\BrightNight\BiglowCanyon\Maps\NoxiousWeedsReport_20240828\BrightNight_Biglow_NoxiousWeeds_20241212.aprx

Noxious Weed

- Bull Thistle (*Cirsium vulgare*)
- Canada Thistle (*Cirsium arvense*)
- Common Rye (*Secale cereale*)
- Diffuse Knapweed (*Centaurea diffusa*)
- Jointed Goatgrass (*Aegilops cylindrica*)
- Marestail (*Conyza canadensis*)
- Prickly Lettuce (*Lactuca serriola*)
- Rush Skeletonweed (*Chondrilla juncea*)
- Russian Thistle (*Salsola tragus*)

Population Size

- <0.1 Acres
- 0.1-1 Acres
- 1-5 Acres

Biglow Canyon
Wind Farm Request
for Amendment 4


Figure 2.6
Noxious Weed
Observations

SHERMAN COUNTY, OR

- Solar Micrositing Area/
Survey Area
- Local Roads




Reference Map



1:6,000

WGS 1984 UTM Zone 10N



NOT FOR CONSTRUCTION

Armstrong St

Hilderbrand Ln
Old Wasco-Heppner Hwy

Attachment 1. Vascular Plants Observed During 2024 Field Surveys

This page intentionally left blank

Attachment 1. Vascular Plants Observed During 2024 Field Surveys

Scientific Name	Common Name	Native or Introduced? ¹	Synonym	Notes
<i>Achillea millefolium</i>	common yarrow, milfoil	N		
<i>Aegilops cylindrica</i>	jointed goatgrass	I		ODA Noxious Weed, B List; Sherman County Noxious Weed, B List
<i>Agoseris grandiflora</i>	large-flowered agoseris	N		
<i>Agoseris heterophylla</i>	annual agoseris	N		
<i>Agropyron cristatum</i>	crested wheatgrass	I		
<i>Amaranthus albus</i>	tumbling pigweed	I		
<i>Amaranthus blitoides</i>	prostrate pigweed, tumbleweed amaranth	U		
<i>Amsinckia</i> spp.	fiddleneck	N		
<i>Artemisia tridentata</i>	big sagebrush	N		
<i>Asclepias fascicularis</i>	narrowleaf milkweed	N		
<i>Astragalus purshii</i>	Pursh's milkvetch, woollypod milkvetch	N		
<i>Balsamorhiza sagittata</i>	arrowleaf balsamroot	N		
<i>Bromus hordeaceus</i>	soft chess	I	<i>Bromus mollis</i>	
<i>Bromus diandrus</i>	ripgut brome	I		
<i>Bromus tectorum</i>	cheatgrass, downy brome, downy chess	I		
<i>Centaurea diffusa</i>	diffuse knapweed, tumble knapweed	I		ODA and Sherman County Noxious Weed, B List
<i>Chenopodium album</i>	lamb's quarters, pigweed	I		
<i>Chondrilla juncea</i>	rush skeletonweed	I		ODA Noxious Weed, B List; Sherman County Noxious Weed, A List
<i>Chrysothamnus viscidiflorus</i>	green rabbitbrush, sticky-flowered rabbitbrush	N		
<i>Cirsium arvense</i>	Canada thistle, creeping thistle	I		ODA Noxious Weed, B List
<i>Cirsium vulgare</i>	bull thistle, common thistle	I		ODA and Sherman County Noxious Weed, B List
<i>Convolvulus arvensis</i>	field bindweed, field morning-glory	I		ODA and Sherman County Noxious Weed, B List
<i>Conyza canadensis</i>	Canadian fleabane, horseweed, marestail	N		Sherman County Noxious Weed, C List
<i>Crataegus douglasii</i>	black hawthorn, Douglas' hawthorn	N		
<i>Crepis atrabarba</i>	slender hawksbeard	N		
<i>Crepis</i> spp.	hawksbeard	N		
<i>Descurainia sophia</i>	flixweed, herb Sophia	I		
<i>Draba verna</i>	spring draba, spring whitlow grass	N		
<i>Elaeagnus angustifolia</i>	oleaster, Russian olive	I		
<i>Eleocharis palustris</i>	common spikerush, marsh spikerush, creeping spikerush	N		
<i>Eleocharis</i> sp.	spikerush	N		
<i>Elymus elymoides</i>	squirreltail	N	<i>Sitanion hystrix</i>	
<i>Elymus trachycaulus</i> ssp. <i>trachycaulus</i>	slender wheatgrass	N		
<i>Epilobium brachycarpum</i>	tall annual willowherb, autumn willowherb	N	<i>Epilobium paniculatum</i>	
<i>Ericameria nauseosa</i>	rubber rabbitbrush	N	<i>Chrysothamnus nauseosus</i>	
<i>Erigeron filifolius</i>	threadleaf fleabane	N		
<i>Eriocoma thurberiana</i>	Thurber's needlegrass	N	<i>Achnatherum thurberianum</i> , <i>Stipa thurberiana</i>	
<i>Eriogonum niveum</i>	snow buckwheat	I		
<i>Erodium cicutarium</i>	redstem filaree	I		

Scientific Name	Common Name	Native or Introduced? ¹	Synonym	Notes
<i>Festuca idahoensis</i>	Idaho fescue	N		
<i>Galium aparine</i>	stickywilly, cleavers, common bedstraw	N		
<i>Gutierrezia sarothrae</i>	broom snakeweed, matchweed	N		
<i>Helianthus annuus</i>	common sunflower	N		
<i>Holosteum umbellatum</i>	jagged chickweed	I		
<i>Hordeum jubatum</i>	squirreltail barley	N		
<i>Hordeum vulgare</i>	barley	I		
<i>Juglans nigra</i>	black walnut	I		
<i>Juncus balticus</i>	baltic rush	N	<i>Juncus arcticus</i> ssp. <i>balticus</i>	
<i>Juncus effusus</i>	pasture rush, soft rush	N		
<i>Lactuca serriola</i>	prickly lettuce	I		Sherman County Noxious Weed, C List
<i>Lagophylla ramosissima</i>	slender hareleaf, common rabbitleaf	N		
<i>Lepidium perfoliatum</i>	clasping pepperweed	I		
<i>Leymus cinereus</i>	Great Basin wildrye	N	<i>Elymus cinereus</i>	
<i>Logfia arvensis</i>	field cottonrose, field filago	I		
<i>Lupinus leucophyllus</i>	velvet lupine, woolly-leaved lupine	N		
<i>Madia exigua</i>	little tarweed, threadstem madia	N		
<i>Madia gracilis</i>	slender tarweed, common tarweed	N		
<i>Matricaria discoidea</i>	pineapple weed	I	<i>Matricaria matricarioides</i>	
<i>Medicago sativa</i>	alfalfa, lucerne	I		
<i>Melilotus officinalis</i>	yellow sweetclover	I		
<i>Mentha canadensis</i>	corn mint, field mint	N		
<i>Microsteris gracilis</i>	annual phlox, slender phlox	N	<i>Phlox gracilis</i>	
<i>Onopordum acanthium</i>	Scotch thistle, cotton thistle	I		ODA and Sherman County Noxious Weed, B List
<i>Phalaris arundinacea</i>	reed canarygrass	I		
<i>Phragmites australis</i> ssp. <i>australis</i>	American common reed	N		
<i>Plantago major</i>	common plantain	I		
<i>Plantago patagonica</i>	Indian wheat, woolly plantain	N		
<i>Plectritus macrocera</i>	longspur white plectritis	N		
<i>Poa bulbosa</i>	bulbous bluegrass	I		
<i>Poa pratensis</i>	Kentucky bluegrass	I		
<i>Poa secunda</i> ssp. <i>juncifolia</i>	alkali bluegrass, big bluegrass, Nevada bluegrass	N	<i>Poa ampla</i>	
<i>Poa secunda</i> ssp. <i>secunda</i>	Sandberg's bluegrass, Canby's bluegrass, pine bluegrass	N		
<i>Polemonium micranthum</i>	annual Jacob's-ladder, annual polemonium	N		
<i>Polygonum aviculare</i>	prostrate knotweed, doorweed, knotgrass	I		
<i>Polypogon monspeliensis</i>	rabbitsfoot grass, annual beardgrass	I		
<i>Populus alba</i>	silverleaf poplar, white poplar	I		
<i>Populus nigra</i>	black poplar, Lombardy poplar	I		
<i>Prunus virginiana</i>	western chokecherry	N		
<i>Pseudoroegneria spicata</i>	bluebunch wheatgrass	N	<i>Agropyron spicatum</i>	
<i>Rhaponticum repens</i>	Russian knapweed	I	<i>Acroptilon repens</i>	ODA and Sherman County Noxious Weed, B List
<i>Robinia pseudoacacia</i>	black locust	I		
<i>Rumex crispus</i>	curly dock	I		
<i>Salix lasiandra</i>	Pacific willow	N		

Scientific Name	Common Name	Native or Introduced? ¹	Synonym	Notes
<i>Salsola tragus</i>	prickly Russian thistle, tumbleweed	I	<i>Salsola kali</i>	Sherman County Noxious Weed, C List
<i>Schoenoplectus acutus</i>	hardstem bulrush, tule	N		
<i>Secale cereale</i>	cereal rye, common rye, rye	I		Sherman County Noxious Weed, C List
<i>Sisymbrium altissimum</i>	Jim Hill mustard, tumble mustard	I		
<i>Solanum triflorum</i>	cut-leaved nightshade	I		
<i>Stephanomeria paniculata</i>	stiff-branched wirelettuce, tufted wirelettuce	N		
<i>Taraxacum officinale</i>	common dandelion	I		
<i>Thinopyrum intermedium</i>	intermediate wheatgrass	I	<i>Agropyron intermedium</i>	
<i>Toxicodendron diversilobum</i>	poison oak	N		
<i>Tragopogon dubius</i>	yellow salsify	I		
<i>Trifolium repens</i>	white clover, Dutch clover	I		
<i>Triticum aestivum</i>	wheat	I		
<i>Typha latifolia</i>	broad-leaf cattail, common cattail	N		
<i>Urtica dioica</i>	stinging nettle	N		
<i>Verbascum thapsus</i>	common mullein, cowboy toilet paper	I		
<i>Verbena bracteata</i>	big-bracted verbenam bracted vervain	N		
<i>Veronica americana</i>	American brooklime	N		
<i>Vulpia bromoides</i>	brome fescue, rattail fescue	I	<i>Festuca bromoides</i>	
<i>Vulpia myuros</i>	rattail fescue, rat-tail six-weeks grass	I	<i>Festuca myuros</i>	
Nomenclature and nativity follows Flora of the Pacific Northwest (Hitchcock and Cronquist 2018) and OregonFlora (https://oregonflora.org/)				
1. I=Introduced, N=Native, U=Unknown				

2024 Habitat Categorization and Special Status Wildlife Survey Report

2024 Habitat Categorization and Special Status Wildlife Survey Report

**Biglow Canyon Wind Farm
Request for Amendment 4
November 2024**

Prepared for

BRIGHT  **NIGHT**

BIGL bn, LLC

Prepared by



TETRA TECH

This page intentionally left blank

Table of Contents

1.0	Introduction	1
2.0	Methods.....	2
2.1	Survey Area.....	2
2.2	Habitat Categorization	2
2.3	Special Status Wildlife	4
3.0	Results	4
3.1	Habitat.....	4
3.1.1	Category 6	5
3.1.2	Category 4	5
3.1.3	Category 3	6
3.1.4	Category 2	6
3.2	Special Status Wildlife Species.....	6
4.0	Summary and Discussion.....	6
5.0	References.....	7

List of Tables

Table 1. Habitat Categorization Types.....	3
Table 2. Habitat Categories, Types, and Subtypes within the Survey Area	5

List of Figures

- Figure 1. Project Location
- Figure 2. Habitat Types and Subtypes within the Survey Area
- Figure 3. Habitat Categories within the Survey Area
- Figure 4. Special Status Wildlife Species Observations

List of Attachments

- Attachment 1. Special Status Wildlife and Plant Species with Potential to Occur
- Attachment 2. Habitat Types and Subtypes Occurring at the Solar Components
- Attachment 3. Habitat Form
- Attachment 4. Select Photographs of Habitat and Wildlife
- Attachment 5. Wildlife Species Occurring in the Solar Components Vicinity

Acronyms and Abbreviations

AC	alternating current
BCWF or Existing Facility	Biglow Canyon Wind Farm
BIGL	BIGL bn, LLC
Certificate Holder / PGE	Portland General Electric Company
Council / EFSC	Oregon Energy Facility Siting Council
GIS	geographic information system
IPaC	Information for Planning and Consultation
MW	megawatt
OAR	Oregon Administrative Rule
ODFW	Oregon Department of Fish and Wildlife
ORBIC	Oregon Biodiversity Information Center
RFA	Request for Amendment
Site Certificate	Site Certificate to Amendment 3
Solar Components	photovoltaic solar energy generation and battery storage
Survey Area	2024 Habitat Categorization Survey Area
Tetra Tech	Tetra Tech, Inc.
U.S.C.	United States Code
USFWS	U.S. Fish and Wildlife Service

1.0 Introduction

The Portland General Electric Company (PGE or Certificate Holder) is submitting a Request for Amendment (RFA) 4 to the Site Certificate on Amendment 3, issued October 31, 2008 (Site Certificate) for the Biglow Canyon Wind Farm (BCWF or Existing Facility) to add photovoltaic solar energy generation and battery storage (Solar Components) to the operating BCWF in Sherman County, OR.

BCWF, owned and operated by PGE, is located within an approved site boundary comprising approximately 25,000 acres, approximately 2.5 miles northeast of the town of Wasco in Sherman County, Oregon (Figure 1). The BCWF operates under the Site Certificate from the Oregon Energy Facility Siting Council (Council or EFSC) as administered by the Oregon Department of Energy. BCWF currently consists of 217 wind turbines, with a maximum blade tip height of 445 feet, and a peak generating capacity of 450 megawatts (MW).

In RFA 4, PGE proposes to add up to 385 MW alternating current (AC) generating capacity from photovoltaic solar arrays and 375 MW in battery storage capacity. RFA 4 seeks to expand the BCWF site boundary to include the Solar Components in portions of the existing site boundary and in the proposed expanded site boundary (together, Solar Micrositing Area or RFA 4 Site Boundary).

The Solar Micrositing Area is approximately 3,980 acres and provides a conservative estimate of the maximum area needed for development, micrositing, and temporary disturbances from the Solar Components during construction, rather than the anticipated disturbance footprint. Solar Components will include solar arrays, inverters, battery energy storage system facilities and their subcomponents (i.e., inverters), two collector substations, a total of approximately 3 miles of 230-kilovolt generation tie transmission line, medium voltage collector lines, operations and maintenance structures, site access roads, internal roads, perimeter fencing, facility entry gates, and temporary laydown areas. The maximum generating capacity from the Solar Components will be 385 MW AC and construction may take place in phases.

PGE will own and operate the Solar Components as a part of the BCWF, which, to date, have been developed by BIGL bn, LLC (BIGL). BIGL, in its capacity as the project developer, supports PGE in this RFA 4 and may construct and temporarily operate the Solar Components on behalf of PGE under a Build-Transfer Agreement. Tetra Tech Inc. (Tetra Tech) is providing support to PGE and BIGL through preparation of RFA 4.

This survey report presents the methods and results of the 2024 habitat categorization and special status wildlife surveys conducted by Tetra Tech for the Solar Micrositing Area. The objective of this survey was to map and classify habitat according to Oregon Department of Fish and Wildlife (ODFW) guidelines set forth in Oregon Administrative Rule (OAR) 635-415-0025 and identify the presence of special status wildlife species and their habitats.

2.0 Methods

Prior to conducting field surveys, Tetra Tech conducted a desktop assessment including a review of existing information regarding habitat types and special status species with the potential to occur within the Solar Micrositing Area. The term “special status species” includes federal and state endangered, threatened, proposed, and candidate fish, wildlife, and plant species, as well as state sensitive wildlife species with distribution in Sherman County. Special status species also include eagles. Attachment 1 includes a list of special status species with the potential to occur in the 2024 Habitat Categorization Survey Area (Survey Area; defined in Section 2.1 below). This report focuses on special status wildlife species; special status plant species are addressed in a separate report. Nests of raptor species not considered special status wildlife are addressed in a separate report.

Information pertaining to federally listed species was obtained from the U.S. Fish and Wildlife Service (USFWS) Information for Planning, and Consultation (IPaC; USFWS 2024) database and site-specific records of sensitive species and habitat occurrences was received from the Oregon Biodiversity Information Center (ORBIC; ORBIC 2024). Other resources consulted during the desktop assessment included the National Land Cover Database, aerial photography, the original application materials for the BCWF, the Oregon Conservation Strategy (2016), and the ODFW Compass mapping tool. The Sherman County Comprehensive Plan (Goal 5) was also reviewed to determine the potential for big game and threatened and endangered species habitat, and eagle or other sensitive bird nesting habitats to occur within the Solar Micrositing Area.

2.1 Survey Area

Figure 1 shows the Solar Micrositing Area of approximately 3,980 acres and the slightly larger Survey Area. The Survey Area covered 4,636 acres including the entire Solar Micrositing Area and some additional lands that have since been omitted from the Solar Micrositing Area. The southern portion of the Survey Area is approximately 1.5 miles east of the town of Wasco, and the northern portion of the Survey Area is approximately 1 mile southwest of the John Day River, in Sherman County, Oregon (Figure 1).

2.2 Habitat Categorization

The objective of conducting surveys was to categorize habitats per the habitat categories set forth in OAR 635-415-0025 (ODFW’s Fish and Wildlife Mitigation Policy) and assign habitat subtypes consistent with the habitat subtypes developed during permitting of the BCWF.

The ODFW Fish and Wildlife Habitat Mitigation Policy provides a framework to categorize habitats based on type, quality, availability, and usefulness/importance to wildlife, and establishes mitigation goals and implementation standards for each. Table 1 defines each of the six habitat category types as presented in the ODFW Habitat and Wildlife Mitigation Policy.

Table 1. Habitat Categorization Types

Category Type	Definition¹	Mitigation Goal
1	Irreplaceable, essential habitat for a fish or wildlife species, population, or a unique assemblage of species and is limited on either a physiographic province or site-specific basis, depending on the individual species, population or unique assemblage.	The mitigation goal for Category 1 habitat is no loss of either habitat quantity or quality.
2	Essential habitat for a fish or wildlife species, population, or unique assemblage of species and is limited either on a physiographic province or site-specific basis depending on the individual species, population or unique assemblage.	The mitigation goal if impacts are unavoidable is no net loss of either habitat quantity or quality and to provide a net benefit of habitat quantity or quality.
3	Essential habitat for fish and wildlife, or important habitat for fish and wildlife that is limited either on a physiographic province or site-specific basis, depending on the individual species or population.	The mitigation goal is no net loss of either habitat quantity or quality.
4	Important habitat for fish and wildlife species.	The mitigation goal is no net loss of either habitat quantity or quality.
5	Habitat for fish and wildlife having high potential to become either essential or important habitat.	The mitigation goal, if impacts are unavoidable, is to provide a net benefit in habitat quantity or quality.
6	Habitat that has low potential to become essential or important habitat for fish and wildlife.	The mitigation goal is to minimize impacts.
1. Source: OAR 635-415-0025.		

Attachment 2 includes a habitat matrix that defines the habitat subtypes and the corresponding habitat categories in which each habitat subtype may fall based on proximity to wildlife resources and/or vegetation composition.

Habitat categorization surveys were conducted within the Survey Area concurrently with special status wildlife surveys. Habitat mapping was performed within a minimum of 1 acre mapping unit (except for specialized habitat such as cliffs or rock outcrops). These surveys were conducted by one field biologist familiar with vegetation of the Columbia Plateau Ecoregion and were scheduled during the growing season to aid in identification of vegetative species.

During surveys, meandering transects were walked by a biologist to categorize the composition and structure of habitat within the Survey Area by noting the dominant vegetation, presence of large trees, existing disturbance, or other habitat features such as shrub and grass composition and density. Each polygon was then assigned a habitat type, subtype, and habitat quality category that was guided by the habitat categorization for the BCWF. Habitat was classified into categories by quality per the ODFW Habitat Mitigation Policy. Spatial information was recorded on Samsung Galaxy tablets using the FieldMaps application; Attachment 3 includes the habitat form that was filled out during surveys to describe habitat category, type, and subtype. The biologist reviewed

high-resolution aerial photographs to locate and visit areas with unique vegetation or habitat features. Habitat types and categories were not assigned to wetlands or waters in the field as they were derived from data collected during wetland and water surveys.

Following the habitat categorization surveys, the digitized habitat boundaries were downloaded and processed in geographic information system (GIS) software, and the data were incorporated into spatial data. Data were then reviewed for quality control and assurance.

2.3 Special Status Wildlife

The objective of the wildlife survey was to document the presence of any special status species within the Survey Area, as well as general wildlife species that may occur. Special status species include species protected by the Endangered Species Act (7 United States Code [U.S.C.] 136), the Bald and Golden Eagle Protection Act (16 U.S.C. 668–668d), the Migratory Bird Treaty Act (16 U.S.C. 703-712), and ODFW (2021b).

Wildlife surveys were conducted by a field biologist familiar with fish and wildlife species of the Columbia Plateau Region and were scheduled to begin early in the morning and continue through late afternoon to allow observations of species most active at dawn and late afternoon (e.g., mule deer [*Odocoileus hemionus*]). Additionally, survey dates were planned to coincide with the period of highest biological activity of neotropical migrant and breeding birds, foraging and breeding wildlife species, and other taxa. During surveys, the biologist walked meandering transects through non-agricultural areas within the Survey Area, searching (scanning and listening) for wildlife species and recognizable signs of wildlife (e.g., scat, tracks, burrows and nests). Agricultural areas were surveyed by visually scanning and listening for wildlife from accessible roads or bordering non-agricultural lands. During surveys, the biologist documented the location of special status species, or recognizable sign, and recorded the number of individuals and their behavior, if observed. Special habitats and unique features (i.e., cliffs, ponds, talus slopes, rock outcrops, raptor nests, and big game) were also documented if encountered.

Following field surveys, the digitized data were downloaded and processed using GIS software and reviewed for quality control and assurance.

3.0 Results

3.1 Habitat

Targeted, field-based habitat categorization surveys were performed from June 10-12, 2024, within the Survey Area concurrently with special status wildlife surveys. These surveys determined habitat types and subtypes as shown in Table 2 and displayed on Figure 2. Land use in the Survey Area consists of existing wind turbines and associated gravel turbine pads and access roads, dryland wheat fields, grassland, and ephemeral drainages (Table 2). Photos of select habitat types and subtypes are provided in Attachment 4.

Table 2. Habitat Categories, Types, and Subtypes within the Survey Area

Habitat Category	Habitat Type	Habitat Subtype	Acreage within Survey Area	Category Subtotal
Category 2	Riparian	Intermittent Streams/Riparian Trees	11.9	11.9
Category 3	Upland	Upland Trees	20.6	70.0
	Shrub-steppe	Sagebrush Shrub-steppe	49.4	
Category 4	Grassland-steppe	Grasslands	94.5	94.5
Category 6	Agriculture	Irrigated Cropland	53.1	4,459.4
		Non-irrigated Cropland	4,343.2	
	Developed		63.3	
Total			4,635.9	4,635.9
Note: Totals may not sum exactly due to rounding.				

The most prominent habitat category in the Survey Area was Category 6, Agriculture habitat type, which included both Irrigated and Non-irrigated Croplands (4,396 acres), and Developed habitat subtypes (63 acres), totaling 96 percent of the Survey Area. The remaining 4 percent contained Category 4 Grasslands habitat subtype (95 acres), Category 3 Sagebrush Shrub-steppe (49 acres) and Upland Trees (21 acres) habitat subtypes, as well as Category 2 Intermittent and Riparian Tree habitat subtype (12 acres; Table 2; Figures 2 and 3).

3.1.1 Category 6

The Category 6 habitat is dominated by the Agriculture habitat type, which primarily contains the habitat subtype of Non-irrigated Croplands as well as one small area of Irrigated Cropland. The Non-irrigated Croplands consisted of 4,343 acres of active dryland wheat and barley fields, as well as fallow fields (Photos 1 and 3 in Attachment 4). The Irrigated Croplands consisted of one fallow and one active barley field, totaling 53 acres (Photo 4 in Attachment 4).

Additionally, approximately 63 acres of Category 6 habitat was classified as Developed, consisting of residences, farm buildings, farm equipment storage and staging areas, farming materials, corrals, turbine strings, paved and gravel county roads, pastures, abandoned buildings, a substation and transmission line, and a rock quarry (Photos 5 to 7 in Attachment 4).

3.1.2 Category 4

The Category 4 Grasslands-steppe contained one habitat subtype: Grasslands. This habitat consisted of a high percentage of non-native plant species. The dominant plants in this habitat were cereal rye (*Secale cereale*) and downy brome (*Bromus tectorum*), with some traces of native green rabbitbrush (*Chrysothamnus viscidiflorus*), balsamroot (*Asteraceae* spp.), wild buckwheat (*Eriogonum ovalifolium*), squirreltail (*Elymus elymoides*), and tall wheatgrass (*Thinopyrum elongatum*; Photos 8 and 9 in Attachment 4). Category 4 habitat is defined as “important habitat.”

3.1.3 Category 3

The Category 3 habitats contained two habitat types, Shrub-steppe and Upland. Both habitat types contain one subtype each: Sagebrush Shrub-steppe and Upland Trees. The Sagebrush Shrub-steppe subtype consisted of moderately disturbed areas with a mix of natives and non-natives and a minor to moderate shrub component. This habitat was dominated by downy brome, followed by basin big sagebrush (*Artemisia tridentata*) and grey rabbitbrush (*Ericameria nauseosa*). One mapped area was dominated by an unidentified annual grass, as well as bluebunch wheatgrass (*Pseudoroegneria spicata*), mature grey rabbitbrush, six-week fescue (*Vulpia octoflora*), sedum (*Crassulaceae* spp.), bulbous bluegrass (*Poa bulbosa*), and sagebrush (*Artemisia* spp.) on the hill tops (Photos 10 and 11 in Attachment 4).

The Upland Trees subtype consisted mostly of live black locust (*Robinia pseudoacacia*) trees and snags containing an understory of downy brome, squirreltail, tall wheatgrass, sagebrush, and rabbitbrush (Photo 12 in Attachment 4).

3.1.4 Category 2

Category 2 Riparian contained a habitat subtype of Intermittent Streams/Riparian Trees and consisted of cattails (*Typhalatifolia*), reed canarygrass (*Phalaris arundinacea*), cereal rye, willow (*Salix exigua*), Russian olive (*Elaeagnus angustifolia*), cottonwood (*Populus trichocarpa*), stinging nettle (*Urtica dioica*) and poison hemlock (*Conium maculatum*) in varying densities (Photos 16 and 17 in Attachment 4).

3.2 Special Status Wildlife Species

Concurrently with habitat surveys on June 10-12, 2024, the biologist documented the occurrence of wildlife species and maintained a running list of wildlife individuals and sign observed in the vicinity of the Solar Components (Attachment 5).

Review of IPaC and ORBIC data returned no occurrences of special status wildlife within the Survey Area. No federal or state endangered, threatened, proposed, or candidate species were observed during the survey. One Oregon special status species, a loggerhead shrike (*Lanius ludovicianus*, state-sensitive) was observed within the Survey Area (Figure 4). According to Holmes and Thibodeaux (2014), the loggerhead shrike breeds in a variety of shrubland and open woodland habitats, but relies on big sagebrush plant communities and breeds primarily in this habitat, which is threatened by high-intensity fires and conversion to agriculture.

4.0 Summary and Discussion

The habitat in the Survey Area was primarily classified as Category 6 Agriculture and Developed lands. The uncultivated and undeveloped portions of the Survey Area were classified as Category 4 Grasslands, Category 3 Sagebrush Shrub-steppe and Upland Trees habitats, or Category 2

Intermittent Streams/Riparian Trees habitat. The Category 2, 3, and 4 habitats provide essential and important wildlife habitat that are limited within the Survey Area.

The mitigation goal for Category 2 habitat, if impacts are unavoidable, is no net loss of either habitat quantity or quality and to provide a net benefit of habitat quantity or quality. The mitigation goal for Category 3 habitat is no net loss of either habitat quantity or quality. The mitigation goal for Category 4 habitat is no net loss of habitat quantity or quality and the mitigation strategy is in-kind, in-proximity mitigation. The mitigation goal for Category 6 habitat is to minimize impact and the mitigation strategy is to minimize direct habitat loss and avoid off-site impacts. Mitigation is typically not required for impacts to Category 6 habitat.

One special status species, the loggerhead shrike, was documented during surveys. This species is likely to use the Shrub-steppe habitat within the Survey Area during breeding and/or migration. No federal or state threatened, endangered, or candidate species were observed in the Survey Area.

5.0 References


- Holmes, A.L., and B. Thibodeaux. 2014. Population Status, Distribution, and Nesting Success of Loggerhead Shrikes in the Boardman Conservation Area, Boardman, Oregon, 2013. Prepared for The Nature Conservancy, Portland, Oregon by Northwest Wildlife Science, LLC. Available online at: https://www.conservationgateway.org/ConservationByGeography/NorthAmerica/UnitedStates/oregon/deserts/cbg/wildlife/Documents/LOSH_2013_Report_Final_editted_jr.pdf. Accessed July 2024.
- ODFW (Oregon Department of Fish and Wildlife). 2021a. Oregon Department of Fish and Wildlife Sensitive Species List. Available online at: https://www.dfw.state.or.us/wildlife/diversity/species/docs/Sensitive_Species_List.pdf. Accessed July 2024.
- ODFW. 2021b. Threatened, Endangered and Candidate Fish and Wildlife Species, Revised October 2021. Available online at: https://www.dfw.state.or.us/wildlife/diversity/species/docs/Threatened_and_Endangered_Species.pdf. Accessed July 2024.
- ORBIC (Oregon Biodiversity Information Center). 2024. Element Occurrence Record Digital Data Set for rare, threatened, or endangered species for the state of Oregon, May 2, 2024. ORBIC, Institute for Natural Resources. Portland State University, Portland, Oregon. Accessed July 2024.
- USFWS (U.S. Fish and Wildlife Service). 2024. IPaC – Information for Planning and Consultation Resource List for the Project location in Sherman County, Oregon. Available online at: <https://ipac.ecosphere.fws.gov/> Accessed July 2024.

This page intentionally left blank.

Figures

This page intentionally left blank

\\css706g\ists\ICES\Projects\BrightNight\BiglowCanyon\Maps\HabitatWildlifeReport_20240701\BrightNight_BiglowSolar_HabitatWildlife_20241212.aprx



Biglow Canyon Wind Farm Request for Amendment 4


**Figure 1
Project Location**

SHERMAN COUNTY, OR

- Amended Site Boundary
 - Survey Area
 - Proposed Transmission Interconnect
 - Existing Transmission Lines (230-kV)
 - Existing Biglow Canyon Substation
 - Existing Biglow Canyon Wind Farm Turbine
- Oregon Energy Facility Council Facilities (Operating)**
- Biglow Canyon Wind Farm
 - Golden Hills Wind Project
 - Klondike III Wind Project
- Land Status**
- Bureau of Land Management
 - Private



Reference Map




1:55,000

WGS 1984 UTM Zone 10N

0 0.5 1 2 Miles

NOT FOR CONSTRUCTION

\\cess706gists\1\CES\Projects\BrightNight\BiglowCanyon\Maps\HabitatWildlifeReport_20240701\BrightNight_BiglowSolar_HabitatWildlife_20241212.aprx



Biglow Canyon Wind Farm Request for Amendment 4

Figure 2
Habitat Types and
Subtypes within
the Survey Area

SHERMAN COUNTY, OR

- Survey Area
- US Highway
- State Highway
- Habitat Type; Subtype
 - Agriculture; Non-Irrigated Cropland
 - Agriculture; Irrigated Cropland
 - Grassland-steppe; Grasslands
 - Riparian; Intermittent Stream / Riparian Trees
 - Shrub-steppe; Sagebrush Shrub-Steppe
 - Upland; Upland Trees
 - Developed



Reference Map




1:44,000

WGS 1984 UTM Zone 10N



NOT FOR CONSTRUCTION

\\cess706gists\1\CES\Projects\PD\BrightNight\BiglowCanyon\Maps\Habitat\WildlifeReport_2024070\BrightNight_BiglowSolar_HabitatWildlife_20241212.aprx



Biglow Canyon Wind Farm Request for Amendment 4


Figure 3
Habitat Categories
within the Survey Area

SHERMAN COUNTY, OR

- Survey Area
- US Highway
- State Highway
- Habitat Category
 - 2
 - 3
 - 4
 - 6



Reference Map




1:44,000

WGS 1984 UTM Zone 10N

0 0.5 1 2 Miles

NOT FOR CONSTRUCTION

\\cess706gists\1\ICES\Projects\PD\BrightNight\BiglowCanyon\Maps\Habitat\WildlifeReport_2024070\BrightNight_BiglowSolar_HabitatWildlife_20241212.aprx



Biglow Canyon Wind Farm Request for Amendment 4


Figure 4
Special Status Wildlife
Species Observations

SHERMAN COUNTY, OR

-  Survey Area
-  US Highway
-  State Highway
- Special Status Wildlife Observation
-  Loggerhead Shrike



Reference Map



1:44,000

WGS 1984 UTM Zone 10N

0 0.5 1 2 Miles

NOT FOR CONSTRUCTION

Attachment 1. Special Status Wildlife and Plant Species with Potential to Occur

This page intentionally left blank

Attachment 1: Special Status Wildlife and Plant Species with Potential to Occur

Common Name	Scientific Name	Federal Status ¹	Oregon Status ²
Birds			
Brewer's sparrow	<i>Spizella breweri</i>	-	S, CSS
Burrowing owl (Western)	<i>Athene cunicularia hypugaea</i>	SOC	SC, CSS
Common nighthawk	<i>Chordeiles minor</i>	-	S, CSS
Ferruginous hawk	<i>Buteo regalis</i>	SOC	SC, CSS
Golden eagle	<i>Aquila chrysaetos</i>	BGEPA	-
Grasshopper sparrow	<i>Ammodramus savannarum</i>	-	S, CSS
Bald eagle	<i>Haliaeetus leucocephalus</i>	BGEPA	-
Lewis's woodpecker	<i>Melanerpes lewis</i>	BCC	SC, CSS
Loggerhead shrike	<i>Lanius ludovicianus</i>	-	S, CSS
Long-billed curlew	<i>Numenius americanus</i>	-	SC, CSS
Swainson's hawk	<i>Buteo swainsoni</i>	-	S, CSS
Reptiles			
Northern sagebrush lizard	<i>Sceloporus graciosus</i>	-	S, CSS
Western painted turtle	<i>Chrysemys picta bellii</i>	-	SC, CSS
Mammals			
Hoary bat	<i>Lasiurus cinereus</i>	-	S
Pallid bat	<i>Antrozous pallidus</i>	-	S
Silver-haired bat	<i>Lasionycteris noctivagans</i>	-	S, CSS
Spotted bat	<i>Euderma maculatum</i>	-	S, CSS
Townsend's big-eared bat	<i>Corynorhinus townsendii</i>	-	SC, CSS
Plants			
Henderson's ricegrass	<i>Eriocoma [Achnatherum] hendersonii</i>	-	C
Hepatic monkeyflower	<i>Erythranthe jungermannioides</i>	-	C
Lawrence's milkvetch	<i>Astragalus collinus var. laurentii</i>	-	T
Sessile mousetail	<i>Myosurus sessilis</i>	-	C
<p>Sources: ODFW 2021a, ODFW 2021b, ORBIC 2024, USFWS 2024</p> <p>1. Federal Status: BGEPA = Bald and Golden Eagle Protection Act, BCC = Bird of Conservation Concern, SOC = Species of Concern, C = Candidate, T = Threatened, PT = Proposed as Threatened.</p> <p>2. Oregon Department of Fish and Wildlife Status in the Columbia Plateau/Columbia Basin: CSS = Conservation Strategy Species, SC = Sensitive Critical, S = Sensitive. Oregon Department of Agriculture Plant listing, T = Threatened, C = Candidate.</p>			


This page intentionally left blank

Attachment 2. Habitat Types and Subtypes Occurring at the Solar Components

This page intentionally left blank

Habitat Type	Habitat Sub-type	Category 1	Category 2	Category 3	Category 4	Category 5	Category 6
Riparian	Intermittent Streams			Essential or important fish and wildlife habitat which is limited.			
	Riparian Trees		Essential and limited habitat for wildlife (documented nest/roost habitat)				
	Intermittent Streams / Riparian Trees		Essential and limited habitat for wildlife (documented nest/roost habitat)				
Surface Water	Ponds			Essential or important wildlife habitat which is limited (e.g. wetland feature).			
Upland	Upland Trees	Irreplaceable, essential habitat for a wildlife species (e.g. Swainson's hawk) and limited within a physiogeographic province (documented food/cover/nest habitat, and active nests).		Essential or important habitat for wildlife that is limited.			
Shrub-steppe	Sagebrush Shrub-Steppe		Essential and limited wildlife habitat (e.g. fairly undisturbed, old-growth shrub structure; moderate grazing).	Essential or important wildlife habitat which is limited (e.g. fairly undisturbed habitat; moderate grazing).	Important wildlife habitat (e.g. moderate-heavy grazing or weedy habitat.		
Grassland-steppe	Grasslands				Important wildlife habitat (e.g. moderate-heavy grazing or weedy habitat.		
Agriculture	Irrigated Cropland						Cultivated cropland with the artificial application of water with low potential for becoming essential or important habitat.
	Non-Irrigated Cropland						Cultivated croplands with low potential for becoming essential or important habitat.
	Conservation Reserve Program (CRP)			Croplands planted to grassland / shrub-steppe in the CRP program that provide important wildlife habitat.	Croplands planted to grassland / shrub-steppe in the CRP program that lack seral stage vegetative communities and/or are of less importance as wildlife habitat because of the land management or topographic locale.		
Developed							Low potential for becoming essential or important habitat (e.g. residences, storage bins, farm equipment storage, grain elevators, industrial/commercial facilities, gravel quarries).

		Category 2	Category 3	Category 4	Category 5
Eastside Grassland	Native % (Without Sage)	> 75 %	50-75 %	15-50 %	< 15 %
	Native % (With Sage)	50-75 %	15-50 %	< 15 %	None
	Disturbance	None – Low	Moderate	High	High - Extreme



Fish and Wildlife Habitat Mitigation - Key Definitions

Essential habitat	Any habitat condition or set of habitat conditions which, if diminished in quality or quantity, would result in depletion of a fish or wildlife species.
Limited habitat	An amount insufficient or barely sufficient to sustain fish and wildlife populations over time.
Important habitat	Any habitat recognized as a contributor to sustaining fish and wildlife populations over time.
Irreplaceable habitat	Successful in-kind habitat mitigation to replace lost habitat quantity and/or quality is not feasible within an acceptable period of time or location, or involves an unacceptable level of risk or uncertainty.
Habitat with High Restoration Potential	Previous uses or activities that have reduced habitat values need to be able to be eliminated or severely reduced.

The Fish and Wildlife Habitat Mitigation Policy establishes mitigation goals for each category of habitat and, depending upon the importance of the habitat, identifies preferred strategies to avoid or mitigate the impact of proposed actions on fish and wildlife habitat. The policy sets sideboards within which ODFW considers recommended options and alternatives for mitigation. The less important the habitat is, the more options that may be considered for mitigation.

Habitat Categories and Mitigation Strategies

Habitat Category	Definition	Example	Goal for Mitigation	Mitigation Strategy
Category 1	Irreplaceable, essential and limited habitat	Bogs and fens, certain springs and pools	No loss of habitat quantity or quality	Avoidance
Category 2	Essential and limited habitat	Salt marshes, cottonwood galleries, big game winter range, salmonid migration corridors, some spawning and rearing areas.	No net loss of habitat quantity or quality and to provide a net benefit of habitat quantity or quality	In-kind, in-proximity mitigation
Category 3	Essential habitat, or important and limited habitat	Older forested areas, reed canary grass wetland, spawning and rearing areas.	No net loss of habitat quantity or quality	In-kind, in-proximity mitigation
Category 4	Important habitat	Isolated or degraded wetlands, big game summer range, spawning, rearing and foraging areas.	No net loss of habitat quantity or quality	In-kind or out-of-kind, in-proximity or off-proximity mitigation
Category 5	Habitat having high potential to become either essential or important habitat	Restorable rye grass fields or diked or drained coastal marshes, marshes, some types of reservoirs.	Net benefit in habitat quantity or quality	Actions that improve habitat conditions
Category 6	Habitat that has low potential to become essential or important habitat	Urban areas and other areas with little or no restoration potential, artificial ponds without native species.	Minimize impacts	Minimize direct habitat loss and avoid off-site impacts

Source: OAR 635-415-0025

Attachment 3. Oregon Habitat Categorization Datasheet

This page intentionally left blank

Oregon Habitat Categorization Datasheet

Data Sheet ID #: _____

Project: _____ County: _____ Surveyor(s): _____ Date: _____

Habitat Category: 1 / 2 / 3 / 4 / 5 / 6

Habitat Type & Subtype

Riparian	Surface Water	Upland	Shrub-steppe	Agriculture <i>*specify crops in "Description"</i>
☐ Intermittent Stream ☐ Riparian Trees ☐ Intermittent Stream / Riparian Trees ☐ _____	☐ Ponds ☐ _____	☐ Upland Trees ☐ _____	☐ Sagebrush Shrub-steppe ☐ _____	☐ Non-irrigated Cropland ☐ Irrigated Cropland ☐ Conservation Reserve Program (CRP) ☐ _____
		Grassland-steppe ☐ Grasslands ☐ _____		Developed <i>*specify features in "Description"</i> ☐ Developed ☐ _____

Datasheet Description: _____**Additional Notes:** _____

Overall Habitat Community *(Total cover amongst all strata, can equal over 100%)*

% Absolute Canopy Closure: Trees: _____ Shrubs: _____ Grasses + Forbs: _____

% Bare Ground: _____ % Duff: _____ % Rock: _____ % Moss: _____ Cryptobiotic Crust: ☐ Yes ☐ No

Species Composition *(Mark natives with *, noxious weeds with **)*

Dominant ≥20%, Subdominant 10-20% of stratum cover

TREES

Dominant: _____

Subdominant: _____

Trace / Other noteworthy species: _____

Subcanopy Layers: _____ Percent of Stratum that is Native: _____ Max Vegetation Height (ft.): _____

Avg. DBH (in.): _____ Stumps Present? ☐ Yes ☐ No Forest Phase: GF / SHR / OSP / CSPP / LGSAW / OGDDSnags Present? ☐ Yes ☐ No Abundance: ____/ac Snag Stage: 1 / 2 / 3 / 4 / 5

Notes: _____

SHRUBS

Dominant: _____

Subdominant: _____

Trace / Other noteworthy species: _____

Subcanopy Layers: _____ Percent of Stratum that is Native: _____ Max Vegetation Height (in.): _____

Notes: _____

GRASSES + FORBS

Dominant: _____

Subdominant: _____

Trace / Other noteworthy species: _____

Subcanopy Layers: _____ Percent of Stratum that is Native: _____ Max Vegetation Height (in.): _____

Notes: _____

Other Descriptions*Check all that apply within or outside but in view of the habitat polygon(s).**List the approximate distance from polygon edge, and specify units (ft, m, mi), or write "in" for features within the polygon(s).*

Disturbance		Sensitive Species/Habitat	Unique Features	Anticipated Recovery Timeline*
☐ None	☐ Row Crop _____	☐ None	☐ None	☐ 0-5 years
☐ Wind Facility _____	☐ Farm _____	☐ Bald Eagle _____	☐ Caves _____	☐ 5-30 years
☐ Solar Facility _____	☐ Corral _____	☐ Golden Eagle _____	☐ Mines / Quarries _____	☐ >30 years
☐ Railroad _____	☐ Pasture _____	☐ Raptor Nest(s) _____	☐ Cliffs / Rimrock _____	
☐ Dirt Road _____	☐ Grazing _____	☐ _____	☐ Rock Outcrop _____	Recovery Rationale:
☐ Gravel Road _____	☐ Erosion _____	☐ _____	☐ Abandoned Buildings _____	☐ Mature Sagebrush
☐ Paved Road _____	☐ Construction _____	☐ _____	☐ Large Snag(s) _____	☐ Mature Riparian
☐ Highway _____	☐ Invasive Plants _____	☐ _____	☐ Wood Bridge(s) _____	☐ _____
☐ Industrial _____	☐ Logging _____	☐ _____	☐ Balds _____	☐ _____
☐ Residence _____	☐ Thinning _____	☐ _____	☐ Bluffs _____	☐ _____
☐ Campground _____	☐ Quarry _____	☐ _____	☐ Wetlands / Waters _____	☐ _____
☐ _____	☐ _____	☐ _____	☐ _____	

*Anticipated recovery timeline should be estimated during surveys for the purposes of determining the need for mitigation to compensate for temporary impacts to habitat that would be long-term or difficult to restore.

For example, disturbed annual grassland typically recovers in 2-5 years, high quality native grassland typically recovers in ~3 years, or > 5 years depending on the site (e.g., existing disturbance, precipitation, and quality of planting/seeding), rabbitbrush typically recovers <5 years, and sagebrush typically recovers in 5-30 years.

See Brown 1985 for forest recovery/phase timelines.

Additional Notes / Categorization Rationale:

Attachment 4. Select Photographs of Habitat and Wildlife

This page intentionally left blank



Photo 1. Category 6: Non-irrigated cropland with active wheatfield.



Photo 3. Category 6: Non-irrigated cropland with fallow field.



Photo 4. Category 6: Irrigated cropland with fallow field.



Photo 5. Category 6: Development with farm building and equipment/materials storage area.



Photo 6. Category 6: Development with abandoned buildings.



Photo 7. Category 6: Development with substation and large transmission line.



Photo 8. Category 4: Grassland-steppe habitat dominated by cereal rye and downy brome.



Photo 9. Category 4: Grassland-steppe habitat dominated by cereal rye, downy brome, and traces of native shrubs and grasses.



Photo 10. Category 3: Moderately disturbed shrub-steppe habitat dominated by downy brome, with some rabbitbrush and sagebrush.



Photo 11. Category 3: Shrub-steppe habitat dominated by annual grasses, native grasses such as six-week fescue and some sagebrush on hill tops.



Photo 12. Category 3: Upland habitat with black locust trees and snags.



Photo 13. Category 3: Upland habitat with black locust trees and snags with raptor nest (great-horned owl).



Photo 14. Category 3: Upland habitat with black locust trees and snags without raptor nests.



Photo 15. Category 3: Upland habitat showing wildlife guzzler location.



Photo 16. Category 2: Riparian habitat with cattails, reed canarygrass, cereal rye, and other riparian vegetation.



Photo 17. Category 2: Riparian habitat within drainage.

Attachment 5. Wildlife Species Occurring in the Solar Components Vicinity

This page intentionally left blank

Attachment 5: Wildlife Species Occurring in the Solar Components Vicinity

Common Name*	Scientific Name
Birds	
American crow	<i>Corvus brachyrhynchos</i>
American goldfinch	<i>Spinus tristis</i>
Barn swallow	<i>Hirundo rustica</i>
California quail	<i>Callipepla californica</i>
Common raven	<i>Corvus corax</i>
European starling	<i>Sturnus vulgaris</i>
Great horned owl	<i>Bubo virginianus</i>
Horned lark	<i>Eremophila alpestris</i>
House wren	<i>Troglodytes aedon</i>
Loggerhead shrike*	<i>Lanius ludovicianus</i>
Mourning dove	<i>Zenaida macroura</i>
Red-tailed hawk	<i>Buteo jamaicensis</i>
Red-winged blackbird	<i>Agelaius phoeniceus</i>
Ring-necked pheasant	<i>Phasianus colchicus</i>
Song sparrow	<i>Melospiza melodia</i>
Turkey vulture	<i>Cathartes aura</i>
Western meadowlark	<i>Sturnella neglecta</i>
Western wood pewee	<i>Contopus sordidulus</i>
Mammals	
Grey squirrel	<i>Sciurus carolinensis</i>
Mountain cottontail	<i>Sylvilagus nuttallii</i>
Mule deer	<i>Odocoileus hemionus</i>
Porcupine	<i>Erethizon dorsatum</i>
Pronghorn	<i>Antilocapra americana</i>
Invertebrates	
Damselfly	NA
Dragonfly	NA
*ODFW Sensitive Species. See Attachment 1. NA = Not Applicable, unable to identify to species level.	

2024 Raptor Nest Survey Report

(**CONFIDENTIAL**)

This document contains confidential information and is provided under separate cover.

2026 Raptor Nest Survey Report

(**CONFIDENTIAL**)

This document contains confidential information and is provided under separate cover.

Attachment G-2. Draft Habitat Mitigation Plan

Biglow Solar Project Habitat Mitigation Plan

**Biglow Canyon Wind Farm
Request for Amendment 4 and 5
October 2026**

Prepared for



Portland General Electric Company

Prepared by



Tetra Tech

1750 S. Harbor Way, Suite 400
Portland, Oregon 97201

Table of Contents

1.0	Introduction	1
2.0	Finalization of the HMP (PRE-FW-01)	1
3.0	Habitat Categories and Habitat Types.....	3
4.0	Temporary and Permanent Impacts	4
5.0	Methods for Calculating Mitigation	5
6.0	Biglow Canyon Wind Farm Habitat Mitigation Area	7
6.1	Habitat Assessment and Mitigation Accounting.....	7
6.2	Habitat Enhancement Actions.....	8
6.2.1	Restoration and Enhancement Actions in the Event of Fire	8
6.2.2	Noxious Weed Prevention, Inventory, and Control.....	9
6.2.3	Access Control	9
6.2.4	Sagebrush Planting.....	10
6.2.5	Annual Grass Treatment.....	10
6.2.6	Enhancing Pollinator Habitat	10
7.0	HMA Enhancement Monitoring	11
7.1	Monitoring Location Establishment	11
7.2	Photo Points.....	12
7.3	Vegetative Cover.....	12
7.4	Density	12
7.5	Weed Monitoring.....	13
8.0	Success Criteria.....	13
9.0	Data Reporting and Timeline.....	14
10.0	Amendment of the HMP.....	15
11.0	References.....	15

List of Tables

Table 1. Habitat Categorization Types	3
Table 2. Anticipated Temporary and Permanent Impacts by Habitat Category and Habitat Subtype .	5
Table 3. Calculating Mitigation for Permanent Impacts.....	6
Table 4. Calculating Mitigation for Temporary Impacts.....	6
Table 5. Recommended Noxious Weed Control Methods	9

Table 6. Cover Types.....	12
Table 7. HMA Schedule of Implementation and Monitoring.....	14

List of Figures

Figure 1. Habitat Mitigation Area

[Figure 2. -Sagebrush Planting and Photopoint Locations](#)

List of Attachments

Attachment 1. Baseline Photo Monitoring

1.0 Introduction

This Habitat Mitigation Plan (HMP) was submitted by the Portland General Electric Company (PGE or Certificate Holder) for photovoltaic (PV) solar energy generation and battery storage (Solar Components) and a transmission line associated with the Biglow Canyon Wind Farm (BCWF) Solar Facility, which is under development in two areas connected by a generation-tie transmission line. BCWF, owned and operated by PGE, is located within an approved site boundary comprising approximately 25,000 acres, approximately 2.5 miles northeast of Wasco in Sherman County, Oregon. BCWF consists of 217 operating wind turbines with a peak generating capacity of 450 megawatts (MW).

The Oregon Energy Facility Siting Council (EFSC) issued the BCWF Fourth Amended Site Certificate for the Facility on March 27, 2026, to add 125 MW alternating current (AC) generating capacity from solar PV arrays and 500 megawatt-hours (MWh) in battery storage capacity (Request for Amendment [RFA] 4 Solar Facility) within the existing BCWF site boundary. Construction of the RFA 4 Solar Facility commenced in July 2026, with an anticipated commercial operation date of December 2027. BCWF RFA 5 proposes to add 200 MW AC generating capacity from solar PV arrays and 1,000 MWh in battery storage capacity (RFA 5 Solar Facility). The RFA 5 Solar Facility is located across approximately 622 acres of the existing BCWF site boundary and 839 acres of expanded site boundary. Construction of the RFA 5 Solar Facility is anticipated to commence in Q3 of 2027. PGE will own and operate the RFA 4 and RFA 5 Solar Facilities as a part of the BCWF, which to date have been developed by BIGL bn, LLC (BIGL or Project Developer). BIGL, in its capacity as the Project Developer, supports PGE in RFA 4 and RFA 5 to add solar PV energy generation and battery storage at the BCWF and may construct and temporarily operate the Solar Components on behalf of PGE under a Build-Transfer Agreement.

The Certificate Holder's goal is to preserve and maintain in-kind habitat in the Columbia Plateau Ecoregion to achieve a net benefit to Category 2 habitat, and no net loss of Category 4 habitat, through the measures described in this HMP.

2.0 Finalization of the HMP (PRE-FW-01)

Fish and Wildlife Habitat Condition PRE-FW-01 requires finalization of the draft HMP, stating the following:

Prior to construction of a phase or component of the solar facility, as applicable, the certificate holder shall submit a final Habitat Mitigation Plan substantially similar to Final Order on RFA4 Attachment F, based on the finalization requirements included in Section 2.0 of the plan. The final plan must be submitted for review and approval by the Department, in consultation with ODFW.

This HMP was finalized prior to construction of the RFA 4 Solar Facility and will be amended prior to construction of the RFA 5 Solar Facility, if necessary. The HMP was updated to include the estimated permanent habitat impacts associated with the facility, component or phase, as applicable. Final review and approval of the HMP was delegated by EFSC under Oregon Revised Statutes (ORS) 469.402 to the Oregon Department of Energy (ODOE). ODOE, in consultation with the Oregon Department of Fish and Wildlife (ODFW), will review the final HMP based on the following scope:

1. **Final habitat and impact assessment.** Based on final design/layout, update tabular summaries of temporary and permanent disturbance estimates (acres).
 - a. Table 2. Temporary and Permanent Impacts by Habitat Category and Habitat Subtype,
 - b. Table 3. Calculating Mitigation for Permanent Impacts; and
 - c. Table 4. Calculating Mitigation for Temporary Impacts.
2. **Execute legal instrument.** PGE will obtain and submit to ODOE a copy of the conservation easement or legal instrument securing the Habitat Mitigation Area (HMA) for conservation purposes as outlined in this HMP:
 - a. The term of the conservation easement or legal instrument will be for the life of the RFA 4 and RFA 5 Solar Facilities.
 - b. The terms of the conservation easement or legal instrument will be binding upon the landowner, heirs, or successors, and would run with the land if property is sold.
 - c. The conservation easement or legal instrument will identify restricted uses that have the potential to diminish wildlife habitat functions and value, including but not limited to, grazing; grading, mowing, blading or expansion of impervious surfaces or access road networks; division of the Property.
3. **Identify/map baseline condition/monitoring sites and enhancement action areas.** Map baseline habitat conditions of the HMA. Propose number and location of monitoring sites for noxious weeds and shrub-planting. Submit habitat mapping and monitoring site locations to ODOE and ODFW for review and approval.
 - a. Provide baseline habitat maps, photographs, and maps/GIS data for proposed monitoring sites at the HMA with a description of the weed and vegetative cover.
 - i. Baseline monitoring plot photo points within the HMA; include 1) a minimum of five permanent photo points distributed to show general vegetation status throughout the HMA, and 2) additional photo points as needed to monitor success of noxious weed control and shrub-planting;
4. **Finalize enhancement actions and success criteria in Section 6.2.1, 6.2.2, and 6.2.3.** Based on the baseline assessment completed under Sub(3) above:
 - a. Provide maps/GIS data of enhancement action locations.
 - b. Determine sagebrush planting density.
 - c. Finalize the success criteria for weed treatment and shrub planting.

- i. Success criteria for weed treatment must represent a percent increase in native vegetation cover and percent decrease in invasive cover at monitoring sites compared to baseline.
- ii. Shrub planting will be considered successful when sagebrush density in enhancement areas is ~~50 percent of planting density~~ 8 shrubs per acre at 5 years.
- d. Propose restoration methods in the event of fire, which at a minimum include post-burn seeding and noxious weed control.
- e. ODOE and ODFW must evaluate whether the combined enhancement actions satisfy the Category 2 mitigation goal of a net benefit in habitat quality and the Category 3 and 4 mitigation goal of a no net loss of habitat quantity or quality. If the HMA has minimal need or opportunity for the enhancement actions as currently proposed, additional actions may be identified/required.
- 5. **Finalize Implementation Schedule.** Provide an implementation schedule for habitat enhancement actions and monitoring.
- 6. **Finalize Monitoring Protocols and Survey Methods.** Submit final proposed monitoring protocol and survey methods to monitor and evaluate the success of enhancement actions.

Approval of the final HMP will be based on review by ODOE, in consultation with ODFW, based on the adequacy of information provided to address the steps/requirements listed above.

3.0 Habitat Categories and Habitat Types

A habitat survey was conducted in 2024 to verify habitat subtypes and habitat categories of all areas to be affected by the RFA 4 and 5 Solar Facilities. The survey area covered 4,636 acres, including the entire RFA 4 and 5 Solar Facilities and some additional lands that have since been omitted from consideration. The ODFW Fish and Wildlife Habitat Mitigation Policy (Oregon Administrative Rules [OAR] 635-415-0025) provides a framework to categorize habitats based on type, quality, availability, and usefulness/importance to wildlife, and establishes mitigation goals and implementation standards for each. Table 1 defines each of the six habitat category types as presented in the ODFW Habitat Mitigation Policy.

Table 1. Habitat Categorization Types

Category Type	Definition ¹	Mitigation Goal
1	Irreplaceable, essential habitat for a fish or wildlife species, population, or a unique assemblage of species and is limited on either a physiographic province or site-specific basis, depending on the individual species, population or unique assemblage.	The mitigation goal for Category 1 habitat is no loss of either habitat quantity or quality.

Category Type	Definition ¹	Mitigation Goal
2	Essential habitat for a fish or wildlife species, population, or unique assemblage of species and is limited either on a physiographic province or site-specific basis depending on the individual species, population, or unique assemblage.	The mitigation goal, if impacts are unavoidable, is no net loss of either habitat quantity or quality and to provide a net benefit of habitat quantity or quality.
3	Essential habitat for fish and wildlife, or important habitat for fish and wildlife that is limited either on a physiographic province or site-specific basis, depending on the individual species or population.	The mitigation goal is no net loss of either habitat quantity or quality.
4	Important habitat for fish and wildlife species.	The mitigation goal is no net loss of either habitat quantity or quality.
5	Habitat for fish and wildlife having high potential to become either essential or important habitat.	The mitigation goal, if impacts are unavoidable, is to provide a net benefit in habitat quantity or quality.
6	Habitat that has low potential to become essential or important habitat for fish and wildlife.	The mitigation goal is to minimize impacts.
1. Source: OAR 635-415-0025.		

4.0 Temporary and Permanent Impacts

Impacts may be permanent or temporary. Permanent impacts are defined as those impacts that will exist for the life of the RFA 4 and 5 Solar Facilities. Temporary impacts are those impacts that will last for a time less than the life of the RFA 4 and 5 Solar Facilities. The duration of temporary impacts to habitat will vary by habitat subtype. For example, the recovery period for agricultural areas that were temporarily disturbed could be as short as 1 to 3 years, ~~g.~~ Grasslands generally recover within 3 to 7 years, and shrublands may require 10 to 50 years to recover (with the longer recovery periods associated with disturbances in mature sagebrush [*Artemisia* sp.] habitats). The Certificate Holder will restore temporary impacts consistent with the Comprehensive Solar Revegetation and Soil Management Plan (see Exhibit C, Attachment C-2 of RFA 5).

As described in Exhibit P of RFA 4, the Final Order on RFA 4, and Exhibit G of RFA 5, there are no impacts to Category 1 habitat within the RFA 4 and 5 Solar Facilities. Category 2 habitat occurs in the Solar Array Area and will be impacted by the Solar Components (Table 2). Category 2 habitat within the RFA 4 Site Boundary is associated with Shrub-steppe: Sagebrush shrub-steppe habitat along Biglow Canyon and, to a lesser extent, Riparian: Riparian Tree habitat and Grassland-steppe: Grassland habitat. Category 3, 4, and 6 habitat will also be impacted by the Solar Facilities, while Category 5 habitat was not identified in the disturbance area. During a site visit on January 10, 2025, ODFW recommended that impacts to Upland Tree habitat types be mitigated with grassland or shrubland habitat types. Table 2 shows the acres of anticipated permanent and temporary impacts in each habitat category by habitat subtype for the RFA 4 and 5 Solar Facilities.

Table 2. Anticipated Temporary and Permanent Impacts by Habitat Category and Habitat Subtype

ODFW Habitat Category	Habitat Type	Habitat Subtype	Temporary Disturbance (acres)			Permanent Disturbance (acres)		
			RFA 4	RFA 5	Total	RFA 4	RFA 5	Total
1	N/A	N/A	-	-	-	-	-	-
2	Grassland-steppe	Grassland	0.1	-	0.1	-	0.9	0.9
	Riparian	Riparian Trees	<0.1	-	<0.1	-	-	-
	Shrub-steppe	Sagebrush Shrub- Steppe	2.7	-	2.7	11.8	-	11.8
	Category 2 Subtotal		2.8	-	2.8	11.8	-	12.7
3	Upland	Upland Trees	-	-	-	-	5.0	5.0
	Category 3 Subtotal		-	-	-	-	5.0	5.0
4	Grassland-steppe	Grasslands	6.1	-	6.1	36.2	-	36.2
	Category 4 Subtotal		6.1	-	6.1	36.2	-	36.2
5	N/A	N/A	-	-	-	-	-	-
6	Agriculture	Irrigated Cropland	4.9	-	4.9	39.4	-	39.4
	Agriculture	Non-Irrigated Cropland	173	43.9	217	525	1,237	1,762
	Developed	Developed	3.5	0.2	3.7	0.2	1.1	1.3
	Category 6 Subtotal		181	44.1	225	565	1,238	1,803
Total			190	44.1	234	613	1,244	1,857
Note: Totals in this table may not sum correctly due to rounding. "-" = no impact; <0.1 = greater than zero but less than 0.05 acres.								
1. There are no disturbances in surface water habitat types. Refer to Exhibit J of RFA 4 for detailed analysis of disturbances to wetlands and waters.								
2. N/A = not applicable; ODFW = Oregon Department of Fish and Wildlife								

5.0 Methods for Calculating Mitigation

RFA 4 and 5 Solar Facility impacts would result in a mitigation need of 71.9 acres, including 66.5 acres of mitigation for permanent impacts and 5.4 acres of mitigation for temporary impacts (Table 3 and Table 4).¹ The actual acres of temporary and permanent impacts and the associated mitigation requirements will be confirmed following construction. The Certificate Holder is not proposing compensatory mitigation under the ODFW Fish and Wildlife Habitat Mitigation Policy for impacts to Category 6 habitat.

¹ Temporary impacts result in temporal loss of habitat function and values from the time an impact occurs to the time when the restored habitat provides a pre-impact level of habitat function. Sagebrush steppe and riparian tree habitat subtypes are reasonably expected to require a longer restoration timeframe (5+ years) and therefore would be expected to result in temporal loss requiring compensatory mitigation beyond the Certificate Holder's revegetation obligation.

Table 3. Calculating Mitigation for Permanent Impacts

Habitat Category	Impact Acres	Mitigation Ratio ¹	Mitigation Need	Mitigation Description
Category 2	12.7	2:1	25.4	The mitigation goal for Category 2 habitat is “no net loss” and “net benefit.” Accordingly, mitigation for permanent impacts on Category 2 habitat needs to demonstrate a net benefit in quality or quantity.
Category 3	5.0	1:1	5.0	The mitigation goal for Category 3 habitat is “no net loss” in quantity or quality.
Category 4	36.2	1:1	36.2	The mitigation goal for Category 4 habitat is “no net loss” in quantity or quality.
Category 6	1,803	0:1	--	The mitigation goal for impacts on Category 6 habitat is minimization; no compensatory mitigation proposed.
Grand Total	1,857	--	66.5	--
1. Acres mitigation per acres impacted. Note: Totals in this table may not sum correctly due to rounding.				

Table 4. Calculating Mitigation for Temporary Impacts

Habitat Category	Habitat Subtype	Impact Acres	Mitigation Ratio ¹	Mitigation Need	Mitigation Description
Category 2	Riparian Trees	0.01	2:1	0.02	The mitigation goal for Category 2 habitat is “no net loss” and “net benefit.” Accordingly, mitigation for permanent impacts on Category 2 habitat needs to demonstrate a net benefit in quality or quantity.
	Grassland-steppe	0.1	0:1	-	
	Sagebrush Shrub-steppe	2.7	2:1	5.4	
Category 4	Grassland-steppe	6.1	0:1	-	The mitigation goal for Category 4 habitat is “no net loss” in quantity or quality. The proposed mitigation ratio would result in a lesser amount of acreage of mitigation than what is impacted by the Solar Components. Combined with restoration of temporary disturbances, the proposed mitigation ratio is intended to account for the temporary loss of habitat functionality and meet the “no net loss” goal. Temporary disturbances to Category 4 Grassland-steppe are not mitigated beyond restoration.
Category 6	Agriculture, Developed	225	0:1	-	The mitigation goal for impacts on Category 6 habitat is minimization; no compensatory mitigation proposed.
Grand Total		234	--	5.4	--
1. Acres mitigation per acres impacted. Note: Totals in this table may not sum correctly due to rounding. “-” = none.					

6.0 Biglow Canyon Wind Farm Habitat Mitigation Area

The HMA is the area where the Certificate Holder is proposing to perform enhancement and preservation actions that are in addition to the revegetation of areas of temporary disturbance associated with the RFA 4 and 5 Solar Facilities. The HMA must be large enough and have the characteristics to meet the standards set in OAR 635-415-0025.

According to ODFW standards, areas appropriate for mitigation of Category 2 habitat impacts must provide “in-kind” mitigation that creates similar structure and function to that being disturbed, be “in-proximity” to the RFA 4 and 5 Solar Facilities, and have potential for habitat enhancement. The mitigation goal for Category 3 habitat is “no net loss” in quantity or quality and mitigation for impacts to Category 4 habitat may be “in-kind” or “out-of-kind” and “in-proximity” or “out-of-proximity.”

The Certificate Holder identified 75 acres for the HMA located within 880 acres of land owned by the Certificate Holder adjacent to the Carty Generating Facility in Morrow County to fulfill the mitigation need for the RFA 4 and 5 Solar Facilities (Figure 1). The Certificate Holder identified an area larger than its mitigation need to allow for flexibility in management of the HMA. The Certificate Holder discussed the location of the HMA with ODFW and received assurance that the mitigation goals of “in-kind” and “in-proximity” can be met.

6.1 Habitat Assessment and Mitigation Accounting

The Certificate Holder identified 75 acres of suitable in-kind and in-proximity habitat within an 880-acre parcel (Figure 1). The HMA is adjacent to the existing PGE-owned Conservation Area and a conservation area maintained by The Nature Conservancy (South Farm Conservation Area; Figure 1). Habitat mapped within the HMA includes rabbitbrush shrubland and broom snakeweed shrubland (Figure 1). The HMA is dominated by Sandberg’s bluegrass (*Poa secunda*), bluebunch wheatgrass (*Pseudoroegneria spicata*), and intermittent areas of needle-and-thread grass (*Hesperostipa comata*), as well as cheatgrass (*Bromus tectorum*). There are also occasional occurrences of green rabbitbrush (*Chrysothamnus viscidiflorus*), gray rabbitbrush (*Ericameria nauseosa*), big sagebrush (*Artemisia tridentata*), fiddleneck (*Amsinckia menziesii*), and yarrow (*Achillea millefolium*).

The HMA is within a Priority Wildlife Connectivity Area (PWCA) connector (ODFW 2026).² The HMA also contains Category 1 and Category 2 habitat associated with Washington ground squirrel (*Urocitellus washingtoni*) colonies that are immediately adjacent to the HMA.

² The Oregon Conservation Strategy (ODFW 2025) defines PWCA “regions” as the combined top one percent of connectivity priorities across all 54 Oregon Connectivity Assessment and Mapping Project (OCAMP) species’ connectivity models. Regions are large, contiguous areas and represent the highest-value habitat for facilitating species movement throughout the state. PWCA “connectors” are described as pathways that provide the best available habitat for facilitating movement between regions. Connectors may pass through high-quality habitat in intact, relatively undisturbed parts of

The Certificate Holder plans to mitigate for the habitats impacted by placing a conservation easement on the appropriate acreage and by providing habitat uplift through the habitat enhancement and monitoring activities described below.

6.2 Habitat Enhancement Actions

The objectives of habitat enhancement and restoration are to protect habitat within the HMA from degradation and improve the habitat quality of the HMA. The Certificate Holder shall initiate habitat enhancement actions for the RFA 4 and 5 Solar Facilities before beginning operation. The Certificate Holder shall restrict uses of the HMA that are inconsistent with the goal of no net loss and net benefit of Category 2 habitat and no net loss of Category 4 habitat, as designated in the conservation easement. The Certificate Holder shall implement habitat enhancement actions as described in this HMP and as specified in the Site Certificate. Enhancement actions and monitoring procedures are detailed below, as finalized in this HMP prior to construction. Besides legal protection to ensure no development, enhancement actions for the HMA include those detailed in the following subsections.

6.2.1 *Restoration and Enhancement Actions in the Event of Fire*

The following actions will be implemented at the HMA if the site is damaged from fire or fire suppression efforts (e.g., vehicular disturbance or chemical application):

1. GIS mapping of fire scar, fire lines, and/or chemical disturbance areas.
2. Photopoint monitoring at all established monitoring points.
3. Assessment of fire severity and native vegetation damage to inform restoration needs.
4. Invasive annual grass surveys will be performed in fall and early spring for a minimum of two years after the fire. Noxious weed surveys will be performed according to HMP noxious weed survey requirements.
5. Annual grass treatments as needed (pre- and/or post-seeding) as native vegetation establishes, especially during the first two growing seasons.
6. Noxious and nuisance weed treatments as needed until noxious weeds are controlled at pre-fire levels.
7. Seeding where warranted during the first seeding window following the fire (based on timing of fire and climatic conditions but likely September–November) using an appropriate native bunchgrass seed mix at a drill seeding rate of 8–10 pounds pure live seed per acre. Appropriate seeding method will be determined based on the size and severity of the disturbance area and presence of Washington ground squirrel colonies. Seeding methods may include drill seeding or broadcast seeding (hand, ~~atv~~ ATV with harrow, and/or aerial). Low intensity fire is often beneficial to native vegetation and may not require reseeding.

the landscape, as well as the best remaining marginal habitat in developed or degraded areas. The Solar Components impact a PWCA region along Biglow Canyon.

8. Sagebrush will be planted to replace burned sagebrush stands to pre-fire conditions.
9. Secondary seeding of native forbs, as needed, after native grasses have reestablished. Need for forb seeding will be determined based on the presence/absence and diversity of forbs naturally established in the burn area.

6.2.2 Noxious Weed Prevention, Inventory, and Control

Weeds will be controlled as needed to maintain and enhance habitat quality within the mitigation area, with the goal of working toward eradication of targeted noxious weeds or, if eradication is not practical, decreasing their abundance to minimize impacts on native plant communities. Weeds known to occur within the HMA include rush skeletonweed (*Chondrilla juncea*), knapweed (*Centaurea* spp.), and yellow starthistle (*Centaurea solstitialis*). Recommended control methods are provided in Table 5. The Certificate Holder shall obtain ODFW's approval prior to the use of pesticides. Controlling weeds in the HMA should promote growth of native vegetation.

Table 5. Recommended Noxious Weed Control Methods

Scientific Name	Common Name	Recommended Treatment
<i>Centaurea</i> sp.	knapweed	Mechanical: Manually remove entire plant, including taproot, prior to flowering. Chemical: Apply to actively growing plants in fall or spring from rosette to bolting stage. Biocontrol: Agents include several seed feeding flies and weevils and a root-boring beetle.
<i>Centaurea solstitialis</i>	yellow starthistle	Mechanical: Manually remove entire plant including taproot. Chemical: Post-emergent herbicide application to rapidly growing plants prior to flowering (rosette to bolting stage). Biocontrol: Agents include two seed weevils, a seed fly, and a rust.
<i>Chondrilla juncea</i>	rush skeletonweed	Mechanical: Regenerates from root fragments. Manually remove entire plant, including taproot, prior to flowering. Will require multiple efforts during the growing season for more than one year. Chemical: Apply to rosettes in spring immediately before or during bolting. Biocontrol: Agents include a gall midge, gall mite, rust fungus, and root-mining moth.
Sources: DiTomaso et al. 2013; ODA 2025; Prather et al. 2019.		

6.2.3 Access Control

The Certificate Holder will monitor and control access to the HMA and will post informative signs depicting the area(s) as "protected" and will include natural resources information as appropriate for the life of the Facility. Primary access to the PGE property is controlled by a gate off Tower Road

north of the Carty Generating Station (currently used by PGE and The Nature Conservancy), the gated entrance to the Boardman Plant, and a gated road from Ione to the south. The Nature Conservancy and Three Mile Canyon Farms may occasionally use the two-track access crossing PGE's property to access the Farm's conservation area. Approved access to the HMA is currently limited to such occasional approved use of access roads and PGE's general operational needs. No livestock grazing is currently occurring within the HMA, and grazing is a restricted use identified in the conservation easement, as established in Section 2.0 of this HMP. Periodic monitoring will be conducted to evaluate the effectiveness of access control measures and signage maintenance needs.

6.2.4 Sagebrush Planting

To mitigate for temporal and permanent impacts to Category 2 sagebrush habitat, the Certificate Holder will plant 29.1 acres of sagebrush in the HMA, focusing on enhancing and expanding remnant stands of shrubs that were impacted by past wildfires (Figure [24](#)). Prior to construction, shrub density within the Category 2 sagebrush shrub-steppe disturbance area at the RFA 4 Solar Facility ~~within the Facility permanent disturbance area~~ was 5 shrubs per acre as of May 26, 2026 (51 sagebrush individuals were observed across 11.5 acres). Sagebrush will be planted at the HMA in fall ~~during the first year of operations~~ 2028 to achieve 150 percent of shrub density in the RFA 4 Solar Facility disturbance area prior to construction for a total of 8 shrubs per acre after 5 years. This planting density was approved by ODFW on August 27, 2026 (Jamie Bowles, ODFW, pers. comm.). Shrubs will be planted in clusters of 10 plants throughout the shrub planting area and monitoring locations will be established at a subset of the planting clusters as described in Section 7.4.

6.2.5 Annual Grass Treatment

The Certificate Holder will treat the entire HMA with a pre-emergent herbicide such as Rejuvra to control invasive annual grasses in the fall of 2027. Additional herbicide spot applications will be conducted as necessary to control annual grasses prior to seeding or planting. The timing of sagebrush planting and pollinator habitat enhancement will depend on the timing and efficacy of invasive annual grass control.

6.2.6 Enhancing Pollinator Habitat

Many native forb species that provide pollinator habitat are well established throughout the HMA and are expected to increase after invasive annual grasses are controlled. If vegetation monitoring does not demonstrate an increase in native forb cover within 3 to 5 years of annual grass treatment, then the Certificate Holder will perform a single seeding of native forbs to enhance pollinator habitat. The location of pollinator habitat enhancement will be determined, if necessary, after 3 to 5 years of monitoring.

7.0 HMA Enhancement Monitoring

The Certificate Holder will use a qualified investigator (botanist, wildlife biologist, or revegetation specialist) to conduct monitoring at the HMA. The purpose of this monitoring is to evaluate, on an ongoing basis, the protection of habitat quality and the results of enhancement actions. Final monitoring methods are presented below for ODOE concurrence. The vegetation monitoring methods for cover and density, including fixed radius plots and Daubenmire cover scale, were recommended and reviewed by Tim Walters of Haley Aldrich as the consulting botanist for ODOE in May 2026.

The investigator will visit the HMA during the growing season for the first ~~five~~5 years post planting/weed treatment. If success is achieved at year ~~five~~5, monitoring may be reduced to every fifth year thereafter (in years divisible by ~~five~~5, unless otherwise specified for specific measures) for the life of the Facility. Monitoring shall include an assessment of the following:

1. General quality of vegetation cover (dominant species, structural stage, etc.), as determined by ocular estimates, measurements, and photo points;
2. Success of weed control efforts;
3. Sagebrush density in enhancement areas;
4. Success of remedial actions to restore habitat quality in damaged areas (such as managed weed infestations, fire damage, and any necessary seeding/planting areas), as determined by vegetation cover (ocular estimate) and photo points;
5. Incidental wildlife occurring within the HMA;
6. Environmental factors found on site during monitoring activities and annual precipitation records; and
7. Surveys of resident special status wildlife species (Washington ground squirrel) that have been documented during previous monitoring or survey efforts within the HMA, using existing protocols approved by ODFW.

7.1 Monitoring Location Establishment

Monitoring locations were established using the following methods and are shown in Figure ~~24~~21. The HMA was divided into four quadrants (NW, NE, SW, SE). The monitoring location within each quadrant (photo point and fixed radius plot) was randomly generated using the ESRI ArcGIS Pro Create Random Points tool. If a point was located within 100 feet of the proposed HMA boundary or was located within line of sight of another monitoring location, an alternate randomly generated location was used. The fifth monitoring location was placed at the centroid (C) of the four quadrants. A monitoring location (W1) was also set up at one moderately dense diffuse knapweed and yellow starthistle infestation incidentally located during surveys, and representative photos of the infestation were taken at this location. All monitoring locations were monumented with a t-post and survey nail. The NW and NE monitoring locations are in the general area designated for shrub

plantings. Additional monitoring locations to monitor weed infestations and shrub planting success will be established as needed.

7.2 Photo Points

At each monitoring location (NW, NE, SW, SE, C), a photo will be taken in each of the four cardinal directions. Baseline photos were taken and the general vegetative condition was described at each monitoring location on May 13, 2026. Baseline photos are provided in Attachment 1.

7.3 Vegetative Cover

To determine vegetative cover, measurements will be made using 100 meter squared (m², 0.025 acre) fixed radius plots at the established monitoring locations. Each plot will be sampled using a 5.64-meter radius string secured around the t-post located at the center of the monitoring plot. Within each plot, percent cover measurements of ~~seven~~ cover types described in Table 6 will be recorded using the Daubenmire cover scale modified by Bailey and Poulton to account for the presence of small forbs commonly found in grasslands (Bonham 2013). The Class midpoints are then averaged for each plot to estimate the percent cover for each cover type. To ensure consistency during classification, a botanist will perform an initial plant species inventory of the HMA, categorizing all species as desirable, undesirable, or noxious.

Table 6. Cover Types

Cover Type	Description
Desirable grasses/ forbs	Seeded or naturally recruited native and desirable non-native herbaceous grasses species.
<u>Desirable forbs</u>	<u>Seeded or naturally recruited native and desirable non-native herbaceous forbs.</u>
Shrubs	Species such as rabbitbrush (<i>Chrysothamnus</i> sp.) and big sagebrush (<i>Artemisia tridentata</i>).
Undesirable species	Invasive, non-native species not found on ODA or county A, B, and T noxious weeds lists, such as cheatgrass (<i>Bromus tectorum</i>) and Russian thistle (<i>Salsola tragus</i>).
Noxious weeds	Species found on the ODA and/or county A, B, and T noxious weed lists.
Litter/mulch	Natural litter layer or mulch from hydroseeding.
Rock	Gravel, cobble, boulders, and bedrock.
Bare ground	Exposed soil with no litter or vegetation.

7.4 Density

Due to the patchy and uneven distribution of shrub and weed species throughout the HMA, density will be used to help determine success of shrub plantings and noxious weed management at monitoring locations. Measurements will be made using 100 m² fixed radius plots at each shrub and weed infestation monitoring point. Each plot will be sampled by using a 5.64-meter radius string secured around the t-post located at the center of the monitoring plot. Occasionally, a scaled down radius may be used for dense populations. Noxious weed species densities will be assessed at the

five established monitoring locations and at all weed infestation monitoring locations. Shrub density will be collected at monitoring locations within delineated shrub planting areas and only for planted species, such as big sagebrush. Shrub monitoring locations will be set up at a subset of sagebrush planting clusters throughout the shrub planting area.

For species density measurements, every plant species of interest that is rooted within the circular 100 m² plot will be counted. Larger plants that are on the line will be included if the majority of their stem is within the plot. The total number of individuals per acre will then be calculated by multiplying the density count per 100 m² by 40.4686.

7.5 Weed Monitoring

Annual noxious weed inventories will be conducted within the HMA during construction and in preparation for sagebrush planting, then every year for the first 5 years following implementation of initial enhancement. Noxious weed monitoring will be conducted every fifth year thereafter (in years divisible by 5) for the life of the Facility, concurrent with Washington ground squirrel monitoring. Control and monitoring of specific noxious weed occurrences will be conducted more frequently (at least every 2 years) in areas prioritized based on the 5-year monitoring results. Weed monitoring will to identify new noxious weed infestations~~occurrences~~, the status of existing ~~infestations~~~~occurrences~~, and the progress of weed control efforts in eradicating or reducing ~~infestations~~~~populations~~. Weed inventories will be conducted on foot. Detailed information on noxious weed infestations~~occurrences~~ will be collected, including, but not limited to, species, location, growth form, number of individuals, and density. In addition, the surveyor will identify potential weed control strategies for each ~~infestation~~~~occurrence~~, as well as constraints that could limit the effectiveness of weed control efforts.

Results from the surveys will be incorporated into a GIS database and provided to ODOE with the Monitoring Report. Weed ~~infestations~~~~occurrences~~ will also be photographed over time to document progress of weed control efforts.

8.0 Success Criteria

Enhancement actions will be assessed using the following success criteria. The sagebrush success criteria was adapted from Section 2 for ODOE review to reflect sagebrush density in shrub habitat impacted by the Facility.

- **Native and Invasive Vegetative Cover** - Decrease in invasive plant cover and increase in native/desirable plant cover at monitoring sites compared to baseline.
- **Sagebrush** - Sagebrush density in enhancement areas is 150 percent of pre-construction shrub density at the Facility 5 years after planting, for a total of 8 shrubs per acre.

9.0 Data Reporting and Timeline

The Certificate Holder will submit a report including wildlife and habitat monitoring data and analysis to the ODOE and ODFW during each monitoring year according to the schedule in Table 7. The Certificate Holder will notify the U.S. Fish and Wildlife Service and ODFW within one business day if any federal or state endangered or threatened species are killed or injured within the HMA. The Certificate Holder may include the reporting of wildlife monitoring data and analysis in the report required under OAR 345-026-0080 or submit this information as a separate document concurrently with the submittal of the report. In addition, the Certificate Holder will provide ODOE with any data or record generated by the investigators in carrying out this Plan upon request by ODOE.

Table 7. HMA Schedule of Implementation and Monitoring

Task	Schedule
Baseline Monitoring	During the first year of construction.
General Monitoring	Every year for the first five 5 years post planting/weed treatment. If success is achieved at year five 5, monitoring may be reduced to every fifth year <u>(in years divisible by 5)</u> thereafter for the life of the Facility.
Washington Ground Squirrel Monitoring	During the first, third, and fifth years after RFA 4 and 5 Solar Facility construction, and otherwise every fifth year <u>(in years divisible by 5)</u> thereafter for the life of the Facility.
General Weed Inventory, Control and Monitoring	During construction and in preparation for sagebrush planting, then every year for the first 5 years following implementation of initial enhancement. Long-term monitoring frequency to be determined after year 5 will be conducted every fifth year thereafter (in years divisible by 5) for the life of the Facility. <u>Long-term monitoring frequency to be determined after year 5 will be conducted every fifth year thereafter (in years divisible by 5) for the life of the Facility, concurrent with Washington ground squirrel monitoring. Control and monitoring of specific noxious weed occurrences will be conducted more frequently (at least every 2 years) in areas prioritized based on the 5-year monitoring results.</u>
<u>Annual Grass Treatment</u>	<u>Initial treatment in fall 2027 with follow up spot applications as necessary prior to seeding or planting.</u>
Sagebrush Planting	Fall of the first year of operations. <u>2028.</u>
Sagebrush Habitat Monitoring	Every year for the first 5 years following implementation of initial enhancement <u>and then every fifth year (in years divisible by 5) thereafter for the life of the Facility.</u> long-term monitoring frequency to be determined after year 5.
<u>Pollinator Habitat Enhancement</u>	<u>Native forbs will be seeded or planted to enhance pollinator habitat if vegetation monitoring does not demonstrate an increase in native forb cover within 5 years of annual grass treatment.</u>

10.0 Amendment of the HMP


This HMP may be amended from time to time by agreement of the Certificate Holder and EFSC, or ODOE, acting within its delegated authority of EFSC under ORS 469.402. Such amendments may be made without amendment of the site certificate. EFSC authorizes ODOE to agree to amendments to this HMP. ODOE shall notify EFSC of all amendments, and EFSC retains the authority to approve, reject, or modify any amendment of this plan agreed to by ODOE.

11.0 References

- Bonham, C. 2013. *Measurements for Terrestrial Vegetation*, 2nd Edition. Wiley. 272 p.
- DiTomaso, J.M., G.B. Kyser, S.R. Oneto, R.G. Wilson, S.B. Orloff, L.W. Anderson, S.D. Wright, J.A. Roncoroni, T.L. Miller, T.S. Prather, C. Ransom, K.G. Beck, C. Duncan, K.A. Wilson, and J.J. Mann. 2013. *Weed Control in Natural Areas in the Western United States*. Weed Research and Information Center, University of California. 544 pp.
- ODA (Oregon Department of Agriculture). 2025. Oregon Noxious Weed Profiles. Noxious Weed Control Program. <https://www.oregon.gov/oda/weeds/oregon-noxious-weeds/Pages/default.aspx>.
- ODFW (Oregon Department of Fish and Wildlife). 2026. Priority Wildlife Connectivity Areas. <https://oregonconservationstrategy.org/success-story/priority-wildlife-connectivity-areas-pwcas/>.
- Prather, T., T. Miller, A. Hulting, and E. Peachey. 2019. Control of Problem Weeds. In: Peachey, E. editor. *Pacific Northwest Weed Management Handbook*. Oregon State University. Corvallis, OR. P. Y-1 to Y-58.

Figures

\\tt.local\GIS\US\CES\Projects\PD\X\BrightNight\BiglowCanyon\Maps\HMA_Report_20250821\BrightNight_BiglowCanyon_HMAReport_20250924.aprx



**Biglow Canyon
Wind Farm
Solar Facility**

**Figure 1
Habitat Mitigation Area**

MORROW COUNTY, OR

- Habitat Mitigation Area
- County Boundary
- City/Town
- Carty 1 HMA boundary
- PGE Conservation Area
- South Farm Conservation Area
- Oregon PWCA Type**
 - Connector
 - Region
- WAGS Habitat**
 - Category 1 Habitat (785-foot Colony Buffer)
 - Category 2 Habitat (1500-meter Colony Buffer)
- Habitat**
 - Broom Snakeweed Shrubland
 - Developed
 - Rabbitbrush Shrubland



Reference Map



1:25,000

WGS 1984 UTM Zone 10N

0 1,250 2,500 5,000 US Feet


NOT FOR CONSTRUCTION

**Biglow Canyon
Wind Farm
Solar Facility**

**Figure 2
Sagebrush Planting and
Photopoint Locations**

SHERMAN COUNTY, OR

-  Habitat Mitigation Area
-  Sagebrush Planting Area
-  Photo Monitoring Location



1:2,500

WGS 1984 UTM Zone 10N

0 250 500 1,000 Feet

NOT FOR CONSTRUCTION

Attachment 1. Baseline Photo Monitoring



Photo 1. Photo Point C. Facing North. 5/13/2026.




Photo 2. Photo Point C. Facing East. 5/13/2026.



Photo 3. Photo Point C. Facing South. 5/13/2026.



Photo 4. Photo Point C. Facing West. 5/13/2026.




Photo 5. Photo Point NE. Facing North. 5/13/2026.




Photo 6. Photo Point NE. Facing East. 5/13/2026.



Photo 7. Photo Point NE. Facing South. 5/13/2026.



Photo 8. Photo Point NE. Facing West. 5/13/2026.



Photo 9. Photo Point NW. Facing North. 5/13/2026.



Photo 10. Photo Point NW. Facing East. 5/13/2026.



Photo 11. Photo Point NW. Facing South. 5/13/2026.



Photo 12. Photo Point NW. Facing West. 5/13/2026.




Photo 13. Photo Point SE. Facing North. 5/13/2026.



Photo 14. Photo Point SE. Facing East. 5/13/2026.



Photo 15. Photo Point SE. Facing South. 5/13/2026.



Photo 16. Photo Point SE. Facing West. 5/13/2026.



Photo 17. Photo Point SW. Facing North. 5/13/2026.



Photo 18. Photo Point SW. Facing East. 5/13/2026.



Photo 19. Photo Point SW. Facing South. 5/13/2026.



Photo 20. Photo Point SW. Facing West. 5/13/2026.



Photo 21. Photo Point W1. Site Detail 1. 5/13/2026.



Photo 22. Photo Point W1. Site Detail 2. 5/13/2026.



Photo 23. Photo Point W1. Site Detail 3. 5/13/2026.



Photo 24. Photo Point W1. Site Detail 4. 5/13/2026.