

<b>Model Purpose</b>	
Starting Inventory	ODF Stand Level Inventory (SLI) Data from the 2017 SLI Year-End Updates. Operations updates are current through December 2017. Stand growth is current through June 2018
Harvest Model Type	Linear Program Model
Model Period	Model Period: 100 years; Model interval: 5 years
Model Goals	Achieve desired future conditions Maintain Non-declining even flow; Finish with a minimum ending inventory Maximize Net Present Value
Spatial Data	GIS Data August 2019
General Costs	
Pond Values	10-year Forest 2 Market 3rd quarter 2019
Conservation Goals	As consistent with take avoidance and FMP landscape strategies
Aquatic Goals	As consistent with take avoidance and FMP aquatic strategies

# Harvest Model Data Prep and Information

09/18/2020

Field	Source	Responsibility	Process	Performed on	Export Date
<i>Unique_ID</i>	Calculated	Division	Calculated at the end for each polygon	Final polygon	20200503
<i>Acres</i>	Calculated	Division	Calculated at the end for each polygon	Final polygon	20200503
<i>District</i>	Harvest Units	Division	From Model_HarvestUnits	Original HU	20200429
<i>Fund</i>	Ownership Managed Lands	Division	HU identity to Ownership ManagedLands dissolved by type. 51 = BOF, 52 = CSL/CSL decert, 0 = not in ownership	HU_SLI_STD	20200429
<i>County</i>	County Boundaries	Division	HU Identity to Boundaries_County_Limit. Number = TRAScode	HU_SLI_STD	20200429
<i>Mbcode</i>	Management Basins	Division	HU polygon centroid joined to Boundaries_ManagementBasins. Number = MBcode	Original HU	20200429
<i>Std_id</i>	SLI_poly	Division	HU identity to SLI_poly	Original HU	20200429
<i>UPDATEYEAR</i>	SLI_poly	Division	HU identity to SLI_poly	HU_SLI_STD	20200429
<i>SLIVersion</i>	SLI_poly	Division	HU identity to SLI_poly	HU_SLI_STD	20200429
<i>HU_ID</i>	Harvest Units	Division	From Model_HarvestUnits	Original HU	20200429
<i>AEYD</i>	Calculated	Division	From Model_HarvestUnits	Original HU	20200429
<i>Avg_Slope</i>	Calculated	Division	Calculated for each Harvest Unit or HU identity to SLI_poly	Original HU	20200429
<i>Spur_Cost</i>	Harvest Units	Division	From Model_HarvestUnits	Original HU	20200429
<i>Node</i>	Harvest Units	Division	From Model_HarvestUnits	Original HU	20200429
<i>Thinable</i>	Non Thinnable Areas	Division	Polygon identity with Model_NonThinnableAreas. 0 = non thinnable, 1 = thinnable	HU_SLI_STD	20200429
<i>Logsys</i>	Harvest Units	Division	From Model_HarvestUnits	Original HU	20200429
<i>ROW</i>	Statewide Roads	Agency	csv file to Greg - Acres of final polygon by uid in 15ft buffer created from subset of statewide roads (within 15ft of NW managed lands, Status in ( NULL , ' ', 'B', 'Blocked', 'O', 'Unknown', 'Ex-exiting', 'Dormant', 'blocked', 'null' )). Created near end point by intersect.	final uid	20200723
<i>Easement</i>	Previous model data	District	HU polygon centroid joined to HUs designated as requiring easements in previous model runs (WL only). Number = number of model period HU is unavailable	Original HU	20200429
<i>Rec_thins</i>	Management Harvest Operations	District	HU identity to recent thin layers supplied by districts or extracted from Management_harvest_operations (status = completed, type= partial cut, year completed >= 2006). For each HU of: 1 - 9 = how many years since thinning (2014-2006), 0 = not recently thinned. Then joined to HU polygons.	Original HU	20190819
<i>H_NoThin</i>	Previous model data	District	HU centroid joined to HUs designated as H_NoThin in previous model runs (TL only). 1 = no thin, 0 = thinning ok	Original HU	20200429
<i>SNC_zone</i>	Biology SNC Zone	Division	HU centroid joined to Biology_SwissNeedleCastZone. SNCZONE = 1 (Severe Impact), SNCZONE = 2 (Light to Mod Impact), SNCZONE = 0 (Outside zone of impact). BIOLOGY_SwisNeedleCastZone 1 closest to ocean, 2 farther from ocean.	Original HU	20200429
<i>Site_grp</i>	SLI points/Tod Haren	Division	Average of site_phy values > 0 for sli_points in HU.	Original HU	Tod
<i>AdminRem</i>	FLMC	Division	Polygon identity to all special use (except riparian/ops limited/wildlife) and HVCA plants (except old growth) in FLMC. 1 = removed, 0 = not removed	HU_SLI_STD	20200503

Field	Source	Responsibility	Process	Performed on	Export Date
<i>OldGrowth</i>	FLMC	District	Polygon identity to all plants identified as old growth in FLMC. 1 = old-growth, 0 = not old growth	HU_SLI_STD	20200503
<i>Rip_Code</i>	Terrainworks modeled stream lines/ODF CFMP Buffer widths 20200512	Division	csv file provided to Greg - RCA acres by final uid. ROW erased from RCA.	final uid	20200723
<i>AA_ID</i>	Aquatic Anchors	Division	HU centroid in Biology_Aquatic Anchor. Number = aquatic anchor id, 0 = Not in AA	Original HU	20200429
<i>In_Gorge</i>	Raster Layers/modeled streams	Division	Polygon identity with modeled inner gorge polygons. 1 = inner gorge, 0 = not inner gorge	HU_SLI_STD	20200430
<i>FPA_Wild</i>	FLMC	Division	Polygon identity with HVCA wildlife habitat areas identified as Eagles/BT pigeons/GB Herons/Ospreys in comments field of FLMC. 1 = designated wildlife habitat, 0 = not wildlife habitat	HU_SLI_STD	20200503
<i>Ta_mmma</i>	Murrelet Management Areas/MMMA final	Division	Current FMP (CFMP) run we switched to using 2 csv files to Greg, acres by final uid and percent by final uid.	final uid	20200503
<i>Tas_Id</i>	Terrestrial Anchors	Division	HU centroid joined to Biology_TerrestrialAnchor. Number = TA id, 0 = not in a TA site	Original HU	20200429
<i>LSPS_Bid</i>	Raster Layers/modeled streams	District	Polygon identity with district LSPS layer. Number = Object ID, 0 Not in a LSPS basin	HU_SLI_STD	20200503
<i>LSPSRisk</i>	Raster Layers/modeled streams	District	Polygon identity with district LSPS layer. 2 = intermediate risk, 3 = substantial risk, all others = 0	HU_SLI_STD	20200503
<i>HLHL_Code</i>	Buren reviewed 2014 HLHL data	Division	csv file provided to Greg - acres by final uid. ROW and RCA erased from HLHL.	final uid	20200723
<i>Elev</i>	Harvest Units	Division	Use HU Centroid, raster tool to get elevation of centroid		20200429
<i>DistOcean</i>	Harvest Units	Division	Create near table calc tool, gives lookup table with feature ID's From/To Distance, join back to the HU Centroids, create field.		20200429

Keep from original HU's

District  
P\_Ground  
P\_Cable  
Spur\_Cost  
LogSYS

## General Model Rules

Number	Value Group	Model Rule	Implementation	GIS Check	Reports Check	Who makes decision	Relative Impact on Model outputs	Comparative Analysis Run	Notes
1	Admin rules	NA	No harvest > 120 acre patch size.	Visually check patch sizes for large patch sizes	NA	<a href="#">OAR 629-630-0000</a>	TBD	Yes	
3.2	Admin rules	Landslide and public safety	No regen harvest in HLHL polygons within high risk basins; If over 50% of a harvest unit is encumbered by HLHL, exclude entire unit from harvest	Visually inspect moderate and high risk basins. Summarize acres harvested by LSPS basin.	Check classification reports for HLHL breakout.	<a href="#">OAR 629-630-0500</a>	Low	Yes	
4	Economic	Merchantability Specifications	Up to five log grades identified by species group. 40ft log grades merchandise consistent with the official rules for log scaling and grading bureaus	Check a subset of stands that are regen harvested. All stands should be greater than 40 years old.		Kevin	Medium	Yes	
5	Economic	Pond Values	Pond values are applied by species, grade, and region. Ten year average price from F2M market reports, PPI adjusted.	N/A		Kevin	High	Yes	
6	Economic	Stumpage Values	Stumpage is currently estimated as pond value - cost	Check GIS polygon layer for estimated pond/logging costs, and transport layer for estimated road costs	Check harvest reports for net revenue estimates	Kevin	High	Yes	
7	Economic	Breakage and Defect	Size based percent defect is removed from net volume estimates.	N/A	Compare gross volume with	Kevin	Medium	Yes	
8	Economic	In-Unit Spur Costs	Applied based on LEI HU estimates	N/A	Check spur costs reported in model outputs	Kevin	Low	Yes	
9	Economic	Road Maintenance Costs	Applied based on LEI model road estimates	N/A		Kevin	Low	Yes	
10	Economic	Hauling Costs	Calculated by species and size for each harvest unit. Each species and district is assigned to a typical mill site.	N/A	Review MSR Reports	Kevin	Low	Yes	
11	Economics	Logging costs	Taken from an OSU survey of local contractors	Check cost estimates in documentation folder	Review average logging cost by harvest type	Kevin	Medium to High	Yes	
12	Economics	Young Stand Silviculture	Average free to grow cost by district	N/A	Review MSR Report	Kevin	Low to Medium	Yes	
13	Growth & Yield	SNC growth&yield	Growth adjustments are applied to regeneration stand yield tables based on SNC Zone. Zone 1 = severe (code=1) Zone 2 = light/moderate (code=2)	N/A	N/A	Robbie	Medium	Yes	
14	Growth & Yield	Growth calibration	FVS adjustments include basal area growth modifiers by species and maximum SDI modifiers by species	Check stands at 2 different time periods to ensure that FVS growth assumptions reflect actual stand capabilities.	Check inventory reports	Tod	Medium	Yes	
15	Growth & Yield	Imputation	Non-measured stands assigned to similar measured stands	N/A	N/A	Tod	Medium	Yes	
16	Growth & Yield	Silvicultural pathways	Yield tables include multiple silviculture pathways such that the model can choose no harvest, regen, regen+pct, thinnings, and combinations of those options over multiple stand rotations.	Check stands at 2 different time periods to ensure that FVS growth assumptions reflect actual stand capabilities.	Check inventory reports	<a href="#">Tod</a>	High	Yes	

Number	Value Group	Model Rule	Implementation	GIS Check	Reports Check	Who makes decision	Relative Impact on Model outputs	Comparative Analysis Run	Notes
17	Growth & Yield	Initial inventory	Stands are projected from their measured date to 2023.	Ensure that current stand inventory estimates are reasonably close to actual volume.	N/A	Tod	Medium	Yes	
18	Growth & Yield	FVS implementation	Stands are grown using regionally appropriate FVS variants.	Check that growth rates are reasonable for a range of stands.	Growth rates represented by MAI and PAI	Tod	Medium	Yes	
19	Growth & Yield	Planting regimes	Site specific planting regimes have been developed and have been included by district.	N/A	N/A	Tod	Medium	Yes	
20	Growth & Yield	Green Tree Retention	5 green TPA > 16" DBH (Approx = QMD of stand)	Input growth/yield tables include a set amount of green trees/acre. The decision here is a policy choice, but can still check to ensure that leave tree volume projections are reasonable.	Use reports to check the amount of leave tree volume/tpa, etc.	Tod	Low	Yes	
22	Operations	Hemlock no-thin zones	Where the attribute "H_NoThin" = 1, the harvest unit receives 'grow only' prescriptions until the first clearcut. Tillamook specific	Check H_NoThin=1, and ensure no thinning in initial periods before a regeneration harvest.	No report to directly check no-thin rule.	Colleen/Jodi	Low	Yes	
23	Operations	Harvest Units	Harvest units have been delineated to identify likely harvest blocks	N/A	N/A	Colleen/Jodi	Medium	Yes	
25	Economics	In-unit spur cost	The in-unit spur construction costs are in \$/acre harvested for the first entry. On subsequent entries, 20% of the original construction cost was applied to account for the cost of reopening the road.	Spur costs are not included in GIS info.	Spur costs are included in road construction costs in the road cost summary report.	Kevin	Low-Medium(de pending on district)	Yes	
27	Economics	Road maintenance	<ul style="list-style-type: none"> <li>Maintenance costs are based on the long-term maintenance of a road segment and included the cost for replacing culverts over time.</li> <li>Maintenance costs are listed in the Road Segment Table in units of \$/MBF/Segment.</li> <li>Maintenance Costs were deducted from the gross value of all harvests that used the road.</li> </ul>	N/A	Review Model Reports	Kevin	Low-Medium(de pending on district)	Yes	
28	Economics	Hauling/Loading Costs	Defined in road inputs	N/A	Review Model Reports	Kevin	Medium	Yes	
29	Operations	Thinning	Minimum age for thinning set to 35 years				Medium	Yes	
30	Growth & Yield	Ending Inventory Level	Minimum ending inventory: 20k bdf/ac * operable acres (or 0.5 * avg CMAI)	N/A	Verify Ending inventory exceeds this constraint		Medium	Yes	

## Policy Model Rules

Number	Value Group	Model Rule	Implementation	GIS Check	Reports Check	Who makes decision	Relative Impact on Model outputs	Comparative Analysis Run	Notes
1	Economic	Discount Rate and Real Price Appreciation	3% discount rate; cost and price appreciation assumed constant.	N/A	Compare nominal cashflow with NPV	<a href="#">Finicial Metrics Issue Paper (ODFnet)</a>	Medium	Yes	
2	Harvest Objectives	Non-Declining Even Flow	NDEF relative to current IP, by district	N/A	Check volume report to ensure departure is within specified limits	Brian	High	Yes	
3	Operations	Minimum regen harvest age	Miminum regen stand age is 40 years for all stands	Check a subset of stands that are regen harvested. All stands should be greater than 40 years old.	Harvest volume by age class	Brian	Medium	Yes	
4	Planning unit	Scale of Plannning Unit	Planning area is at district-level	N/A		Brian	Medium	Yes	related to NDEF

## Model Scenario Rules

Number	Value Group	Model Rule	Implementation	GIS Check	Reports Check	Who makes decision	Relative Impact on Model outputs	Date	Comparative Analysis Run	Notes
1	Aquatic/Riparian	No harvest in identified riparian buffers	No harvest allowed in riparian buffers.	No harvest in any polygon with RIP_FMP > 0	Look at Classification reports - there should be no harvest in any of the RIP_FMP categories	<a href="#">FMP Aquatic Stratgies</a>	Low-High, depending on buffer width		Yes	
2	Aquatic/Riparian	Potential Debris Flow Tracks (PDFT)	No harvest allow in PDFT	No harvest in any polygon with RIP_FMP=4	Look at Classification reports - there should be no harvest in any of the RIP_FMP=4 categories	Brian	Low		Yes	
3	Aquatic/Riparian	Inner Gorge	No harvest allowed in inner gorge.	No harvest in In_Gorge = 1	Classification Summary Reports= no harvest in inner gorge	Brian	Medium		Yes	
4	DFC Free-Up	Landscape design is freed once complex structure goal is attained	Mapped DFC is released once Complex Structure is achieved across landscape. DFC is released by district: Astoria, Forest Grove, West Oregon & Tillamook : After year 70 North Cascade & Western Lane: After Year 35 Coos: N/A - no LD has been designated Southwest: After year 20	no clearcut harvest within DFC prior to free-up	clearcut within DFC throughout duration		High		Yes	
5	Complex Structure Goal	mimimize time to achieve desired ammount of complex structure	Total Complex Structure must meet or exceed levels present when mapped DFC is released. Once a stand becomes complex, LYR or OFS, it may not be harvested for 20 years	Check a subset of stands that are regen harvested. All stands should be greater than 40 years old.	Amount of complex struture at end of time period per district		High		Yes	
6	T&E Red Tree Vole	T&E projected encumberance	adding potential incumberance based on habitat suitability models (After year 10 )	Model doesn't harvest suitable habitat after year 10	Acres of high suitable red tree vole habitat harvested	Nick	Medium-High (depending on district)		Yes	
7	T&E Marbled Murrelet	T&E projected encumberance	adding potential incumberance based on habitat suitability models (After year 10 )	Model doesn't harvest suitable habitat after year 10	Acres of high suitable Marbled Murrelet harvested	Nick	Medium-High (depending on district)		Yes	
8	T&E NSO	T&E projected encumberance	adding potential incumberance based on habitat suitability models (After year 10 )	Model doesn't harvest suitable habitat after year 10	Acres of high suitable NSO harvested	Nick	Medium-High (depending on district)		Yes	
9	Take Avoidance (scenario)	NSO Take Avoidance	Use NSO 60% and 40% habitat model "floating" circles for outer/inner circles	Visually check GIS layer to ensure that near-term entries of circles are reasonable.	Check Summary reports, both individual circle reports (including both inner/outer circles) and aggregated owl circle reports.	Nick	Medium-High (depending on district)		Yes	

Number	Value Group	Model Rule	Implementation	GIS Check	Reports Check	Who makes decision	Relative Impact on Model outputs	Date	Comparative Analysis Run	Notes
10	Take Avoidance (scenario)	Marbled Murrelet Take Avoidance	No action in MMMA's or additional incumbered acres	Visually check GIS layer to ensure that no harvest occurred within listed areas	Acres of MMMA's and Additional incumbered acres harvested	Nick	Medium-High (depending on district)		Yes	
11	Species of Concern (SOC)	Terrestrial Anchors	No clearcuts nor partials cuts that results in less than 80ft <sup>2</sup> /ac BA nor residual SDI less than 35% of the max	Check implimentation of harvest prescriptions were followed	acres harvested by description type	Nick	High		Yes	
12	Species of Concern (SOC)	Aquatic Anchors	Increased buffers within aquatic anchors	Check buffer width	acres harvested within aquatic anchor buffers	Mike	Medium		Yes	

## Reforestation Prescriptions

District	Regime	Species	TPA
Astoria	WXR1	DF	120
		RA	15
		RC	15
		WH	150
		<i>Total</i>	<i>300</i>
	WXR2	DF	150
		RA	6
		RC	15
		WH	129
		<i>Total</i>	<i>300</i>
	DXR3	DF	193
		RA	35
		RC	18
		WH	105
		<i>Total</i>	<i>350</i>
1DR4	DF	225	
	RA	15	
	WH	60	
	<i>Total</i>	<i>300</i>	
DXR5	DF	210	
	RA	17.5	
	NF	35	
	WH	87.5	
	<i>Total</i>	<i>350</i>	
Forest Grove	1DR3	DF	392
		GF	44
		<i>Total</i>	<i>436</i>
	DXR1	DF	174
		NF	87
		RC	87
		WH	87
		<i>Total</i>	<i>436</i>
	DXR2	DF	327
		NF	109
<i>Total</i>		<i>436</i>	
Tillamook	1WR1	DF	35
		NF	35
		RA	35
		RC	35

\* All regimes include 2 dominant species green trees representing the minimum FPA requirement

\* All stands include 2 snags representing the minimum FPA requirement

District	Regime	Species	TPA
		WH	210
		<i>Total</i>	<i>350</i>
	1WR2	DF	35
		RA	35
		RC	35
		SS	20
		WH	225
		<i>Total</i>	<i>350</i>
	WXR3	DF	105
		NF	35
		RA	35
		RC	35
		WH	140
		<i>Total</i>	<i>350</i>
	WXR4	DF	105
		RA	35
		RC	35
		SS	35
		WH	140
		<i>Total</i>	<i>350</i>
West Oregon	DXR1	DF	200
		RA	20
		RC	40
		WH	100
		<i>Total</i>	<i>360</i>
	DXR2	DF	210
		GF	20
		RA	20
		RC	40
		WH	70
		<i>Total</i>	<i>360</i>
	DXR3	DF	180
		RA	20
		RC	50
		SS	20
		WH	90
		<i>Total</i>	<i>360</i>
	OTR4	DF	110
		RA	70
		RC	50
		SS	40

District	Regime	Species	TPA
		WH	90
		<i>Total</i>	<i>360</i>
North Cascade	1DR1	DF	435
		RA	35
		WH	30
		<i>Total</i>	<i>500</i>
	DXR2	DF	348
		RA	30
		RC	87
		<i>Total</i>	<i>465</i>
	OTR3	DF	60
		NF	240
		WH	30
		<i>Total</i>	<i>330</i>
	WXR4	RA	10
RC		150	
WH		150	
<i>Total</i>		<i>310</i>	
Western Lane	DXR1	BM	15
		DF	265
		RA	15
		RC	25
		WH	30
		<i>Total</i>	<i>350</i>
	DXR2	BM	20
		DF	150
		RA	20
		RC	70
		WH	100
		<i>Total</i>	<i>360</i>
	OTR3	BM	35
DF		70	
RA		50	
RC		70	
WH		125	
<i>Total</i>		<i>350</i>	
Coos	1DR1	DF	285
		RA	15
		<i>Total</i>	<i>300</i>

District	Regime	Species	TPA
	DXR2	DF	200
		RC	200
		<i>Total</i>	<i>400</i>
	DXR3	DF	200
		WH	200
		<i>Total</i>	<i>400</i>
	DXR4	DF	180
		RA	40
		RC	92
		WH	88
		<i>Total</i>	<i>400</i>
Southwest	DXR1	DF	335
		PP	40
		SP	40
		IC	20
		<i>Total</i>	<i>435</i>

## Yield Table Prescriptions

### Moderate Commercial Thins

Thin from below, relative SDI constrained  
Conifer Stands Only

Dominant SPP Max SDI	DF	600	* SDI max for thinning is different than SDIMAX growth modifier
	WH	720	
Trigger RD	55,60		
Target RD	35,30		
Max DBH	45		
Min DBH	8		
Min Harv BdFt	Tillamook	6000	
	Others	7000	
Species preference	DF,WH,RA,BM,SS,RC		
Re-entry delay	15		

### Precommercial Thins

Thin from below, residual TPA target

Minimum age	10
Maximum age	23
Triger TPA	300
Residual TPA	220

### HCA Prescriptions

Maximum of 2 thinnings through age 90  
Less than 40% of basal area removed (surrogate for cover)

## Delivered Log Prices

10 year average from F2M reports, adjusted for inflation using the logging producers price index

Prices as of 9/15/2019

### Westside

Species	Prod-1	Prod-2	Prod-3	Prod-4	Prod-5
DF	615	600	429	110	110
WH	505	502	403	110	110
RA	682	510	510	110	110
SS	419	465	465	110	110
RC	1076	1032	1032	110	110
BM	453	453	453	110	110
NF	505	502	403	110	110
SF	505	502	403	110	110
CX	505	502	403	110	110
HX	453	453	453	110	110

### Southwest

Species	Prod-1	Prod-2	Prod-3	Prod-4	Prod-5
DF	649	553	500	140	0
PP	400	350	350	100	0
SP	400	350	350	100	0
WF	473	436	402.5	112.5	0
IC	1057	279	0	0	0
CX	473	436	402.5	112.5	0
HX	635	471	372	0	0

## Transportation

hu\_trans\_costs\_20200124.csv

Typical mill assigned by species and log size - Provided by Bodi D. on 1/2020

Road network combines HnH era roads, state highway layer, and "stubs" in Coos to complete the network. This is to provide reasonable haul costs, but is inconsistent with the official roads data

ESRI network analysis extension was used to calculate a distance and cost matrix between each HU node and mill node.

Field	Source	Responsibility	Process	Performed on	Export Date	Comments
<i>hu_id</i>	<i>Harvest Units</i>	Division	Update HU polygon layer and export	01/24/2020	01/24/2019	
<i>node</i>	<i>Harvest Units</i>	Division	Update HU polygon layer and export	01/24/2020	01/24/2019	Links each HU to the model roads network Coos HUs stubbed from centroid to end of nearest primary road HUs with missing or incorrect nodes assigned to nearest existing node
<i>df_sw_mill</i>	<i>Asset Mgt</i>	GIS/Analyst	Update Mill nodes table, road network, and re-run distance matrix analysis	01/24/2020	01/24/2019	An "expected" mill destination was identified for each species and log size on each district. Each mill was identified on the model road network.
<i>df_lw_mill</i>	<i>Asset Mgt</i>	GIS/Analyst	"	01/24/2020	01/24/2019	
<i>wh_sw_mill</i>	<i>Asset Mgt</i>	GIS/Analyst	"	01/24/2020	01/24/2019	
<i>wh_lw_mill</i>	<i>Asset Mgt</i>	GIS/Analyst	"	01/24/2020	01/24/2019	
<i>alder_mill</i>	<i>Asset Mgt</i>	GIS/Analyst	"	01/24/2020	01/24/2019	
<i>pine_mill</i>	<i>Asset Mgt</i>	GIS/Analyst	"	01/24/2020	01/24/2019	
<i>cedar_mill</i>	<i>Asset Mgt</i>	GIS/Analyst	"	01/24/2020	01/24/2019	
<i>hwd_mill</i>	<i>Asset Mgt</i>	GIS/Analyst	"	01/24/2020	01/24/2019	
<i>pulp_mill</i>	<i>Asset Mgt</i>	GIS/Analyst	"	01/24/2020	01/24/2019	
<i>df_sw_haul</i>	<i>GIS Analysis</i>	GIS/Analyst	Update Hauling Rate and Rerun cost matrix	01/24/2020	01/24/2019	Hauling costs calculated as a function of travel time along the shortest path to each mill. Haul cost=2x travel time * \$65/hr
<i>df_lw_haul</i>	<i>GIS Analysis</i>	GIS/Analyst	"	01/24/2020	01/24/2019	
<i>wh_sw_haul</i>	<i>GIS Analysis</i>	GIS/Analyst	"	01/24/2020	01/24/2019	
<i>wh_lw_haul</i>	<i>GIS Analysis</i>	GIS/Analyst	"	01/24/2020	01/24/2019	
<i>alder_haul</i>	<i>GIS Analysis</i>	GIS/Analyst	"	01/24/2020	01/24/2019	
<i>pine_haul</i>	<i>GIS Analysis</i>	GIS/Analyst	"	01/24/2020	01/24/2019	
<i>cedar_haul</i>	<i>GIS Analysis</i>	GIS/Analyst	"	01/24/2020	01/24/2019	
<i>hwd_haul</i>	<i>GIS Analysis</i>	GIS/Analyst	"	01/24/2020	01/24/2019	

<b>Field</b>	<b>Source</b>	<b>Responsibility</b>	<b>Process</b>	<b>Performed on</b>	<b>Export Date</b>	<b>Comments</b>
<i>pulp_haul</i>	<i>GIS Analysis</i>	GIS/Analyst	"	01/24/2020	01/24/2019	
<i>miles</i>	<i>GIS Analysis</i>	GIS/Analyst	Rerun distance matrix	01/24/2020	01/24/2019	Total distance along the shortest path to the DF SW mill
<i>maintenance</i>	<i>GIS Analysis</i>	GIS/Analyst	Update road network with maintenance rate and rerun cost matrix	01/24/2020	01/24/2019	Total maintenance cost along the shortest path to the DF SW mill
<i>construction</i>		GIS/Analyst		NA		There is no mechanism to allocate construction costs between harvest units in the LP model

## Silvicultural Actions

### Regeneration Harvests

The intent of a Regeneration Harvest is to develop a new stand. In general, residual trees left after a Regeneration Harvest are intended to remain on the site through the life of the new stand and subsequent stands. All types of Regeneration Harvests retain less than 80 square feet of basal area per acre (based on trees greater than 11 inches DBH). The Harvest Types (within the Regeneration Harvest Goals) are best defined using residual trees per acre or square feet of basal area per acre; in either case, only trees greater than 11 DBH are counted.

Prescription	Description	Requirements
Clearcut (cc)	A Clearcut removes all (or nearly all) trees in a stand; however, the FMP and the Forest Practices Act (FPA) require that at least a few live trees be retained in each unit. Clearcuts will provide the best conditions for successful plantation establishment on almost all sites on State Forests.	Retain between 2 to 5 green trees or snags per acre Are subject to FPA rules for type 3 harvest (maximum size is 120 acres with green-up requirement)
	Retention Cuts look more like a partial cut or the first stage of a shelter wood harvest than a clearcut, however the focus of future management will be on the new/young trees in the stand, rather than the residual trees. At its highest density, a Retention Cut leaves nearly as much basal area as a Heavy Thinning, and the management focus may be on the existing cohort, the new cohort, or both. In this harvest type, regeneration is achievable and will help develop complex structure more quickly than after a clearcut. A Retention Cut will result in a stand with two distinct age classes that are well-distributed across the stand.	Retains between 33 and 80 square feet of basal area per acre (on Site Clas I, II, or III) Are subject to the FPA Rules for Type 1 Harvest Are designed to meet and exceed landscape goals for structural components
Retention Cut (RC)		

### Parital Cuts

The intent of a Partial Cut Harvest is to manage the growth and density of an existing stand. A prescription for a Partial Cut may be designed to increase the structural complexity of a stand, maximize volume growth, or capture tree mortality. A stand may be Partial Cut several times throughout its life. All Partial Cut harvest types retain at least 80 square feet of basal area per acre of trees greater than 11 inches DBH.

There are several forms and intensities of Partial Cuts, however the most common form of Partial Cut is thinning. Thinning prescriptions are often designed using measures of Stand Density Index or Relative Density and remove a portion of the trees from a stand in a generally uniform pattern. Sometimes thinning prescriptions are developed to increase the horizontal diversity within a stand; a diameter limit prescription often results in a stand with variable density.

The structure of a stand immediately after a partial cut (1 to 3 years) is very dependent on both the harvest prescription and the structure of the stand prior to harvest. Generally, the stand structure will remain the same or become more complex.

Prescription	Description	Requirements
Heavy Thin	A Heavy Thinning approaches the harvest intensity of a Retention Cut, and the management focus will be on developing a new cohort of trees to speed up understory development which leads to a new cohort throughout the thinning area. A heavy thinning results in the fast growth of individual trees, but reduces the total volume growth of the stand	Retains at least 80 square feet of basal area per acre and an SDI% of less than 30.
Moderate Thin	A Moderate Thinning provides for optimal stand growth and allows vigorous growth of the individual trees. Stand structure will continue to develop with a Moderate Thinning, and depending on species composition and site index, a new cohort of trees may be initiated.	Retains an SDI% of greater than or equal to 30 and less than 40.
Light Thin	A Light Thinning focuses on maintaining stand growth and health, however in order to achieve these goals, it must occur more frequently than a Heavy or Moderate Thinning in the same stand. More complex stand structure may not be developed with a Light Thinning and a new cohort of trees will may not be initiated. Early commercial thinning (ECT) falls under a light thinning.	Retains an SDI% of greater than or equal to 40 and less than 50.

Harvest Type Definitions Table

Harvest Types	Site Class <sup>3</sup>	Regeneration Harvest		Partial Cut	
		Clearcut	Retention Cut (RC)	Thinning	Group Selection
Residual Basal Area (sq. ft / acre)	I, II, III		$\geq 33$ & $< 80$	$\geq 80$	Residual BA/Acre of entire unit must meet PC BA standard <sup>4</sup>
	IV, V		$\geq 20$ & $< 50$	$\geq 50$	
	VI		$\geq 10$ & $< 40$	$\geq 40$	
Residual Trees Per Acre <sup>1</sup>		$\geq 2$ & $< 5$	$> 5$		
Residual Stand Density Index <sup>2</sup>				Generally, $\geq 30\%$ & $\leq 55\%$	
Unit Size (acres)		$\geq 5$ & $< 120$	$\geq 5$		$> 0.5$ & $< 5.0$ $< 50\%$ of total unit in patch cuts.

1. The NW and SW FMPs do not set a minimum size for residual trees, so use the FPA standard of an 11 DBH and 30 feet tall.
2. SDI based on all trees >5.6 DBH and the SDI% is based on a weighted average of the species that comprise a significant component of the stand.
3. The FPA defines harvest types based on the Cubic Foot Site Class
4. When a harvest unit is comprised of a combination of partial cut and patch cuts, then the entire unit must meet the Residual Basal Area standard for a Partial Cut.

## **Reforestation and Young Stand Management**

### **Planting**

#### Initial Planting

Initial planting occurs after a regeneration harvest. Planted seedlings will be well suited and adapted to the reforestation site, and where appropriate a mixture of species will be planted to increase diversity across the permit area. Density will vary from 250-536 trees per acre (TPA). Stocktype will be site specific and take in factors such as soil type, soil quality, and animal browse potential. Species selection will be on a site by site basis with goal of increasing diversity across the landscape to increase resiliency in the uncertainty of climate change. In areas of disease, such as Swiss needle cast (SNC) or laminated root rot, planted species will be of tolerant stock or from a resistant species with an emphasis on resistant species

#### Interplanting

Interplanting will occur when stocking levels fall below Forest Practice Act minimums. In certain instances, interplanting will occur to increase stocking on high quality sites to fully capture the site. In other areas, lower stocking will be acceptable as it will provide high quality early seral habitat while still meeting FPA requirements. Density will be site dependent, but range from 200-400 TPA.

#### Site Preparation

Site preparation is any planned measure to prepare a site for the favorable conditions for newly planted seedlings. Site preparation should not cause detrimental or excessive soil disturbance, and should be carried out in a cost-effective manner. Through site preparation, factors that are limiting for seedling survival and growth may be overcome. Such factors may include limited soil moisture, low light levels, and compacted soil. Logging slash can have positive and negative benefits and should be evaluated on a site by site basis. The three main site preparation techniques are mechanical, chemical, and broadcast burning, which are described below.

#### Mechanical

Mechanical site preparation is the use of mechanized equipment to rearrange or alter forest slash and/or disturb the forest surface layer and vegetation to create seedbeds or planting spots. Mechanical site preparation reduces competition of other vegetation with crop trees for light, water and nutrients. It can alter wildlife habitat, both positively and negatively, and this should be taken into consideration before use at each site. It can also be used to treat the adverse effects of past activities, such as compaction.

## Chemical

Chemical site preparation, which involves the application of herbicides, controls competing vegetation before planting or natural regeneration and during the early stages of seedling establishment. Applications occur by two primary methods, aerially by helicopter or ground based with the use of backpack application equipment. In general, herbicides are effective in suppressing most undesirable vegetation and are well suited for use on many sites. Herbicides, when properly applied, also have little site impact. Proper application implies application according to label instructions. However, in some areas, application of herbicides are of particular concern to other resource users and members of the public.

## Broadcast Burning

When properly applied on appropriate sites, prescribed burning can achieve many site preparation objectives. Fire can be used on steep terrain, does not compact the soil, and improves access for planting. Fire impacts can also improve seedling survival and growth by reducing competing vegetation. Prescribed burning is also used to remove slash piles throughout the site and on landings.

However, it also has disadvantages. The biggest disadvantage is the risk of escape, and intense fire can reduce the amount of soil nutrients. Prescribed burning can also reduce the amount of downed wood in a unit, decreasing the amount of suitable habitat for a number of species of concern. Burning can also increase the amount of unwanted vegetation, such as ceanothus and senecio, in certain parts of the permit area.

## Release Treatments

Release treatments usually occur in young stands and are designed to reduce competition for desirable tree species. They can also be used to alter species composition under pressure from insect and disease and favor species that are tolerant or resistant to threat. There are two types of release treatments, manual and chemical, and they are described below.

### Manual

Manual release treatments are used to reduce competition from unwanted vegetation, usually, but not limited to other tree species. The two main release treatments are precommercial thinning (PCT) and hardwood release.

### Precommercial Thinning

A common manual release treatment is PCT. This silviculture activity is used to manipulate the density, structure or species composition of overstocked young forest stands. Generally, the purpose of a PCT operation is to release the biggest and best growing trees so they can maintain their growth. PCT decisions are not made at the planting planning process, as planting density should not require PCT. This tool is used when ingrowth from natural regeneration, both conifer and hardwood, occurs reducing the growth and vigor of planted saplings. PCT is normally conducted in a stand between the ages of 10 and 20 years. Remaining density should be appropriate for the site, and range from 250-350 TPA. In areas of disease, such as SNC, PCT can be used to favor western hemlock and other resistant species over Douglas-fir to help ensure a healthy future stand.

### Hardwood Release

Hardwood release is used when ingrowth of hardwoods, mainly red alder in the northwest and madrone and tanoak in the southwest, threaten to change the

stand from conifer dominate to hardwood dominate. While hardwoods are important on the landscape and for local mills, long term conifer production is the goal for many stands across the planning area. In this treatment, hardwoods are removed leaving all conifer. This differs from a PCT in the fact that conifer spacing and species are not manipulated.

#### Chemical

Chemical release treatments involve the application of herbicides to control undesirable vegetation. Typical application methods are broadcast, directed spray, and hack and squirt, and are described below.

#### Broadcast

Broadcast application treatments are sprayed over the top of seedlings and undesirable vegetation. These applications usually occur in the first one to two years after planting and are designed to reduce competition from annual forbs and grasses. The two main application methods are aerial and backpack. Aerial is more efficient and cost effective than backpack, however, aerially application methods are a concern to certain resource users, wildlife professionals, and members of the public. Broadcast applications can also occur later in the stand as a release treatment from hardwoods that have overgrown the planted conifer, however, this is rarely used.

#### Directed Spray

Directed spray (spot spray) applications are made with a backpack and target individual plants. This treatment is often used to remove invasive species, such as Scotch broom, from young stands.

#### Hack N Squirt

Hack and squirt (basal or stem-injection) is typically applied as way to release conifers from hardwood competition. This method selects certain species, such as red alder, bigleaf maple, madrone, myrtle, tanoak and chinquapin for treatment.

#### Animal Damage Control

Animal damage on newly planted seedlings reduces their overall size, health, and vigor. Extensive damage can lead to interplanting, extend the time to achieve free to grow, potentially violating the Forest Practice Act. Animal damage occurs in many forms, but the most common is from ungulates (deer and elk) and mountain beaver.

#### Ungulates

Ungulate browse ranges from minor to severe. Minor browse damage usually has little impact to growth and survival. Repeated severe browse damage to seedlings, sometimes seen with western redcedar, can have major impacts on growth and occasionally lead to mortality. Control measures include Vexar tubing to protect from browse and controled hunts.

#### Mountain Beaver

Mountain beavers clip the seedling at its base, causing mortality. As the seedling ages, the diameter becomes too large and the animal climbs the stem and clips branches. Mountain beaver browse will occur in most stands in the norther part of the planning area with little damage in the southern part. Trapping is used to reduce damage from mountain beavers.

## Rip\_Code

This field name is used for the FPA and also the other versions of the model runs/Alts, so no need to change field name for different types of runs.

*Buffer distances are from the outer edge of the channel. Lines with 'BufClass' field and channels were generated by Terrainworks 4/24/2020, FMP total buffers generated by BRM 5/12/2020.*

'BufClass' from Terrainworks stream lines

Current FMP buffer widths will be applied on the Terrainworks channels

Code	Description
1	Yes Buffer
0	No Buffer

### Terrainworks BufClass field reclassified to current FMP strategies:

'BufClass'	No AA	Yes AA	Terrainworks "BufClass" Description
F_P	115	115	F perennial, 120' buffer
F_S	115	115	F seasonal, 50' buffer
N_P_LM	115	115	N perennial large and medium, 120' buffer
N_P_Temp	30	55	N small perennial temperature zone (500' upstream from F channel edge, if a confluence, or from end-of-fish, measured as slope distance), two scenarios: 50' and 120' buffers
N_P_S or N_P	30	55	N small perennial, upstream of N_F_Temp, 35' buffer
N_DF_Temp	30	55	N debris-flow, within 500' slope length upstream of any Type F, 50' and 120' buffers
N_DF	30	55	N debris-flow, upstream of N_DF_Tran, 35' buffer
N_HE_Temp	30	55	N high energy, within 500' slope length upstream of any Type F, 50' and 120' buffers
N_HE	30	55	N high energy, upstream of N_HE_Tran, 35' buffer
N_P_S_Tran	0	0	N perennial to seasonal transition zone (100' length), 35' buffer
N_S	0	0	N seasonal, no buffer

*\*Randy did the first edits, then Colleen and Jodi reviewed and made minor changes, 5/12/2020.*

Below is example from the Calibration FPA model run stream buffers

**Aquatic Zone** is based off size of stream only (each side of line), and will apply to all buffer strategies below

Size	Aquatic Zone
Large	20
Med	10
Small	5

**FPA Buffer size:**

Query	Code	Description	Buffer Ft	Added to outside of aquatic zone
Fish & Large	1	Fish	120	100
Fish & Med	1	Fish	80	70
Fish & Small	1	Fish	55	50
Perennial & Large	2	Non-fish	90	70
Perennial & Med	2	Non-fish	60	50
Perennial & Small	3	Perennial	5	0

**35??? Buffer Size:**

Query	Code	Description	Buffer Ft	outside of aquatic zone
Fish & Large		Fish	135	115
Fish & Med		Fish	125	115
Fish & Small		Fish	120	115
Large <> Fish		Non-fish	135	115
Med <> Fish		Non-fish	125	115
Fpsize = 'Small' AND Fishpres <> 'Fish' AND Duration in ( 'Perennial', 'PERENNIAL')		Perennial	40	35