

2020 CCO Incentive Measures

Updated 30 January 2020

#	Measure	NQF Number	Measure Description	Measure Steward	Data Source*	Population Characteristics		Sector				
						Patient Population	Sex	Dental Health	Behavioral Health	Primary Care	Specialty Physical Health**	Hospital
Prevention/Early Detection - Physical Health Conditions												
1	Childhood Immunization Status (Combo 2)	0038	Percentage of children that turned 2 years old during the measurement year and had the Dtap, IPV, MMR, Hib, HepB, and VZV vaccines by their second birthday	NCQA	Claims/Clinical Data	Children	All			Y		Y
2	*NEW* Immunizations for Adolescents (Combo 2)	1407	Percentage of adolescents that turned 13 years old during the measurement year and had the meningococcal, Tdap, and HPV vaccines by their 13th birthday	NCQA	Claims/Clinical Data	Adolescent	All			Y		Y
3	*NEW* Well-Child Visits in the 3rd, 4th, 5th, and 6th Years of Life (kindergarten readiness)	1516	Percentage of children ages 3 to 6 that had one or more well-child visits with a PCP during the measurement year	NCQA	Claims/Clinical Data	Children	All			Y		
4	Prenatal & Postpartum Care - Postpartum Care	1517	Percentage of deliveries of live births between November 6 of the year prior to the measurement year and November 5 of the measurement year that had a postpartum visit on or between 21 and 56 days after delivery	NCQA	Claims/Clinical Data	Adolescent, Adult, Older Adult	Female			Y		Y
Prevention/Early Detection - Mental Health Conditions												
5	Screening for Clinical Depression and Follow-Up Plan	0418	Percentage of patients aged 12 years and older screened for clinical depression on the date of the encounter using an age appropriate standardized depression screening tool AND if positive, a follow up plan is documented on the date of the positive screen	CMS	Claims/Clinical Data (eCQM measure)	Adolescent, Adult, Older Adult	All			Y		
Prevention/Early Detection - Substance Use Disorder (SUD) Conditions												
6	Cigarette Smoking Prevalence	NA	Percentage of Medicaid members ages 13 and older who currently smoke cigarettes or use other tobacco products	OHA	Clinical Data	Adult, Older Adult	All			Y		Y
7	Alcohol and Drug Misuse: Screening, Brief Intervention and Referral for Treatment (SBIRT)	NA	Percentage of patients ages 12 years and older who have had a qualifying outpatient visit or home visit during the measurement year with one or more screening, brief intervention, and referral to treatment services	OHA	Clinical Data	Adolescent, Adult, Older Adult	All			Y		
Prevention/Early Detection - Oral Health Conditions												
8	*NEW* Members Receiving Preventive Dental Services, ages 1-5 (kindergarten readiness) and 6-14	NA	Percentage of enrolled children ages 1-5 (kindergarten readiness) and 6-14 who received a preventive dental service during the measurement year	OHA	Claims	Children, Adolescent	All	Y				
9	Oral Evaluation for Adults with Diabetes	NA	Percentage of adults with diabetes who received at least one oral evaluation within the reporting year	OHA (modified from DQA/ NCQA)	Claims	Adult, Older Adult	Adults	Y				
Prevention/Early Detection - All Conditions												
10	Mental and Physical Health and Oral Health Assessment Within 60 Days for Children in DHS Custody	NA	Percentage of children ages 0-17 who received a physical health assessment, children ages 1-17 who received a dental health assessment, and children ages 4-17 who received a mental health assessment within 60 days of the state notifying CCOs that the children were placed into custody with the Department of Human Services (foster care)	OHA	Claims/Social Service Data	Children, Adolescent	All	Y	Y	Y		
Chronic Disease and Special Health Needs - Physical Health Conditions												
11	Comprehensive Diabetes Care: Hemoglobin A1c (HbA1c) Poor Control (>9.0%)	0059	Percentage of patients 18-75 years of age with diabetes who had hemoglobin A1c > 9.0% during the measurement period	NCQA	Claims/Clinical Data (eCQM measure)	Adult, Older Adult	All			Y		
Chronic Disease and Special Health Needs - Substance Use Disorder (SUD) Conditions												
12	*NEW* Initiation and Engagement of Alcohol and Other Drug Abuse or Dependence Treatment	0004	Percentage of adolescent and adult patients with a new episode of alcohol or other drug (AOD) dependence who (1) initiated treatment through an inpatient AOD admission, outpatient visit, intensive outpatient encounter or partial hospitalization within 14 days of the diagnosis and (2) who initiated treatment and who had two or more additional services with a diagnosis of AOD within 34 days of the initiation visit	NCQA	Claims	Adolescent, Adult, Older Adult	All		Y	Y		Y
Acute, Episodic and Procedural Care (Includes Maternity and Hospital)												
13	Disparity Measure: Emergency Department Utilization among Members with Mental Illness	NA	Number of ED visits per 1,000 member months for adult members enrolled within the organization who are identified as having experienced mental illness	Homegrown CCO	Claims	Adult, Older Adult	All		Y	Y		Y

Notes

*Clinical data includes electronic health records, registry data, and paper medical records. Claims/clinical data includes measures that require claims and clinical data, and measures that require claims or claims and clinical data. eCQM measures are indicated using italic font.

Measure Steward Abbreviations

AHRQ: Agency for Healthcare Research and Quality
 AMA-PCPI: American Medical Association-convened Physician Consortium for Performance Improvement
 CMS: Centers for Medicare & Medicaid Services
 DQA: Dental Quality Alliance
 MNCM: Minnesota Community Measurement

NCQA: National Committee for Quality Assurance
 OHA: Oregon Health Authority
 OHSU: Oregon Health & Science University
 POA: Pharmacy Quality Alliance
 TJC: The Joint Commission

Version Control

* This document was updated on 30 January 2020 to include a more accurate description of the specifications for the Oral evaluation for adults with diabetes measure.

* This document was updated on 10 October 2019 to correct an error in the description of the cigarette smoking cessation measure; the earlier version erroneously indicated that this measure is limited to adults. This measure includes Medicaid members ages 13 and older; the correction has been made above.