

SUICIDALITY, SELF-HARM AND PROBLEM GAMBLING

- Among individuals with problematic gambling behaviors, suicidal behaviors (e.g., suicide attempts, completed suicides) are more common among those who also engage in disordered substance use than those who engage in problem gambling without disordered substance use.¹³
- Suicidality is 5.3 times higher in individuals with problematic gambling than those without problematic gambling behaviors.¹⁴
- Even after controlling for conditions such as depression, substance use, ADHD, and anxiety, suicidality rates remained 2.9 times higher in those with problematic gambling than in the general population.¹⁴
- Although the research is limited, some data suggest a positive relationship - that is, when one increases so does the other - between non-suicidal self-harm (e.g., cutting, self-burning, or self-mutilation) and problem gambling.¹⁵



PERSONALITY DISORDERS AND PROBLEM GAMBLING

- Nearly one half (47.9%) of individuals seeking treatment for problem gambling had comorbid personality disorders, with the most prevalent being narcissistic (16.6%), antisocial (14.0%), avoidant (13.4%), obsessive-compulsive (13.4%), and borderline (13.1%) personality disorders.⁷
- In a large epidemiological study of over 10,000 adults, individuals with problematic gambling were significantly more likely to be diagnosed with a comorbid personality disorder (42.96%) than individuals who do not problem gamble (7.3%).¹⁶

7. Dowling, N., Cowlishaw, S., Jackson, A., Merkouris, S., Francis, K., & Christensen, D. (2015). Prevalence of psychiatric co-morbidity in treatment-seeking problem gamblers: a systematic review and meta-analysis. *Australian & New Zealand Journal of Psychiatry*, 49(6), 519-539

13. Håkansson, A. & Karlsson, A. (2020). Suicide attempts in patients with gambling disorder- associations with comorbidity in cluding substance use disorders. *Frontiers in Psychology*, 11, 593533.

14. Wardle, H., John, A., Dymond, S., McManus, S. (2020). Problem gambling and suicidality in England: secondary analysis of a representative cross-sectional survey. *Public Health*, 184, 11-16.

15. Gray, H. M., Edson, T. C., Nelson, S. E., Grossman, A. B., & LaPlante, D. A. (2020). Association between gambling and self-harm: a scoping review. *Addiction Research & Theory*, 1-13.

16. Pietrzak, R. H., Morasco, B. J., Blanco, C., Grant, B. F., & Petry, N. M. (2007). Gambling level and psychiatric and medical disorders in older adults: Results from the National Epidemiologic Survey on alcohol and related conditions. *American Journal of Geriatric Psychiatry*, 15(4), 301-313.

ANXIETY DISORDERS AND PROBLEM GAMBLING

An estimated 17.7% of individuals seeking treatment for problem gambling have an anxiety disorder.⁷ Of those engaged in problematic gambling, individuals are twice as likely to experience any anxiety condition (i.e., PTSD, panic disorder, phobia, generalized anxiety) than the general population.¹

GENERALIZED ANXIETY DISORDER



- Approximately 14.4% of individuals seeking treatment for problem gambling also have a generalized anxiety disorder.⁷
- Generalized anxiety has been shown to predict problem gambling 1 year later.¹⁰

POST-TRAUMATIC STRESS DISORDER



- A recent review found that across studies, PTSD symptom severity heavily contributed to gambling disorder severity.¹⁷
- Approximately 12.3% of individuals who sought treatment for problem gambling met the criteria for post-traumatic stress disorder.⁷
- People who engage in problematic gambling with a history of PTSD are more likely to use gambling to cope with unwanted emotions.¹⁸

OTHER ANXIETY-RELATED DISORDERS



- Roughly 13.7% of individuals who sought treatment for problem gambling also had a diagnosis of panic disorder.⁷
- 14.9% of individuals who sought treatment for problematic gambling also had a social phobia.⁷
- An estimated 8.2% of individuals seeking treatment for problematic levels of gambling also have obsessive-compulsive disorder.⁷

Women who experienced any anxiety condition before gambling onset, *tripled* their risk of developing problematic gambling behaviors.¹

1. Sundqvist, K. & Rosendahl, I. (2019). Problem gambling and psychiatric comorbidity - risk and temporal sequencing among women and men: Results from the Swelogs case-control study. *Journal of Gambling Studies*, 35, 757-771.

7. Dowling, N., Cowlishaw, S., Jackson, A., Merkouris, S., Francis, K., & Christensen, D. (2015). Prevalence of psychiatric co-morbidity in treatment-seeking problem gamblers: a systematic review and meta-analysis. *Australian & New Zealand Journal of Psychiatry*, 49(6), 519-539.

10. Dowling, N. A., Butera, C. A., Merkouris, S. S., Youssef, G. J., Rodda, S. N., Jackson, A. C. (2019). The reciprocal association between problem gambling and mental health symptoms/substance use: cross-lagged path modelling of longitudinal cohort data. *Journal of Clinical Medicine*, 8, 1888.

17. Moore III, L. H., & Grubbs, J. B. (2021). Gambling disorder and comorbid PTSD: a systematic review of empirical research. *Addictive Behaviors*, 114.

18. Ledgerwood, D. M., & Milosevic, A. (2015). Clinical and personality characteristics associated with post traumatic stress disorder in problem and pathological gamblers recruited from the community. *Journal of Gambling Studies*, 31, 501-5012.

ATTENTION-DEFICIT/HYPERACTIVITY DISORDER (ADHD) AND PROBLEM GAMBLING

- One in five individuals with problem gambling having clinical levels of ADHD symptoms.¹⁹
- Individuals who engage in problem gambling are 4.18 times more likely to have ADHD than those who do not engage in problem gambling. Individuals with ADHD were 2.85 times more likely to experience problem gambling than individuals without ADHD.¹⁹
- 18.46% of individuals who engage in problem gambling also have ADHD and 11.75% of individuals with ADHD also engage in problem gambling.¹⁹
- Among males with ADHD, impulsivity traits (1) lack of premeditation and (2) sensation seeking are not significant risk factors for problem gambling.²⁰

Emotion dysregulation in people with ADHD contributes to problem gambling²⁰

OTHER HARMFUL BEHAVIORS AND PROBLEM GAMBLING

Individuals who engage in problem gambling tend to also engage in more harmful risk taking and impulsive behaviors compared to the general population²³.

- Problem gambling is significantly related to intimate personal violence (IPV), with one-third of individuals with problem gambling reporting being a survivor of IPV or perpetrators of IPV.³
- Individuals who engage in problem gambling are more likely to lose control of their anger and engage in aggressive behaviors than individuals who do not problem gamble.²¹
- There is an increased risk for lying, cheating, and stealing with those who engage in problem gambling compared to those who do not.²²

3. Dowling NA, Jackson AC, Suomi A, et al. (2014b) Problem gambling and family violence: Prevalence and patterns in help-seeking populations. *Addictive Behaviors* 39: 1713–1717.
 19. Theule, J., Hurl, K. E., Cheung, K., Ward, M., Henrikson, B. (2019). Exploring the relationships between problem gambling and ADHD: a meta-analysis. *Journal of Attention Disorders*, 23(12), 1427-1437.
 20. Brunault, P., Mathieu, S., Faussat, C., Barrault, S., Varescon, I. (2020). Impulsivity facets and cognitive distortions associated with problem gambling: differences between ADHD and non-ADHD gamblers. *Europeans Review of Applied Psychology*, 70(4), 100559
 21. Aymami, N., Granero, R., Penelo, E., Fernández-Aranda, F., Krug, I., Gunnard, K., Santamaría, J. J., Bueno, B., Jaurieta, N., Gómez-Peña, M., Álvarez-Moya, E., Claes, L., Fagundo, A. B., Menchón, J. M., & Jiménez-Murcia, S. (2014). Anger in pathological gambling: clinical, psychopathological, and personality correlates. *The Spanish Journal of Psychology*, 17. <https://doi.org/10.1017/sjp.2014.40>
 22. Lind, K., & Kääriäinen, J. (2018). Cheating and stealing to finance gambling: analysis of screening data from a problem gambling self-help program. *Journal of Gambling Issues*, 39. <https://doi.org/10.4309/jgi.2018.39.9>
 23. Wong, G., Zane, N., Saw, A., & Chan, A. K. K. (2013). Examining gender differences for gambling engagement and gambling problems among emerging adults. *Journal of Gambling Studies / Co-Sponsored by the National Council on Problem Gambling and Institute for the Study of Gambling and Commercial Gaming*, 29(2), 171–189. <https://doi.org/10.1007/s10899-012-9305-1>

PROBLEM GAMBLING PREVENTION AND INTERVENTION STRATEGIES

Given the harmful relationship problem gambling has with mental health, prevention strategies and methods to intervene are important for community stakeholders to know about.

- Learn more about the harmful relationship between mental health and problematic gambling, including prevention, treatment, and ways to support individuals who are seeking help.
- The sequence of comorbidity onset for mental health symptoms and problematic gambling differ between sexes. Females tend to begin gambling after the onset of depression, anxiety, and/or substance use problems; conversely, males tend to initiate gambling behaviors prior to mental health symptom onset.¹
- Integrate the harmful impact of problem gambling into education opportunities when discussing mental health concerns and/or developing partnerships.

- Use a brief assessment measure to identify gambling problems such as the Brief Biosocial Gambling Screen (BBGS) and assess functionality in areas of life (e.g., occupational, financial, social, interpersonal, and intrapersonal).^{24,25}
- Develop a harm minimization strategy. Take a responsible gambling approach and collaboratively plan what best aligns with their goals, while also promoting health and reducing harm.²⁴

Interventions and Treatment

- Brief, early intervention (e.g. motivational interviewing or motivational enhancement)
- Social support groups such as Gambling Anonymous (GA) or SMART Recovery
- Cognitive-behavioral therapy from specialists in addictive behaviors
- Improve problem gambling behaviors by integrating gambling screening and referral processes into substance abuse and mental health treatment.^{12,26}
- Gambling-specific treatments may be most beneficial for treatment success of problem gambling behaviors, including in a substance use treatment setting.²⁶

1. Sundqvist, K. & Rosendahl, I. (2019). Problem gambling and psychiatric comorbidity - risk and temporal sequencing among women and men: Results from the Swelogs case-control study. *Journal of Gambling Studies*, 35, 757-771.

12. Kennedy, S. H., Welsh, B. R., Fulton, K., Soczynska, J. K., McIntyre, R. S., O'Donovan, C., Milev, R., le Melleo, J., Bissler, J., Zimmerman, M., Martin, N. (2010). Frequency and correlates of gambling problems in outpatients with major depressive disorder and bipolar disorder. *Canadian Journal of Psychiatry*, 55(9), 568-576.

24. Price, A., Hilbrecht, M., & Billi, R. (2021). Charting a path towards a public health approach for gambling harm prevention. *Journal of Public Health*, 1-17. <https://doi.org/10.1007/s10389-020-01437-2>

25. Yao, Y. H. C., & Potenza, M. N. (2015). Gambling disorder and other behavioral addictions: Recognition and treatment. *Harvard Review of Psychiatry*, 23(2), 134. <https://doi.org/10.1097/HRP.0000000000000051>

26. Rash, C. J., Weinstock, J., & Van Patten, R. (2016). A review of gambling disorder and substance use disorders. *Substance abuse and rehabilitation*, 7, 3.

PROBLEM GAMBLING AND MENTAL HEALTH: RESEARCH CONTRADICTING THE LARGER BODY OF WORK

Some research suggests that the relationship between problem gambling and mental health is not consistent.

- Among adults over the age of 40, severity of specific phobias were negatively associated with gambling activity. In fact, that the more severe the phobia, the less often individuals engaged in gambling.²⁷
- Among young adults (adults under 40), greater use of substances, and excessive computer use were associated with less gambling.²⁷

Key Findings

- Those who engage in problematic gambling and also have a comorbid mental health condition, specifically PTSD or depression, are more likely to use gambling as a coping strategy.
- Individuals at-risk of and who engage in problematic gambling are at an increased risk of suicide and self-harm behaviors.
- Gambling behaviors are strongly associated with mental health conditions. Studies consistently report that there are high comorbidities between individuals who engage in problematic gambling and mental health conditions.
- For women, there is a greater risk of psychiatric and problematic gambling comorbidities.¹

1. Sundqvist, K. & Rosendahl, I. (2019). Problem gambling and psychiatric comorbidity - risk and temporal sequencing among women and men: Results from the Swelogs case-control study. *Journal of Gambling Studies*, 35, 757-771.

27. Black, D. W., Allen, J., Bormann, N. L. (2021). Are comorbid disorders associated with changes in gambling activity? A longitudinal study of younger and older subjects with DSM-IV pathological gambling. *Journal of Gambling Studies*.