

Oregon Behavioral Risk Factor Surveillance System (BRFSS), 1997

At Risk for Overweight/BMI
(Weighted Column Percents)

| | | Total | Age Groups | | | | | |
|--|---------|-------|------------|-------|-------|-------|-------|------|
| | | | 18-24 | 25-34 | 35-44 | 45-54 | 55-64 | 65+ |
| No risk | | 65.9 | 80.1 | 70.0 | 63.4 | 57.6 | 58.8 | 68.4 |
| At risk | | 31.7 | 18.3 | 27.3 | 33.4 | 39.6 | 39.6 | 29.7 |
| Unknown | | 2.4 | 1.5 | 2.7 | 3.2 | 2.8 | 1.7 | 1.9 |
| Male | No risk | 67.3 | 80.9 | 69.8 | 64.3 | 58.7 | 60.7 | 73.1 |
| | At risk | 32.1 | 18.3 | 28.8 | 35.7 | 41.0 | 38.8 | 26.2 |
| | Unknown | .6 | .9 | 1.4 | | .3 | .4 | .7 |
| Female | No risk | 64.6 | 79.4 | 70.3 | 62.5 | 56.5 | 56.9 | 65.1 |
| | At risk | 31.3 | 18.4 | 25.8 | 31.1 | 38.1 | 40.2 | 32.2 |
| | Unknown | 4.2 | 2.3 | 3.9 | 6.4 | 5.4 | 2.9 | 2.7 |
| Denominator is: All survey respondents | | | | | | | | |

At Risk for Overweight/BMI
(Unweighted Frequencies)

| | | Total | Age Groups | | | | | |
|--|---------|-------|------------|-------|-------|-------|-------|-----|
| | | | 18-24 | 25-34 | 35-44 | 45-54 | 55-64 | 65+ |
| Total | | 3308 | 292 | 607 | 750 | 592 | 391 | 676 |
| No risk | | 2206 | 235 | 437 | 480 | 354 | 232 | 468 |
| At risk | | 1022 | 52 | 155 | 247 | 221 | 152 | 195 |
| Unknown | | 80 | 5 | 15 | 23 | 17 | 7 | 13 |
| Male | Total | 1399 | 153 | 273 | 327 | 241 | 170 | 235 |
| | No risk | 955 | 124 | 198 | 209 | 146 | 101 | 177 |
| | At risk | 435 | 28 | 72 | 118 | 94 | 67 | 56 |
| | Unknown | 9 | 1 | 3 | | 1 | 2 | 2 |
| Female | Total | 1909 | 139 | 334 | 423 | 351 | 221 | 441 |
| | No risk | 1251 | 111 | 239 | 271 | 208 | 131 | 291 |
| | At risk | 587 | 24 | 83 | 129 | 127 | 85 | 139 |
| | Unknown | 71 | 4 | 12 | 23 | 16 | 5 | 11 |
| Denominator is: All survey respondents | | | | | | | | |

Oregon Behavioral Risk Factor Surveillance System (BRFSS), 1997

Obese: BMI greater than 27.8 (for Males) or 27.3 (for Females)
(Weighted Column Percents)

| | | Total | Age Groups | | | | | |
|----------------|----------------|-------|------------|-------|-------|-------|-------|------|
| | | | 18-24 | 25-34 | 35-44 | 45-54 | 55-64 | 65+ |
| No risk | | 65.9 | 80.1 | 70.0 | 63.4 | 57.6 | 58.8 | 68.4 |
| At risk | | 31.7 | 18.3 | 27.3 | 33.4 | 39.6 | 39.6 | 29.7 |
| Unknown | | 2.4 | 1.5 | 2.7 | 3.2 | 2.8 | 1.7 | 1.9 |
| Male | No risk | 67.3 | 80.9 | 69.8 | 64.3 | 58.7 | 60.7 | 73.1 |
| | At risk | 32.1 | 18.3 | 28.8 | 35.7 | 41.0 | 38.8 | 26.2 |
| | Unknown | .6 | .9 | 1.4 | | .3 | .4 | .7 |
| Female | No risk | 64.6 | 79.4 | 70.3 | 62.5 | 56.5 | 56.9 | 65.1 |
| | At risk | 31.3 | 18.4 | 25.8 | 31.1 | 38.1 | 40.2 | 32.2 |
| | Unknown | 4.2 | 2.3 | 3.9 | 6.4 | 5.4 | 2.9 | 2.7 |

Denominator is: All survey respondents

Obese: BMI greater than 27.8 (for Males) or 27.3 (for Females)

(Unweighted Frequencies)

| | | Total | Age Groups | | | | | |
|----------------|----------------|-------|------------|-------|-------|-------|-------|-----|
| | | | 18-24 | 25-34 | 35-44 | 45-54 | 55-64 | 65+ |
| Total | | 3308 | 292 | 607 | 750 | 592 | 391 | 676 |
| No risk | | 2206 | 235 | 437 | 480 | 354 | 232 | 468 |
| At risk | | 1022 | 52 | 155 | 247 | 221 | 152 | 195 |
| Unknown | | 80 | 5 | 15 | 23 | 17 | 7 | 13 |
| Male | Total | 1399 | 153 | 273 | 327 | 241 | 170 | 235 |
| | No risk | 955 | 124 | 198 | 209 | 146 | 101 | 177 |
| | At risk | 435 | 28 | 72 | 118 | 94 | 67 | 56 |
| | Unknown | 9 | 1 | 3 | | 1 | 2 | 2 |
| Female | Total | 1909 | 139 | 334 | 423 | 351 | 221 | 441 |
| | No risk | 1251 | 111 | 239 | 271 | 208 | 131 | 291 |
| | At risk | 587 | 24 | 83 | 129 | 127 | 85 | 139 |
| | Unknown | 71 | 4 | 12 | 23 | 16 | 5 | 11 |

Denominator is: All survey respondents