

Oregon 2016 BRFSS  
Immunization

**During the past 12 months, have you had either a seasonal flu shot or seasonal flu vaccine that was sprayed in your nose?**

(Weighted Column Percents)

		Age Groups						
		Total	18-24	25-34	35-44	45-54	55-64	65+
Yes		35.6	23.9	28.9	30.5	31.3	38.1	51.2
No		64.4	76.1	71.1	69.5	68.7	61.9	48.8
Male	Yes	32.2	22.3	23.7	25.4	28.2	36.9	49.5
	No	67.8	77.7	76.3	74.6	71.8	63.1	50.5
Female	Yes	38.9	25.7	34.7	35.4	34.5	39.4	52.7
	No	61.1	74.3	65.3	64.6	65.5	60.6	47.3

Denominator is: All survey respondents.

**Risk factor for respondents aged 65+ who have had a flu shot within the past 12 months.**

(Weighted Column Percents)

Flu shot in past 12 months		51.2
No flu shot in past 12 months		48.8
Male	Flu shot in past 12 months	49.5
	No flu shot in past 12 months	50.5
Female	Flu shot in past 12 months	52.7
	No flu shot in past 12 months	47.3

Denominator is: All survey respondents aged 65 or older who have had a flu shot in the past 12 months.

**A pneumonia shot or pneumococcal vaccine is usually given only once or twice in a person's lifetime and is different from the flu shot. Have you ever had a pneumonia shot?**

(Weighted Column Percents)

		Age Groups						
		Total	18-24	25-34	35-44	45-54	55-64	65+
Yes		41.3	45.4	26.0	20.2	22.8	35.7	76.6
No		58.7	54.6	74.0	79.8	77.2	64.3	23.4
Male	Yes	40.8	49.4	29.8	22.4	24.2	34.0	70.8
	No	59.2	50.6	70.2	77.6	75.8	66.0	29.2
Female	Yes	41.8	41.1	22.3	18.3	21.6	37.3	81.4
	No	58.2	58.9	77.7	81.7	78.4	62.7	18.6

Denominator is: All survey respondents.

Oregon 2016 BRFSS  
Immunization

**Risk factor for respondents aged 65+ who  
have ever had a pneumonia shot.**

(Weighted Column Percents)

Had pneumococcal shot		76.6
No pneumococcal shot		23.4
Male	Had pneumococcal shot	70.8
	No pneumococcal shot	29.2
Female	Had pneumococcal shot	81.4
	No pneumococcal shot	18.6

Denominator is: All survey respondents aged 65 or older who have ever had a pneumonia shot.