Oregon 2002 - 2005	BRFSS Data -	Behavior.	Risk Survey	Topics l	by County

Data are adjusted to the 2000 U.S.	Census Population.		
County	Asthma		
OREGON	9.3%		
Baker	9.6%		
Benton	8.7%		
Clackamas	9.6%		
Clatsop	9.5%		
Columbia	9.7%		
Coos	8.4%		
Crook	9.2%		
Curry	7.8%		
Deschutes	8.9%		
Douglas	11.8%		
Grant	10.3%		
Harney	5.5%		
Hood River	3.6%**		
Jackson	8.3%		
Jefferson	5.8%		
Josephine	11.2%		
Klamath	10.7%		
Lake	13.6%		
Lane	10.5%		
Lincoln	9.1%		
Linn	12.0%**		
Malheur	5.7%		
Marion	8.9%		
Morrow	10.5%		
Multnomah	8.9%		
Polk	6.9%		
Tillamook	9.4%		
Umatilla	7.1%		
Union	11.0%		
Wallowa	6.9%		
Washington	9.1%		
Yamhill	10.7%		
Gilliam/Wheeler	2.7%†		
Sherman/Wasco	9.8%		

Age-adjusted Prevalence of Selected Chronic Diseases and Related Risk Factors Among Adults by County, Oregon 2002-2005

Tables were created using combined BRFSS survey results from 2002 through 2005. Two numbers are provided for each condition or disease: an unadjusted rate and an age-adjusted rate.

Unadjusted rates, provided in separate tables, give a description of the absolute burden of a disease or risk factor for an individual county.

Age-adjusted rates supplied in this table allow you to compare rates for a county to the State's rate. They adjust for differences that could result from a county's population being, on average, older or younger than the State's. When comparing counties with Oregon or with each other, use these age-adjusted rates.

† Rate based on less than 50 respondents overall or fewer than 12 persons in one age group; may not accurately reflect behavior of entire county and results should be interpreted with caution. Estimates based on 10 or fewer respondents are not reported.

** Statistically significant difference compared to Oregon.

Three age groups were used for age adjusting: 18-34yrs., 35-54yrs, and 55+yrs.

Source: Oregon Behavioral Risk Factor Surveillance System - healthoregon.org/brfss