

Table III: Age-Adjusted and Unadjusted Rates for Preventive Health Screening among Adults, by County, Oregon 2004-2007

County	% who had their cholesterol checked within past 5 years (≥ 18 years old)		% who had a mammogram within past 2 years (women ≥40 years old)		% who had a PAP test within past 3 years (women ≥ 18 years old)	
	unadjusted	age-adjusted	unadjusted	age-adjusted	unadjusted	age-adjusted
OREGON		69.4%		73.3%		84.4%
Baker	68.9%	64.1%	74.0%	73.6%	73.2%†	71.5%†
Benton	64.5%	70.3%	79.2%	78.4%	91.5%	92.3%**
Clackamas	73.4%	71.4%	75.4%	74.2%	87.0%	86.4%
Clatsop	64.9%	60.9%**	64.0%	64.5%	70.9%	70.4%**
Columbia	80.6%	79.3%**	73.2%	74.0%	85.4%	84.2%
Coos	72.7%	68.2%	75.1%	72.2%	83.5%	84.4%
Crook	83.9%	82.8%**	69.6%†	68.4%†	91.1%†	91.0%†
Curry	79.3%	72.5%	80.8%†	78.2%†	82.1%†	87.3%†
Deschutes	74.7%	72.2%	68.7%	67.1%	81.1%	81.1%
Douglas	75.6%	71.5%	68.5%	66.8%**	78.8%	79.8%
Grant	62.4%	52.8%**	55.9%†	54.4%†	---	---
Harney	62.7%	61.9%	---	---	---	---
Hood River	53.4%	52.2%**	80.2%†	80.9%†	89.1%†	86.3%†
Jackson	71.4%	68.1%	72.3%	70.4%	84.1%	84.0%
Jefferson	71.7%	69.3%	59.2%†	57.0%†	79.2%†	79.1%†
Josephine	67.8%	61.0%**	70.6%	69.7%	79.7%	79.8%
Klamath	69.8%	67.6%	76.4%	73.1%	88.7%	88.5%
Lake	75.4%	72.5%	89.9%†	90.3%†	78.4%	81.0%
Lane	68.6%	68.3%	75.0%	73.5%	87.2%	87.1%
Lincoln	76.8%	69.3%	67.8%	66.1%	75.3%	75.4%
Linn	73.8%	71.7%	73.6%	72.1%	81.2%	81.2%
Malheur	66.9%	67.1%	66.2%	64.7%	78.8%	75.9%
Marion	66.0%	66.5%	77.3%	75.7%	79.6%	79.6%
Morrow	59.0%	59.4%	---	---	---	---
Multnomah	69.4%	70.6%	77.4%	76.7%	86.6%	86.0%
Polk	75.1%	74.1%	77.7%	72.7%	79.3%	78.2%
Tillamook	74.2%	67.2%	62.6%	60.9%	85.4%†	86.1%†
Umatilla	67.8%	67.7%	72.3%	70.8%	85.3%	84.9%
Union	60.5%	59.4%	74.8%	72.7%	90.3%	89.1%
Wallowa	73.1%	60.0%	62.0%†	60.5%†	---	---
Washington	70.9%	72.0%	77.1%	76.5%	87.8%	87.1%
Yamhill	65.4%	66.0%	70.2%	68.6%	82.7%	80.8%
Gilliam/Wheeler	82.3%	63.7%†	---	---	---	---
Sherman/Wasco	57.3%	51.4%**	75.9%	77.0%	78.0%†	79.1%†

* Unadjusted rates provide a description of the absolute burden of disease or risk factor for a county. Age-adjusted rates allow you to compare rates for a given county to the State's rate. When comparing counties with Oregon or with each other, it's best to use age-adjusted rates.

** Statistically significant difference compared to Oregon.

† % based on less than 50 respondents; may not accurately reflect behavior of entire county.

Sources: BRFSS 2004-2007

Table III (continued): Unadjusted Rates for Preventive Health Screening among Adults, by County, Oregon 2004-2007

County	% who had a blood stool test within the past year (≥50 years old)	% who had a sigmoidoscopy or colonoscopy within 5 years (≥50 years old)
	unadjusted	unadjusted
OREGON	18.9%	47.0%
Baker	21.7%	24.8%
Benton	20.9%	60.3%
Clackamas	14.6%	46.0%
Clatsop	15.3%	45.1%
Columbia	23.1%	36.5%
Coos	26.3%	46.4%
Crook	10.0%	31.4%
Curry	26.2%	45.1%
Deschutes	16.6%	51.7%
Douglas	26.0%	46.1%
Grant	4.6% †	48.4% †
Harney	---	---
Hood River	16.9% †	28.3% †
Jackson	18.2%	45.2%
Jefferson	14.6% †	30.4% †
Josephine	8.8%	42.5%
Klamath	24.1%	42.4%
Lake	11.3% †	25.6% †
Lane	24.4%	53.4%
Lincoln	11.4%	32.3%
Linn	28.6%	43.6%
Malheur	4.7%	32.3%
Marion	23.1%	44.5%
Morrow	---	---
Multnomah	15.6%	47.5%
Polk	26.5%	50.1%
Tillamook	18.6%	44.6%
Umatilla	18.9%	40.6%
Union	9.2% †	47.0% †
Wallowa	13.7% †	34.6% †
Washington	19.5%	55.9%
Yamhill	14.1%	47.0%
Gilliam/Wheeler	---	---
Sherman/Wasco	20.5%	38.7%

* Unadjusted rates provide a description of the absolute burden of disease or risk factor for a county. Age-adjusted rates allow you to compare rates for a given county to the State's rate. When comparing counties with Oregon or with each other, it's best to use age-adjusted rates.

** Statistically significant difference compared to Oregon.

† % based on less than 50 respondents; may not accurately reflect behavior of entire county.

Sources: BRFSS 2004-2007