Oregon 2004 -2007 BRFSS Data - Behavior Risk Survey Topics by County

Binge Drinking, Adult Females, Oregon, 2004-2007 (Non-age adjusted)

Binge drinking among adult females: Had 4+ drinks of alcohol on one occasion in the past 30 days.\*\*\*

Source: Oregon Behavioral Risk Factor Surveillance System

		95% Co	nfidence			
Females	PERCENT	Inte	rval	Small	Number	Interviews
18 years & older	weighted %	L. limit %	U. limit %	#	unw eighted N	unweighted N
STATEWIDE	8.3	7.7	8.9		1094	16442
BAKER	6.6	3.1	13.5		8	136
BENTON	9.2	5.6	14.7		22	396
CLACKAMAS	6.7	5.3	8.5		94	1603
CLATSOP	7.1	3.8	13.1		13	208
COLUMBIA	7.0	4.0	11.9		15	235
COOS	10.5	6.9	15.8		27	385
CROOK	13.6	5.2	31.0		8	103
CURRY	9.6	5.0	17.7		10	130
DESCHUTES	11.4	8.4	15.4		57	633
DOUGLAS	6.9	4.8	9.9		37	585
GILLIAM/WHEELER	**	**	**	**	2	20
GRANT	22.8	10.5	42.7		9	52
HARNEY	9.6	3.3	24.6	**	4	40
HOOD RIVER	3.5	0.8	13.2		2	102
JACKSON	7.5	5.6	9.9		57	942
JEFFERSON	5.8	1.8	16.8		4	101
JOSEPHINE	7.6	4.5	12.6		21	421
KLAMATH	9.5	5.9	15.1		27	334
LAKE	9.5	3.7	22.5		5	57
LANE	8.3	6.5	10.5		97	1633
LINCOLN	4.3	2.0	9.1		9	239
LINN	8.2	5.4	12.3		36	542
MALHEUR	4.1	1.6	9.8		7	172
MARION	7.6	5.9	9.9		73	1186
MORROW	12.8	5.6	26.6		7	55
MULTNOMAH	11.3	9.8	12.9		251	2717
POLK	5.1	3.1	8.5		18	348
TILLAMOOK	13.6	7.5	23.3		12	141
UMATILLA	7.2	3.9	13.0		20	348
UNION	6.0	2.9	12.0		10	162
WALLOWA	14.0	5.7	30.5		6	52
WASCO/SHERMAN	3.5	1.5	7.9		6	145
WASHINGTON	6.2	4.9	7.8		100	1838
YAMHILL	6.7	4.0	10.8		20	381

<sup>\*\* %</sup> based on less than 50 respondents may not accurately reflect behavior of the entire county. Data are not reported for counties with less than 10 respondents.

<sup>\*\*\*</sup>In 2005, the definition of binge drinking for women changed from 5+ drinks of alcohol to 4+ drinks of alcohol on one occasion. Prior to 2005, the binge drinking definition was 5+ drinks of alcohol on one occasion for both genders. The 2004 binge drinking data, based upon 5+ drinks of alcohol on one occasion for both genders, has been combined with 2005-2007 data, so we can include 2004 data, even though the drinks on one occasion changed for women in

I	_	_	
-	Λ	$\boldsymbol{\epsilon}$	
		_	

Source: Oregon Behavioral Risk Factor Surveillance System - healthoregon.org/brfss