

Oregon Adults 65+ Who Received Any Type of Seasonal Flu Immunization Within the Past Year, Oregon, 2012-2015

Adults 18 years & older	<i>Unadjusted</i>		
	%	95% Confidence Interval ¹	
		L. limit %	U. limit %
STATEWIDE	55.3	53.9	56.8
BAKER	30.4 ~	18.5	45.6
BENTON	51.3	43.1	59.4
CLACKAMAS	59.9	55.2	64.4
CLATSOP	53.0	40.4	65.2
COLUMBIA	65.7	54.1	75.7
COOS	52.5	43.7	61.1
CROOK	67.9	54.9	78.6
CURRY	48.4	35.6	61.4
DESCHUTES	59.0	52.3	65.5
DOUGLAS	53.4	46.4	60.4
GRANT	48.5	30.2	67.1
HARNEY	*	*	*
HOOD RIVER	51.1	33.2	68.7
JACKSON	47.9	42.0	53.9
JEFFERSON	61.2	40.6	78.4
JOSEPHINE	48.1	40.7	55.6
KLAMATH	49.6	39.8	59.4
LAKE	*	*	*
LANE	55.5	51.1	59.8
LINCOLN	56.0	46.9	64.7
LINN	53.2	46.4	60.0
MALHEUR	37.7 ~	24.4	53.1
MARION	57.4	51.9	62.8
MORROW	54.4	31.5	75.7
MULTNOMAH	59.9	55.8	63.9
POLK	49.6	40.1	59.2
TILLAMOOK	56.8	42.9	69.7
UMATILLA	62.4	51.6	72.2
UNION	38.2	24.6	54.0
WALLOWA	58.2	33.1	79.7
WASHINGTON	58.1	53.2	62.9
WHEELER	*	*	*
YAMHILL	58.7	49.9	67.0
GILLIAM/WASCO/SHERMAN	54.2	38.8	68.7

* This number is suppressed because it is statistically unreliable.

~ Statistically Significant: significantly higher or lower than the statewide rate.

¹ Confidence intervals provide a measure of how much an estimate might vary due to chance. We present 95% confidence intervals, which means that if we conducted this survey over and over, 95% of the confidence intervals calculated would contain the true population value. Wider intervals suggest the estimate is more unreliable and should be interpreted with caution, however it is the best estimate available.

Source: Oregon Behavioral Risk Factors Surveillance System 2012-2015 county combined
 NOTE: Age adjustment is based on three age groups: 18-34; 35-54; and 55+, per U.S. Standard Million Population.