

TABLE 3-5. Contraceptive use, number of previous abortions, and number of living children by age of patient, Oregon occurrence, 2016

Contraceptive used, previous abortions, and number of living children	Total	Age groups								
		<15	15-19	20-24	25-29	30-34	35-39	40-44	45+	N.S.
Total	8,942	20	865	2,548	2,478	1,659	997	341	34	–
None used	5,421	13	596	1,543	1,478	959	597	215	20	–
No previous abortion	3,362	13	525	1,102	830	472	305	106	9	–
One	1,147	–	51	290	352	237	147	63	7	–
Two	499	–	8	80	168	139	76	26	2	–
Three	195	–	3	29	65	52	34	12	–	–
Four or more	130	–	–	20	38	41	24	6	1	–
Pills used	1,036	3	91	351	311	175	86	18	1	–
No previous abortion	631	3	80	236	169	89	46	7	1	–
One	237	–	8	71	85	49	17	7	–	–
Two	101	–	2	30	34	23	10	2	–	–
Three	32	–	–	5	12	7	6	2	–	–
Four or more	22	–	–	4	7	5	6	–	–	–
Condoms used	1,227	1	83	336	336	241	162	59	9	–
No previous abortion	732	1	73	241	192	111	73	34	7	–
One	297	–	8	63	86	74	50	14	2	–
Two	112	–	1	20	35	30	18	8	–	–
Three	39	–	–	5	12	10	10	2	–	–
Four or more	28	–	–	1	7	11	8	1	–	–
Other contraceptive	913	3	57	221	257	217	118	37	3	–
No previous abortion	520	3	50	152	144	104	46	20	1	–
One	242	–	6	52	63	71	35	13	2	–
Two	84	–	–	10	29	21	21	3	–	–
Three	38	–	–	4	10	13	11	–	–	–
Four or more	19	–	–	1	5	8	4	1	–	–
Contraceptive use unknown ..	459	–	47	131	121	91	51	17	1	–
No previous abortion	292	–	43	95	75	45	24	9	1	–
One	92	–	1	23	24	30	10	4	–	–
Two	35	–	1	8	12	4	9	1	–	–
Three	13	–	–	1	3	3	5	1	–	–
Four or more	12	–	–	1	3	4	2	2	–	–
Number of living children										
No children¹	4,253	20	738	1,577	1,113	499	236	65	5	–
Total with children	4,673	–	125	966	1,361	1,157	760	275	29	–
One	1,999	–	109	637	587	371	228	63	4	–
Two	1,626	–	13	254	496	453	284	113	13	–
Three	697	–	1	68	197	219	143	61	8	–
Four	248	–	1	7	63	85	67	22	3	–
Five or more	103	–	1	–	18	29	38	16	1	–

– Quantity is zero.

¹ Rows will not add to total due to some patients having an unknown number of children.

N.S. = Not stated.

NOTE: Contraceptive totals include abortions where the number of previous abortions is unknown. Multiple contraceptive methods may be reported for a single patient.